

# High piecewise syndeticity of product sets in amenable groups

Isaac Goldbring

University of Illinois at Chicago

Combinatorics meets ergodic theory  
Banff, July 2015

# 1 A quantitative version of Jin's Theorem

## 2 Amenable groups

## 3 Proof of the first part

# Densities

## Definition

Suppose that  $A \subseteq \mathbb{Z}$ .

- 1 The *upper density* of  $A$  is

$$\bar{d}(A) := \limsup_{n \rightarrow \infty} \frac{|A \cap [-n, n]|}{2n + 1}.$$

- 2 The *Banach density* of  $A$  is

$$\text{BD}(A) := \lim_{n \rightarrow \infty} \sup_{x \in \mathbb{Z}} \frac{|A \cap [x - n, x + n]|}{2n + 1}.$$

# Jin's Theorem

## Theorem (Jin, 2002)

If  $A, B \subseteq \mathbb{Z}$  are such that  $BD(A), BD(B) > 0$ , then  $A + B$  is *piecewise syndetic*: there is  $m \in \mathbb{N}$  such that  $A + B + [-m, m]$  contains intervals of arbitrarily large length.

The proof used nonstandard analysis.

# A quantitative version of Jin's Theorem

## Theorem (DGJLLM, 2013)

Suppose that  $A, B \subseteq \mathbb{Z}$  are such that  $\bar{d}(A) = \alpha > 0$  and  $BD(B) > 0$ . Then  $A + B$  is **upper syndetic of level  $\alpha$** : there is  $m \in \mathbb{N}$  such that, for all  $k \in \mathbb{N}$ , we have

$$\bar{d}(\{x \in \mathbb{Z} : x + [-k, k] \subseteq A + B + [-m, m]\}) \geq \alpha.$$

This proof also used nonstandard analysis. In particular, we had to formulate and prove a version of the Lebesgue Density Theorem for quotients of Loeb measure spaces. The proof actually works for subsets of  $\mathbb{Z}^d$  for any  $d$ .

1 A quantitative version of Jin's Theorem

2 Amenable groups

3 Proof of the first part

# Amenable groups

Let  $G$  be a countable (discrete) group.

## Definition

$G$  is *amenable* if  $G$  admits a (left) **Folner sequence**, namely a sequence  $\mathcal{S} := (S_n)$  of finite subsets of  $G$  such that, for every  $g \in G$ , we have

$$\lim_{n \rightarrow \infty} \frac{|gS_n \Delta S_n|}{|S_n|} = 0.$$

Examples of (countable) amenable groups include finite groups and virtually solvable groups. Free groups are not amenable.

# Densities in amenable groups

## Definition

Suppose that  $G$  is an amenable group and  $A$  is a subset of  $G$ .

- 1 If  $\mathcal{S} = (S_n)$  is a Folner sequence for  $G$ , the (upper)  $\mathcal{S}$ -density of  $A$  is

$$d_{\mathcal{S}}(A) := \limsup_{n \rightarrow \infty} \frac{|A \cap S_n|}{|S_n|}.$$

- 2 The *Banach density* of  $A$  is

$$\text{BD}(A) := \sup\{d_{\mathcal{S}}(A) : \mathcal{S} \text{ a Folner sequence for } G\}.$$

## Remark

Suppose  $G = \mathbb{Z}$ . If  $S_n = [-n, n]$ , then  $d_{\mathcal{S}} = \bar{d}$ . One can also check that the notion of Banach density is the same.



# Amenable group version of Jin's Theorem

## Theorem (Beiglböck, Bergelson, Fish, 2009)

If  $G$  is a countable amenable group,  $A, B \subseteq G$  with  $\text{BD}(A), \text{BD}(B) > 0$ , then  $AB$  is *piecewise syndetic*: there is a finite set  $E \subseteq G$  so that, for all finite sets  $L \subseteq G$ , there is  $x \in G$  with  $Lx \subseteq EAB$ .

- This theorem was originally proven using ergodic theory.
- Later, Di Nasso and Lupini gave a combinatorial proof using nonstandard analysis that also worked for uncountable amenable groups and which showed one could assume  $|E| \leq \lfloor \frac{1}{\text{BD}(A)\text{BD}(B)} \rfloor$ .
- The point of this talk is to show how we can achieve a quantitative version of the above theorem generalizing our theorem for  $\mathbb{Z}^d$ .

# $\mathcal{S}$ -thick and $\mathcal{S}$ -syndetic

## Definition

Suppose that  $G$  is a countable amenable group,  $\mathcal{S}$  is a Folner sequence for  $G$ ,  $A$  is a subset of  $G$ , and  $\alpha > 0$ .

- 1 We say that  $A$  is  **$\mathcal{S}$ -thick of level  $\alpha$**  if, for any finite  $L \subseteq G$ , we have

$$d_{\mathcal{S}}(\{x \in G : Lx \subseteq A\}) \geq \alpha.$$

- 2 We say that  $A$  is  **$\mathcal{S}$ -syndetic of level  $\alpha$**  if there is a finite  $E \subseteq G$  such that  $EA$  is  $\mathcal{S}$ -thick of level  $\alpha$ .

# The main results

## Theorem (DGJLLM, 2015)

*Suppose that  $G$  is a countable amenable group and  $S$  a Folner sequence. Further suppose that  $A, B \subseteq G$  are such that  $d_S(A) = \alpha > 0$  and  $\text{BD}(B) > 0$ . Then  $BA$  is  $S$ -syndetic of level  $\alpha'$  for every  $\alpha' < \alpha$ . If, in addition,  $G$  is abelian, then  $BA$  is  $S$ -syndetic of level  $\alpha$ .*

- When  $G = \mathbb{Z}^d$  and  $S_n = [-n, n]^d$ , this recovers our earlier theorem.
- This also recovers another theorem of ours: if  $A, B \subseteq \mathbb{Z}^d$  are such that  $\underline{d}(A) = \alpha > 0$  and  $\text{BD}(B) > 0$ , then  $A + B$  is  $S'$ -syndetic for any subsequence  $S'$  of  $S$ .
- The moreover part: we may assume that  $d_S(A \cup gA) > \alpha$  for some  $g \in G$  (otherwise you can show that  $A$  is already  $S$ -thick of level  $\alpha$ ) and that  $\text{BD}(B) > 1/2$ , so  $\text{BD}(B \cap Bg^{-1}) > 0$ .

# The main results

## Theorem (DGJLLM, 2015)

*Suppose that  $G$  is a countable amenable group and  $S$  a Folner sequence. Further suppose that  $A, B \subseteq G$  are such that  $d_S(A) = \alpha > 0$  and  $\text{BD}(B) > 0$ . Then  $BA$  is  $S$ -syndetic of level  $\alpha'$  for every  $\alpha' < \alpha$ . If, in addition,  $G$  is abelian, then  $BA$  is  $S$ -syndetic of level  $\alpha$ .*

- When  $G = \mathbb{Z}^d$  and  $S_n = [-n, n]^d$ , this recovers our earlier theorem.
- This also recovers another theorem of ours: if  $A, B \subseteq \mathbb{Z}^d$  are such that  $\underline{d}(A) = \alpha > 0$  and  $\text{BD}(B) > 0$ , then  $A + B$  is  $S'$ -syndetic for any subsequence  $S'$  of  $S$ .
- The moreover part: we may assume that  $d_S(A \cup gA) > \alpha$  for some  $g \in G$  (otherwise you can show that  $A$  is already  $S$ -thick of level  $\alpha$ ) and that  $\text{BD}(B) > 1/2$ , so  $\text{BD}(B \cap Bg^{-1}) > 0$ .

1 A quantitative version of Jin's Theorem

2 Amenable groups

3 Proof of the first part

# A dictionary: ultrapowers and nonstandard analysis

- $A^*$  = an ultrapower of  $A$
- internal subset of  $A^*$  =ultraproduct of subsets of  $A$
- hyperfinite subset of  $A^*$  =ultraproduct of finite subsets of  $A$
- If  $E \subseteq A^*$  is hyperfinite, say  $E = \prod_{\mathcal{U}} E_n$ , then  $|E| = (|E_n|)^\bullet \in \mathbb{N}^*$
- If  $\nu \in \mathbb{N}^*$ , say  $\nu = (a_n)^\bullet$ , then  $\nu > \mathbb{N}$  if and only if  $(a_n)$  is  $\mathcal{U}$ -unbounded
- If  $x \in \mathbb{R}^*$  and  $|x| \leq n$  for some  $n \in \mathbb{N}$ , then there is a unique  $r \in \mathbb{R}$  such that  $|x - r|$  is infinitesimal;  $r$  is called the *standard part* of  $x$  and is denoted  $\text{st}(x)$

# Loeb measure

Suppose that  $X$  is a hyperfinite set. We can define a finitely additive measure  $\mu_X$  on the collection of internal subsets of  $X$  given by

$$\mu_X(A) := \text{st} \left( \frac{|A|}{|X|} \right).$$

$\mu_X$  then extends to a probability measure on the  $\sigma$ -algebra of *Loeb measurable* subsets of  $X$ . For internal  $C \subseteq G^*$ , we write  $\mu_X(C)$  instead of  $\mu_X(C \cap X)$ .

# Folner approximations

## Fact (Di Nasso and Lupini)

$G$  is amenable if and only if  $G$  has a *Folner approximation*, which is a hyperfinite subset  $Y \subseteq G^*$  such that, for all  $g \in G$ , we have

$$\frac{|gY \Delta Y|}{|Y|} \approx 0,$$

or, equivalently, for all  $g \in G$ , we have  $\mu_Y(gY) = 1$ . In this case, for any  $A \subseteq G$ , we have

$$\text{BD}(A) = \max\{\mu_X(A^*) : X \text{ a Folner approximation for } G\}.$$

For example, if  $\mathcal{S} = (S_n)$  is a Folner sequence for  $G$ , then  $S_\nu$  is a Folner approximation for  $G$  whenever  $\nu > \mathbb{N}$ .



# The “process”

- Fix a hyperfinite subset  $\Gamma$  of  $G^*$  and  $\nu > \mathbb{N}$ .
- We define a sequence  $(H_n)$  of subsets of  $G$  and a sequence  $(s_n)$  from  $G$  as follows.
- $H_0 := \{g \in G : \mu_\Gamma(\{x \in \Gamma : gx \in (BA)^*\}) < 1\}$ .
- Suppose that  $H_n$  has been defined and is not empty. Let  $s_n \in H_n$  be arbitrary and define

$$H_{n+1} := \{g \in G : \mu_\Gamma(\{x \in \Gamma : gx \in (\{s_0, \dots, s_n\}BA)^*\}) < 1\}.$$

- **Suppose**  $H_n = \emptyset$  and let  $E := \{s_0, \dots, s_n - 1\}$ . Further **suppose** that  $\mu_{S_\nu}(\Gamma) \geq \alpha$ . We claim that  $EBA$  is  $\mathcal{S}$ -thick of level  $\alpha$ .
- Fix  $L \subseteq G$  finite. Since  $H_n = \emptyset$ , we have that  $Lx \subseteq (EBA)^*$  for almost all  $x \in \Gamma$ , whence  $\mu_{S_\nu}(\{x \in S_\nu : Lx \subseteq (EBA)^*\}) \geq \alpha$ .
- But for any  $C \subseteq G$ , we have  $d_{\mathcal{S}}(C) = \max(\{\mu_{S_\nu}(C^*) : \nu > \mathbb{N}\})$ .

# The “process”

- Fix a hyperfinite subset  $\Gamma$  of  $G^*$  and  $\nu > \mathbb{N}$ .
- We define a sequence  $(H_n)$  of subsets of  $G$  and a sequence  $(s_n)$  from  $G$  as follows.
- $H_0 := \{g \in G : \mu_\Gamma(\{x \in \Gamma : gx \in (BA)^*\}) < 1\}$ .
- Suppose that  $H_n$  has been defined and is not empty. Let  $s_n \in H_n$  be arbitrary and define

$$H_{n+1} := \{g \in G : \mu_\Gamma(\{x \in \Gamma : gx \in (\{s_0, \dots, s_n\}BA)^*\}) < 1\}.$$

- **Suppose**  $H_n = \emptyset$  and let  $E := \{s_0, \dots, s_n - 1\}$ . Further **suppose** that  $\mu_{S_\nu}(\Gamma) \geq \alpha$ . We claim that  $EBA$  is  $\mathcal{S}$ -thick of level  $\alpha$ .
- Fix  $L \subseteq G$  finite. Since  $H_n = \emptyset$ , we have that  $Lx \subseteq (EBA)^*$  for almost all  $x \in \Gamma$ , whence  $\mu_{S_\nu}(\{x \in S_\nu : Lx \subseteq (EBA)^*\}) \geq \alpha$ .
- But for any  $C \subseteq G$ , we have  $d_{\mathcal{S}}(C) = \max(\{\mu_{S_\nu}(C^*) : \nu > \mathbb{N}\})$ .

# The “process”

- Fix a hyperfinite subset  $\Gamma$  of  $G^*$  and  $\nu > \mathbb{N}$ .
- We define a sequence  $(H_n)$  of subsets of  $G$  and a sequence  $(s_n)$  from  $G$  as follows.
- $H_0 := \{g \in G : \mu_\Gamma(\{x \in \Gamma : gx \in (BA)^*\}) < 1\}$ .
- Suppose that  $H_n$  has been defined and is not empty. Let  $s_n \in H_n$  be arbitrary and define

$$H_{n+1} := \{g \in G : \mu_\Gamma(\{x \in \Gamma : gx \in (\{s_0, \dots, s_n\}BA)^*\}) < 1\}.$$

- **Suppose**  $H_n = \emptyset$  and let  $E := \{s_0, \dots, s_n - 1\}$ . Further **suppose** that  $\mu_{S_\nu}(\Gamma) \geq \alpha$ . We claim that  $EBA$  is  $\mathcal{S}$ -thick of level  $\alpha$ .
- Fix  $L \subseteq G$  finite. Since  $H_n = \emptyset$ , we have that  $Lx \subseteq (EBA)^*$  for almost all  $x \in \Gamma$ , whence  $\mu_{S_\nu}(\{x \in S_\nu : Lx \subseteq (EBA)^*\}) \geq \alpha$ .
- But for any  $C \subseteq G$ , we have  $d_{\mathcal{S}}(C) = \max(\{\mu_{S_\nu}(C^*) : \nu > \mathbb{N}\})$ .

# The “process”

- Fix a hyperfinite subset  $\Gamma$  of  $G^*$  and  $\nu > \mathbb{N}$ .
- We define a sequence  $(H_n)$  of subsets of  $G$  and a sequence  $(s_n)$  from  $G$  as follows.
- $H_0 := \{g \in G : \mu_\Gamma(\{x \in \Gamma : gx \in (BA)^*\}) < 1\}$ .
- Suppose that  $H_n$  has been defined and is not empty. Let  $s_n \in H_n$  be arbitrary and define

$$H_{n+1} := \{g \in G : \mu_\Gamma(\{x \in \Gamma : gx \in (\{s_0, \dots, s_n\}BA)^*\}) < 1\}.$$

- **Suppose**  $H_n = \emptyset$  and let  $E := \{s_0, \dots, s_n - 1\}$ . Further **suppose** that  $\mu_{S_\nu}(\Gamma) \geq \alpha$ . We claim that  $EBA$  is  $\mathcal{S}$ -thick of level  $\alpha$ .
- Fix  $L \subseteq G$  finite. Since  $H_n = \emptyset$ , we have that  $Lx \subseteq (EBA)^*$  for almost all  $x \in \Gamma$ , whence  $\mu_{S_\nu}(\{x \in S_\nu : Lx \subseteq (EBA)^*\}) \geq \alpha$ .
- But for any  $C \subseteq G$ , we have  $d_{\mathcal{S}}(C) = \max(\{\mu_{S_\nu}(C^*) : \nu > \mathbb{N}\})$ .

# The “process”

- Fix a hyperfinite subset  $\Gamma$  of  $G^*$  and  $\nu > \mathbb{N}$ .
- We define a sequence  $(H_n)$  of subsets of  $G$  and a sequence  $(s_n)$  from  $G$  as follows.
- $H_0 := \{g \in G : \mu_\Gamma(\{x \in \Gamma : gx \in (BA)^*\}) < 1\}$ .
- Suppose that  $H_n$  has been defined and is not empty. Let  $s_n \in H_n$  be arbitrary and define

$$H_{n+1} := \{g \in G : \mu_\Gamma(\{x \in \Gamma : gx \in (\{s_0, \dots, s_n\}BA)^*\}) < 1\}.$$

- **Suppose**  $H_n = \emptyset$  and let  $E := \{s_0, \dots, s_n - 1\}$ . Further **suppose** that  $\mu_{S_\nu}(\Gamma) \geq \alpha$ . We claim that  $EBA$  is  $\mathcal{S}$ -thick of level  $\alpha$ .
- Fix  $L \subseteq G$  finite. Since  $H_n = \emptyset$ , we have that  $Lx \subseteq (EBA)^*$  for almost all  $x \in \Gamma$ , whence  $\mu_{S_\nu}(\{x \in S_\nu : Lx \subseteq (EBA)^*\}) \geq \alpha$ .
- But for any  $C \subseteq G$ , we have  $d_{\mathcal{S}}(C) = \max(\{\mu_{S_\nu}(C^*) : \nu > \mathbb{N}\})$ .

# The “process”

- Fix a hyperfinite subset  $\Gamma$  of  $G^*$  and  $\nu > \mathbb{N}$ .
- We define a sequence  $(H_n)$  of subsets of  $G$  and a sequence  $(s_n)$  from  $G$  as follows.
- $H_0 := \{g \in G : \mu_\Gamma(\{x \in \Gamma : gx \in (BA)^*\}) < 1\}$ .
- Suppose that  $H_n$  has been defined and is not empty. Let  $s_n \in H_n$  be arbitrary and define

$$H_{n+1} := \{g \in G : \mu_\Gamma(\{x \in \Gamma : gx \in (\{s_0, \dots, s_n\}BA)^*\}) < 1\}.$$

- **Suppose**  $H_n = \emptyset$  and let  $E := \{s_0, \dots, s_n - 1\}$ . Further **suppose** that  $\mu_{S_\nu}(\Gamma) \geq \alpha$ . We claim that  $EBA$  is  $\mathcal{S}$ -thick of level  $\alpha$ .
- Fix  $L \subseteq G$  finite. Since  $H_n = \emptyset$ , we have that  $Lx \subseteq (EBA)^*$  for almost all  $x \in \Gamma$ , whence  $\mu_{S_\nu}(\{x \in S_\nu : Lx \subseteq (EBA)^*\}) \geq \alpha$ .
- But for any  $C \subseteq G$ , we have  $d_S(C) = \max(\{\mu_{S_\nu}(C^*) : \nu > \mathbb{N}\})$ .

# The “process” (continued)

- We are seeking hyperfinite  $\Gamma \subseteq G^*$  with  $\mu_{S_\nu}(\Gamma) \geq \alpha$  such that the process stops at some finite stage.
- **Suppose** there is standard  $r > 0$  and a Folner approximation  $Y$  of  $G$  such that, setting  $C := A^* \cap S_\nu$ ,  $D := B^* \cap Y$ , and

$$\Gamma := \left\{ x \in S_\nu : \frac{|xC^{-1} \cap D|}{|Y|} \geq r \right\},$$

we have  $\mu_{S_\nu}(\Gamma) \geq \alpha$ .

- We then claim that  $H_n = \emptyset$  for  $n > \frac{1}{r}$ .

# The process (continued)

- Suppose that  $H_n \neq \emptyset$ . For  $k = 0, \dots, n$ , fix  $s_k \in H_k$  and take  $\gamma_k \in \Gamma$  so that  $s_k \gamma_k \notin (\{s_0, \dots, s_{k-1}\}BA)^*$ .
- Note then that if  $0 \leq i < j \leq n-1$ , then  $s_j \gamma_j (A^*)^{-1} \cap s_i B^* = \emptyset$ .
- It follows that the sets  $s_k ((\gamma_k C^{-1}) \cap D)$  for  $k = 0, \dots, n-1$  are pairwise disjoint.
- Since  $Y$  is a Folner approximation for  $G$ , we have

$$1 \geq \text{st} \left( \frac{|\bigcup_{k=0}^{n-1} s_k ((\gamma_k C^{-1}) \cap D)|}{|Y|} \right) = \sum_{k=0}^{n-1} \mu_Y((\gamma_k C^{-1}) \cap D) \geq nr.$$

- It follows that  $n \leq \frac{1}{r}$ .



# The key lemma

## Lemma

Suppose  $d_S(A) \geq \alpha$  and  $\text{BD}(B) \geq \beta$ . Then there exists a Folner approximation  $Y$  of  $G$  and  $\nu > \mathbb{N}$  such that, setting  $C := A^* \cap S_\nu$  and  $D := B^* \cap Y$ , we have:

- 1  $\mu_{S_\nu}(C) \geq \alpha$ ;
- 2  $\mu_Y(D) \geq \beta$ ;
- 3  $\text{st} \left( \frac{1}{|S_\nu|} \sum_{x \in S_\nu} \frac{|(xC^{-1}) \cap D|}{|Y|} \right) \geq \alpha\beta$ .

# Finishing the process from the key lemma

- Recall  $d_S(A) > \alpha$  and  $BD(B) > 0$ .
- Fact: there is a finite  $T$  such that  $d_S(A) \cdot BD(TB) > \alpha$ . Since  $BA$  is  $S$ -syndetic of level  $\alpha$  if and only if  $TBA$  is  $S$ -syndetic of level  $\alpha$ , we may suppose that  $T = \{1\}$ .
- Take standard  $r > 0$  so that  $d_S(A) \cdot BD(B) > \alpha + r$  and recall  $\Gamma := \{x \in S_\nu : \frac{|(xC^{-1}) \cap D|}{|Y|} \geq r\}$ .
- Take  $Y$  and  $\nu$  as in the previous lemma. We then have

$$\alpha + r < \frac{1}{|S_\nu|} \sum_{x \in S_\nu} \frac{|(xC^{-1}) \cap D|}{|Y|} \leq \frac{1}{|S_\nu|} \left( \sum_{x \in \Gamma} + \sum_{x \notin \Gamma} \right) \leq \frac{|\Gamma|}{|S_\nu|} + r.$$

# Finishing the process from the key lemma

- Recall  $d_S(A) > \alpha$  and  $\text{BD}(B) > 0$ .
- Fact: there is a finite  $T$  such that  $d_S(A) \cdot \text{BD}(TB) > \alpha$ . Since  $BA$  is  $S$ -syndetic of level  $\alpha$  if and only if  $TBA$  is  $S$ -syndetic of level  $\alpha$ , we may suppose that  $T = \{1\}$ .
- Take standard  $r > 0$  so that  $d_S(A) \cdot \text{BD}(B) > \alpha + r$  and recall  $\Gamma := \{x \in S_\nu : \frac{|(xC^{-1}) \cap D|}{|Y|} \geq r\}$ .
- Take  $Y$  and  $\nu$  as in the previous lemma. We then have

$$\alpha + r < \frac{1}{|S_\nu|} \sum_{x \in S_\nu} \frac{|(xC^{-1}) \cap D|}{|Y|} \leq \frac{1}{|S_\nu|} \left( \sum_{x \in \Gamma} + \sum_{x \notin \Gamma} \right) \leq \frac{|\Gamma|}{|S_\nu|} + r.$$

# Proving the key lemma

- Take any Følner approximation  $Y$  for  $G$  such that  $\mu_Y(D) \geq \beta$ .
- Since  $d_S(A) \geq \alpha$ , the following statement is true: for any finite  $E \subseteq G$  and any  $n_0 \in \mathbb{N}$ , there is  $n > n_0$  such that  $\frac{|A \cap S_n|}{|S_n|} \geq \alpha - 2^{-n_0}$  and for which, for all  $g \in E$ , we have  $\frac{|g^{-1}S_n \Delta S_n|}{|S_n|} < 2^{-n_0}$ .
- Apply the transferred version of this statement to  $Y$  and some given  $\nu_0 > \mathbb{N}$  to get  $\nu > \nu_0$  such that  $\mu_{S_\nu}(C) \geq \alpha$  and for which  $\frac{|g^{-1}S_\nu \Delta S_\nu|}{|S_\nu|} \approx 0$  for all  $g \in E$ .
- We need to check the third condition of the lemma.

# Proving the key lemma

- Take any Folner approximation  $Y$  for  $G$  such that  $\mu_Y(D) \geq \beta$ .
- Since  $d_S(A) \geq \alpha$ , the following statement is true: for any finite  $E \subseteq G$  and any  $n_0 \in \mathbb{N}$ , there is  $n > n_0$  such that  $\frac{|A \cap S_n|}{|S_n|} \geq \alpha - 2^{-n_0}$  and for which, for all  $g \in E$ , we have  $\frac{|g^{-1}S_n \Delta S_n|}{|S_n|} < 2^{-n_0}$ .
- Apply the transferred version of this statement to  $Y$  and some given  $\nu_0 > \mathbb{N}$  to get  $\nu > \nu_0$  such that  $\mu_{S_\nu}(C) \geq \alpha$  and for which  $\frac{|g^{-1}S_\nu \Delta S_\nu|}{|S_\nu|} \approx 0$  for all  $g \in E$ .
- We need to check the third condition of the lemma.

# Proving the key lemma

- Take any Følner approximation  $Y$  for  $G$  such that  $\mu_Y(D) \geq \beta$ .
- Since  $d_S(A) \geq \alpha$ , the following statement is true: for any finite  $E \subseteq G$  and any  $n_0 \in \mathbb{N}$ , there is  $n > n_0$  such that  $\frac{|A \cap S_n|}{|S_n|} \geq \alpha - 2^{-n_0}$  and for which, for all  $g \in E$ , we have  $\frac{|g^{-1}S_n \Delta S_n|}{|S_n|} < 2^{-n_0}$ .
- Apply the transferred version of this statement to  $Y$  and some given  $\nu_0 > \mathbb{N}$  to get  $\nu > \nu_0$  such that  $\mu_{S_\nu}(C) \geq \alpha$  and for which  $\frac{|g^{-1}S_\nu \Delta S_\nu|}{|S_\nu|} \approx 0$  for all  $g \in E$ .
- We need to check the third condition of the lemma.

# We finish by counting

$$\begin{aligned}
 \frac{1}{|S_\nu|} \sum_{x \in S_\nu} \frac{|xC^{-1} \cap D|}{|Y|} &= \frac{1}{|S_\nu|} \sum_{x \in S_\nu} \frac{1}{|Y|} \sum_{d \in D} \chi_{C^{-1}}(x^{-1}d) \\
 &= \frac{1}{|Y|} \sum_{d \in D} \frac{1}{|S_\nu|} \sum_{x \in S_\nu} \chi_{C^{-1}}(x^{-1}d) \\
 &= \frac{1}{|Y|} \sum_{d \in D} \frac{|S_\nu^{-1}d \cap C^{-1}|}{|S_\nu|} \\
 &\geq \frac{1}{|Y|} \sum_{d \in D} \left( \frac{|C^{-1}|}{|S_\nu|} - \frac{|d^{-1}S_\nu \Delta S_\nu|}{|S_\nu|} \right) \\
 &\geq \frac{|C|}{|S_\nu|} \frac{|D|}{|Y|} - \text{an infinitesimal.}
 \end{aligned}$$

# Lower density versions?

## Theorem (DGJLLM, 2013)

Suppose that  $A, B \subseteq \mathbb{Z}^d$ .

- 1 Suppose that  $\underline{d}(A) = \alpha > 0$  and  $\text{BD}(B) > 0$ . Then for any  $\epsilon > 0$ , there is  $m \in \mathbb{N}$  such that, for all  $k \in \mathbb{N}$ , we have

$$\underline{d}(\{x \in \mathbb{Z}^d : x + [-k, k] \subseteq A + B + [-m, m]\}) \geq \alpha - \epsilon.$$

- 2 Suppose that  $d = 1$ ,  $\underline{d}(A) = \alpha > 0$  and  $\underline{d}(B) = \beta > 0$ . Then there is  $m \in \mathbb{N}$  such that, for all  $k \in \mathbb{N}$ , we have

$$\underline{d}(\{x \in \mathbb{Z} : x + [-k, k] \subseteq A + B + [-m, m]\}) \geq \min(\alpha + \beta, 1).$$

## Question

Can we prove amenable group versions of these results?



# References

- M. Di Nasso, I. Goldbring, R. Jin, S. Leth, M. Lupini, and K. Mahlburg, *High density piecewise syndeticity of product sets in amenable groups*, submitted. arXiv 1505.04701
- M. Di Nasso, I. Goldbring, R. Jin, S. Leth, M. Lupini, and K. Mahlburg, *High density piecewise syndeticity of sumsets*, *Advances in Math.* Volume 278 (2015), 1-33.