


This article was downloaded by:[Weizmann Institute Science]
On: 22 July 2008
Access Details: [subscription number 791802962]
Publisher: Taylor & Francis
Informa Ltd Registered in England and Wales Registered Number: 1072954
Registered office: Mortimer House, 37-41 Mortimer Street, London W1T 3JH, UK



Communications in Partial Differential Equations

Publication details, including instructions for authors and subscription information:
<http://www.informaworld.com/smpp/title-content=t713597240>

Regularity Criterion for Solutions of Three-Dimensional Turbulent Channel Flows

Chongsheng Cao ^a; Junlin Qin ^b; Edriss S. Titi ^{cd}

^a Department of Mathematics, Florida International University, University Park, Miami, Florida, USA

^b Department of Mathematics, Xian Jiaotong University, Xian, Shaanxi, China

^c Department of Mathematics and Department of Mechanical and Aerospace Engineering, University of California, Irvine, California, USA

^d Department of Computer Science and Applied Mathematics, Weizmann Institute of Science, Rehovot, Israel

Online Publication Date: 01 March 2008

To cite this Article: Cao, Chongsheng, Qin, Junlin and Titi, Edriss S. (2008) 'Regularity Criterion for Solutions of Three-Dimensional Turbulent Channel Flows', Communications in Partial Differential Equations, 33:3, 419 — 428

To link to this article: DOI: 10.1080/03605300701454859

URL: <http://dx.doi.org/10.1080/03605300701454859>

PLEASE SCROLL DOWN FOR ARTICLE

Full terms and conditions of use: <http://www.informaworld.com/terms-and-conditions-of-access.pdf>

This article maybe used for research, teaching and private study purposes. Any substantial or systematic reproduction, re-distribution, re-selling, loan or sub-licensing, systematic supply or distribution in any form to anyone is expressly forbidden.

The publisher does not give any warranty express or implied or make any representation that the contents will be complete or accurate or up to date. The accuracy of any instructions, formulae and drug doses should be independently verified with primary sources. The publisher shall not be liable for any loss, actions, claims, proceedings, demand or costs or damages whatsoever or howsoever caused arising directly or indirectly in connection with or arising out of the use of this material.

Regularity Criterion for Solutions of Three-Dimensional Turbulent Channel Flows

CHONGSHENG CAO¹, JUNLIN QIN²,
AND EDRISS S. TITI^{3,4}

¹Department of Mathematics, Florida International University,
University Park, Miami, Florida, USA

²Department of Mathematics, Xian Jiaotong University,
Xian, Shaanxi, China

³Department of Mathematics and Department of Mechanical and
Aerospace Engineering, University of California, Irvine, California, USA

⁴Department of Computer Science and Applied Mathematics,
Weizmann Institute of Science, Rehovot, Israel

In this paper we consider the three-dimensional Navier–Stokes equations in infinite channel. We provide a regularity criterion for solutions of the three-dimensional Navier–Stokes equations in terms of the vertical component of the velocity field.

Keywords Regularity criterion; Three-dimensional Navier–Stokes equations; Turbulent channel flows.

Mathematics Subject Classification 35Q35; 65M70.

1. Introduction

Turbulence stands out as a prototype of multi-scale phenomenon that occurs in nature. It involves wide ranges of spatial and temporal scales which makes it very difficult to study analytically and prohibitively expensive to simulate computationally. Turbulent channel flows are considered to be the simplest flows confined within physical boundaries that can be simulated numerically and that demonstrates many of the common features of turbulence. In this paper we consider three-dimensional finite energy turbulent flows of viscous incompressible homogeneous fluids in the infinite channel $\Omega = \mathbb{R}^2 \times [-L, L] \subset \mathbb{R}^3$, subject to the no-slip Dirichlet boundary conditions. These flows are governed by the

Received September 1, 2006; Accepted March 1, 2007

Address correspondence to Edriss S. Titi, Department of Mathematics, University of California, Irvine, CA 92697-3875, USA; E-mail: etiti@math.uci.edu

three-dimensional Navier–Stokes system of equations:

$$\frac{\partial u}{\partial t} - \nu \Delta u + (u \cdot \nabla)u + \nabla p = f \quad \text{in } \Omega \quad (1)$$

$$\nabla \cdot u = 0 \quad \text{in } \Omega \quad (2)$$

$$u = 0 \quad \text{on } \partial\Omega \quad (3)$$

$$\lim_{x \rightarrow \infty} u(t, x) = 0 \quad (4)$$

$$u(0, x) = u_0(x) \quad \text{in } \Omega. \quad (5)$$

Here, $u = (u_1, u_2, u_3)$ represents the unknown velocity vector field, and p is the unknown pressure scalar; where $\nu > 0$, the constant kinematic viscosity, f , the body forcing term, and u_0 , the initial velocity, are given.

Mathematically, it is well-known that the three-dimensional system (1)–(5) has global (for all time and all initial data) weak solutions (see, e.g., Constantin and Foias, 1988; Galdi, 1994; Ladyzhenskaya, 1969; Sohr, 2001; Temam, 1983, 2001 and references therein). The question of well-posedness, in the sense of Hadamard, and in particular the question of uniqueness, of these weak solutions is still an open problem. On the other hand, it is also well-established (see, e.g., Constantin and Foias, 1988; Galdi, 1994; Ladyzhenskaya, 1969; Sohr, 2001; Temam, 1983, 2001 and references therein) that the system (1)–(5) possesses a unique strong (regular) solution, which depends continuously on the initial data, for a short interval of time $[0, T_*)$, where T_* depends on the size of initial datum, u_0 , on f , ν , and L . Moreover, it is also well-known that the existence (for all time) and uniqueness of strong (regular) solutions is guaranteed under suitable additional assumptions (see, e.g., Berselli, 2002; Berselli and Galdi, 2002; Cao and Titi, 2007; Chae and Lee, 2001; He, 2002; Kato, 1984; Kukavica and Ziane, 2006; Moise, 2006; Pokorný, 2003; Prodi, 1959; Serrin, 1962; Zhou, 2002 and references therein). In particular, some of these recent results involve conditions on only one component of the velocity field of the 3D NSE in the whole space \mathbb{R}^3 or under periodic boundary conditions (see, e.g., He, 2002; Kukavica and Ziane, 2006; Pokorný, 2003; Zhou, 2002). In this paper, we study this type of sufficient conditions for the global regularity of the 3D NSE in the infinite channel Ω , subject to no-slip Dirichlet boundary condition on the physical boundary of the channel (see also, Moise, 2006). Using the geophysical terminology, our condition is formulated in terms of the third component of the baroclinic mode \tilde{u}_3 (see (16), below, for the definition of the barotropic mode (vertically averaged mode), \bar{u} , and the baroclinic mode (the fluctuation about the barotropic mode)). Specifically, our results states that if \tilde{u}_3 satisfies

$$\nabla \tilde{u}_3 \in L^\infty([0, \infty), L^2(\Omega)), \quad (6)$$

then the strong (regular) unique solution of the 3D Navier–Stokes equation (1)–(5) exists for all time.

Let us observe that our condition (6) seems to be slightly tighter than the former ones (cf. e.g., He, 2002; Kukavica and Ziane, 2006; Pokorný, 2003; Zhou, 2002). However, unlike the previous works we study here the 3D Navier–Stokes in a domain with physical boundaries under the no-slip Dirichlet boundary conditions. Furthermore, we emphasize that the techniques developed here, which are inspired

by ideas presented in Cao and Titi (2007), are totally different than the previous ones.

Let us denote by $L^q(\Omega)$, $L^q(\mathbb{R}^2)$, and $H^m(\Omega)$, $H^m(\mathbb{R}^2)$ the usual L^q -Lebesgue and Sobolev spaces, respectively (Adams, 1975). We denote by

$$\|\phi\|_q = \begin{cases} \left(\int_{\Omega} |\phi|^q dx_1 dx_2 dx_3 \right)^{\frac{1}{q}}, & \text{for every } \phi \in L^q(\Omega) \\ \left(\int_{\mathbb{R}^2} |\phi|^q dx_1 dx_2 \right)^{\frac{1}{q}}, & \text{for every } \phi \in L^q(\mathbb{R}^2). \end{cases} \tag{7}$$

Let

$$\mathcal{V} = \{v \in C_0^\infty(\Omega) : \nabla \cdot v = 0\}.$$

Since we are interested in flows of finite energy in the infinite channel Ω , we consider the spaces H and V , defined to be the closures of the set \mathcal{V} in $L^2(\Omega)$ under L^2 -topology, and in $H^1(\Omega)$ under H^1 -topology, respectively. Denote by $P : L^2 \rightarrow H$, the orthogonal projection, and let $A = -P\Delta$ be the Stokes operator subject to the homogeneous Dirichlet boundary condition (3). It is well known that the Navier–Stokes equations (NSE) (1)–(5) are equivalent to the functional differential equation (see, e.g., Constantin and Foias, 1988; Sohr, 2001; Temam, 1983, 2001)

$$\frac{du}{dt} + vAu + B(u, u) = f, \tag{8}$$

$$u(0) = u_0, \tag{9}$$

where $B(u, u) = P((u \cdot \nabla)u)$, the nonlinear (bilinear) term. We say u is a Leray–Hopf weak solution to the system (8)–(9) if u satisfies (see, e.g., Constantin and Foias, 1988; Temam, 1983, 2001)

- (1) $u \in C([0, T], H\text{-weak}) \cap L^2([0, T], V)$, and $\partial_t u \in L^1([0, T], V')$, where V' is the dual space of V ,
- (2) the weak formulation:

$$\begin{aligned} & \int_{\Omega} u(t, x) \cdot \phi(x) dx - \int_{\Omega} u(t_0, x) \cdot \phi(x) dx \\ &= - \int_{t_0}^t \int_{\Omega} (v(\nabla u(s, x)) : \nabla \phi(x)) + (u(s, x) \cdot \nabla)u(s, x) \cdot \phi(x) dx ds \\ &+ \int_{t_0}^t \int_{\Omega} (f(s, x) \cdot \phi(x)) dx ds, \end{aligned}$$

for every $\phi \in \mathcal{V}$, and almost every $t, t_0 \in [0, T]$.

- (3) the energy inequality:

$$\|u(t)\|_2^2 - \|u(t_0)\|_2^2 + 2\nu \int_{t_0}^t \|\nabla u(s)\|_2^2 ds \leq 2 \int_{t_0}^t \int_{\Omega} f(s, x) \cdot u(s, x) dx ds, \tag{10}$$

for all $t \in [0, T]$, and for almost every t_0 in the interval $[0, t]$.

Moreover, a weak solution is called strong solution of (8)–(9) on $[0, T]$ if, in addition, it satisfies

$$u \in C([0, T], V) \cap L^2([0, T], H^2(\Omega)).$$

For convenience, we recall the following Gagliardo–Nirenberg, Ladyzhenskaya, and Sobolev inequalities (cf. e.g., Adams, 1975; Constantin and Foias, 1988; Galdi, 1994; Ladyzhenskaya, 1985, 1969; Sohr, 2001) in \mathbb{R}^2 :

$$\|\phi\|_{L^r(\mathbb{R}^2)} \leq C_r \|\phi\|_{L^2(\mathbb{R}^2)}^{2/r} \|\phi\|_{H^1(\mathbb{R}^2)}^{r-2}, \quad r < \infty, \tag{11}$$

for every $\phi \in H^1(\mathbb{R}^2)$, and in \mathbb{R}^3 :

$$\|\psi\|_{L^\alpha(\Omega)} \leq C_\alpha \|\psi\|_{L^2(\Omega)}^{6-\alpha} \|\psi\|_{H^1(\Omega)}^{\frac{3(\alpha-2)}{2\alpha}}, \tag{12}$$

for every $u \in H^1(\Omega)$, $2 \leq \alpha \leq 6$. Here C_r and C_α are scale invariant constants. We also recall the Pioncaré inequality:

$$\|\nabla v\|_2 \geq \frac{C_0}{L} \|v\|_2 \quad \forall v \in V \tag{13}$$

$$\|Av\|_2 \geq \frac{C_0}{L} \|\nabla v\|_2 \quad \forall v \in \mathcal{D}(A), \tag{14}$$

where C_0 is a scale invariant constant. Also, we recall the integral version of Minkowsky inequality for the L^r spaces, $r \geq 1$. Let $\Omega_1 \subset \mathbb{R}^{m_1}$ and $\Omega_2 \subset \mathbb{R}^{m_2}$ be two measurable sets, where m_1 and m_2 are two positive integers. Suppose that $\phi(\xi, \eta)$ is measurable over $\Omega_1 \times \Omega_2$. Then,

$$\left[\int_{\Omega_1} \left(\int_{\Omega_2} |\phi(\xi, \eta)| d\eta \right)^r d\xi \right]^{1/r} \leq \int_{\Omega_2} \left(\int_{\Omega_1} |\phi(\xi, \eta)|^r d\xi \right)^{1/r} d\eta. \tag{15}$$

2. Global Existence of the Strong Solution

In this section we will show the global existence of the strong solutions to the three-dimensional Navier–Stokes system (1)–(5) under assumption (6).

We will denote by

$$\bar{\theta}(x_1, x_2) = \frac{1}{2L} \int_{-L}^L \theta(x_1, x_2, x_3) dx_3 \quad \text{and} \quad \tilde{\theta} = \theta - \bar{\theta}. \tag{16}$$

Following the geophysical fluid dynamics terminology we will call $\bar{\theta}$ the barotropic mode and $\tilde{\theta}$ the baroclinic mode.

From now on, we will denote by $\nabla_h = (\frac{\partial}{\partial x_1}, \frac{\partial}{\partial x_2})$ and $\Delta_h = \frac{\partial^2}{\partial x_1^2} + \frac{\partial^2}{\partial x_2^2}$. First, let us prove the following lemma.

Lemma 1. *Suppose that $\xi(x_1, x_2) \in H^1(\mathbb{R}^2)$, $\phi \in H^1(\Omega)$, and $\psi \in L^2(\Omega)$. Then,*

$$\int_{\Omega} |\xi| |\phi| |\psi| dx_1 dx_2 dx_3 \leq C \|\xi\|_2^{1/2} (\|\xi\|_2 + \|\nabla_h \xi\|_2)^{1/2} \|\phi\|_2^{1/2} (\|\phi\|_2 + \|\nabla_h \phi\|_2)^{1/2} \|\psi\|_2.$$

Proof. Notice that

$$\int_{\Omega} |\xi| |\phi| |\psi| dx_1 dx_2 dx_3 = \int_{\mathbb{R}^2} \left[|\xi| \left(\int_{-L}^L |\phi| |\psi| dx_3 \right) \right] dx_1 dx_2.$$

We will estimate the above term by applying the same method used to establish Proposition 2.2 in Cao and Titi (2003). First, by Cauchy–Schwarz inequality, we obtain

$$\int_{-L}^L |\phi| |\psi| dx_3 \leq \left(\int_{-L}^L |\phi|^2 dx_3 \right)^{\frac{1}{2}} \left(\int_{-L}^L |\psi|^2 dx_3 \right)^{\frac{1}{2}}.$$

Thus, by the above and Hölder inequality, we reach

$$\begin{aligned} \int_{\Omega} |\xi| |\phi| |\psi| dx_1 dx_2 dx_3 &\leq \int_{\mathbb{R}^2} \left[|\xi| \left(\int_{-L}^L |\phi|^2 dx_3 \right)^{\frac{1}{2}} \left(\int_{-L}^L |\psi|^2 dx_3 \right)^{\frac{1}{2}} \right] dx_1 dx_2 \\ &\leq \left(\int_{\mathbb{R}^2} |\xi|^4 dx_1 dx_2 \right)^{\frac{1}{4}} \left[\int_{\mathbb{R}^2} \left(\int_{-L}^L |\phi|^2 dx_3 \right)^2 dx_1 dx_2 \right]^{\frac{1}{4}} \left[\int_{\Omega} |\psi|^2 dx \right]^{\frac{1}{2}}. \end{aligned}$$

By using Minkowsky inequality (15), we get

$$\left[\int_{\mathbb{R}^2} \left(\int_{-L}^L |\phi|^2 dx_3 \right)^2 dx_1 dx_2 \right]^{\frac{1}{2}} \leq \int_{-L}^L \left(\int_{\mathbb{R}^2} |\phi|^4 dx_1 dx_2 \right)^{\frac{1}{2}} dx_3.$$

Thanks to (11) with $r = 4$, for every fixed x_3 we have

$$\left(\int_{\mathbb{R}^2} |\phi|^4 dx_1 dx_2 \right)^{\frac{1}{4}} \leq C \|\phi\|_{L^2(\mathbb{R}^2)}^{\frac{1}{2}} \|\phi\|_{H^1(\mathbb{R}^2)}^{\frac{1}{2}}.$$

As a result of the above and the Cauchy–Schwarz inequality, we obtain

$$\begin{aligned} \int_{-L}^L \left(\int_{\mathbb{R}^2} |\phi|^4 dx_1 dx_2 \right)^{\frac{1}{2}} dx_3 &\leq C \int_{-L}^L \|\phi\|_{L^2(\mathbb{R}^2)} \|\phi\|_{H^1(\mathbb{R}^2)} dx_3 \\ &\leq C \left(\int_{-L}^L \|\phi\|_{L^2(\mathbb{R}^2)}^2 dx_3 \right)^{\frac{1}{2}} \left(\int_{-L}^L \|\phi\|_{H^1(\mathbb{R}^2)}^2 dx_3 \right)^{\frac{1}{2}} \\ &\leq C (\|\phi\|_2 + \|\nabla_h \phi\|_2) \|\phi\|_2. \end{aligned}$$

Therefore,

$$\left[\int_{\mathbb{R}^2} \left(\int_{-L}^L |\phi|^2 dx_3 \right)^2 dx_1 dx_2 \right]^{\frac{1}{4}} \leq C (\|\phi\|_2 + \|\nabla_h \phi\|_2)^{1/2} \|\phi\|_2^{1/2}. \tag{17}$$

By using (11) with $r = 4$, we have

$$\left[\int_{\mathbb{R}^2} |\xi|^4 dx_1 dx_2 \right]^{\frac{1}{4}} \leq C \|\xi\|_{L^2(\mathbb{R}^2)}^{1/2} \|\xi\|_{H^1(\mathbb{R}^2)}^{1/2}.$$

Thus, by the above and (17), we get

$$\int_{\Omega} |\xi| |\phi| |\psi| dx_1 dx_2 dx_3 \leq C \|\xi\|_2^{1/2} (\|\xi\|_2 + \|\nabla_h \xi\|_2)^{1/2} \|\phi\|_2^{1/2} \times (\|\phi\|_2 + \|\nabla_h \phi\|_2)^{1/2} \|\psi\|_2. \quad \square$$

Theorem 2. Let $f \in L^\infty([0, \infty), L^2(\Omega))$, $u_0 \in V$. Let $u = (u_1, u_2, u_3)$ be a weak solution of the system (8)–(9) in $[0, \infty)$. Suppose that for $T > 0$, $\nabla \tilde{u}_3 \in L^\infty([0, T], L^2(\Omega))$; that is, \tilde{u}_3 satisfies

$$\sup_{0 \leq t \leq T} \|\nabla \tilde{u}_3(t)\|_2 < \infty. \quad (18)$$

Then u is the strong solution of the system (8)–(9) on $[0, T]$.

Proof. Let $u_0 \in V$. Following, for instance, the Galerkin method one can show that there exists a unique strong solution u for the system (8)–(9), with the initial datum u_0 , for a short interval of time (see, e.g., Constantin and Foias, 1988; Ladyzhenskaya, 1969; Sohr, 2001; Temam, 1983, 2001. Suppose that $[0, T_*)$ is the maximal interval of existence of this strong solution u . It is also well known (see, e.g., the above references) that there exists a Leray–Hopf weak solution for the system (8)–(9), with the same initial datum u_0 , which exists globally in time, i.e. for all time $t \geq 0$. Most importantly, following the work of Sather and Serrin (1963) one can show that all the Leray–Hopf weak solutions coincide with the unique strong solution, u , on the interval $[0, T_*)$.

To conclude our proof we need to show that $T < T_*$. Suppose, arguing by contradiction, that $T_* \leq T$, and that (18) holds. If we show that the $\limsup_{t \rightarrow T_*^-} \|u(t)\|_{H^1} < \infty$ then $[0, T_*)$ is not a maximal interval of existence, which leads to a contradiction.

For the rest of this proof we consider the strong solution, u , in the the interval $[0, T_*)$. From the energy inequality of (10), which is satisfied by all the Leray–Hopf weak solutions (see, for example, Constantin and Foias, 1988; Ladyzhenskaya, 1969; Sohr, 2001; Temam, 1983, 2001, for details), we have

$$\|u(t)\|_2^2 \leq C \frac{L^4 F^2}{\nu^2} + e^{-\frac{\nu t}{L^2}} \|u_0\|_2^2, \quad (19)$$

$$\nu \int_0^t \|\nabla u(s)\|_2^2 ds \leq C \frac{L^2 F^2 t}{\nu} + \|u_0\|_2^2, \quad (20)$$

for all $t \in [0, T_*)$, where

$$F = \|f\|_{L^\infty([0, \infty), L^2(\Omega))}. \quad (21)$$

In particular, since $t < T_* \leq T$ we have

$$\|u(t)\|_2^2 + \nu \int_0^t \|\nabla u(s)\|_2^2 ds \leq K_1, \quad (22)$$

where

$$K_1 = C \frac{F^2 (L^4 + \nu T)}{\nu^2} + 2\|u_0\|_2^2. \tag{23}$$

Taking the inner product of the equation (8) with $-\Delta_h u$ in H , and notice that $P\partial_{x_i} = \partial_{x_i}P$ for $i = 1, 2$, we get

$$\frac{1}{2} \frac{d\|\nabla_h u\|_2^2}{dt} + \nu\|\Delta_h u\|_2^2 + \nu\|\nabla_h u_z\|_2^2 = - \int_{\Omega} (f - B(u, u)) \cdot \Delta_h u \, dx_1 \, dx_2 \, dx_3.$$

By integration by parts we get

$$\begin{aligned} - \int_{\Omega} B(u, u) \cdot \Delta_h u \, dx_1 \, dx_2 \, dx_3 &= \int_{\Omega} \sum_{l=1}^2 \frac{\partial u_k}{\partial x_j} \frac{\partial u_j}{\partial x_l} \frac{\partial u_k}{\partial x_l} \, dx_1 \, dx_2 \, dx_3 \\ &= \int_{\Omega} \left\{ \left(\frac{\partial u_1}{\partial x_1} \right)^3 + \left(\frac{\partial u_2}{\partial x_2} \right)^3 + \left(\frac{\partial u_1}{\partial x_1} + \frac{\partial u_2}{\partial x_2} \right) \left[\left(\frac{\partial u_1}{\partial x_2} \right)^2 + \left(\frac{\partial u_2}{\partial x_1} \right)^2 + \frac{\partial u_1}{\partial x_2} \frac{\partial u_2}{\partial x_1} \right] \right. \\ &\quad \left. + \sum_{k,l=1}^2 \frac{\partial u_k}{\partial x_3} \frac{\partial u_3}{\partial x_l} \frac{\partial u_k}{\partial x_l} + \sum_{j,l=1}^2 \frac{\partial u_3}{\partial x_j} \frac{\partial u_j}{\partial x_l} \frac{\partial u_3}{\partial x_l} + \sum_{l=1}^2 \frac{\partial u_3}{\partial x_3} \frac{\partial u_3}{\partial x_l} \frac{\partial u_3}{\partial x_l} \right\} dx_1 \, dx_2 \, dx_3 \\ &= \int_{\Omega} \left\{ - \frac{\partial u_3}{\partial x_3} \left[\left(\frac{\partial u_1}{\partial x_1} \right)^2 + \left(\frac{\partial u_2}{\partial x_2} \right)^2 - \frac{\partial u_1}{\partial x_1} \frac{\partial u_2}{\partial x_2} + \left(\frac{\partial u_1}{\partial x_2} \right)^2 + \left(\frac{\partial u_2}{\partial x_1} \right)^2 + \frac{\partial u_1}{\partial x_2} \frac{\partial u_2}{\partial x_1} \right] \right. \\ &\quad \left. + \sum_{k,l=1}^2 \left(\frac{\partial u_k}{\partial x_l} \frac{\partial u_3}{\partial x_3} \frac{\partial u_k}{\partial x_l} + u_k \frac{\partial u_3}{\partial x_3} \frac{\partial^2 u_k}{\partial x_l \partial x_l} - u_k \frac{\partial u_3}{\partial x_l} \frac{\partial^2 u_k}{\partial x_l \partial x_3} \right) \right. \\ &\quad \left. + \sum_{j,l=1}^2 \frac{\partial u_3}{\partial x_j} \frac{\partial u_j}{\partial x_l} \frac{\partial u_3}{\partial x_l} - \sum_{l=1}^2 \left(\frac{\partial u_1}{\partial x_1} + \frac{\partial u_2}{\partial x_2} \right) \frac{\partial u_3}{\partial x_l} \frac{\partial u_3}{\partial x_l} \right\} dx_1 \, dx_2 \, dx_3 \\ &= \int_{\Omega} \left\{ - \frac{\partial \tilde{u}_3}{\partial x_3} \left[\left(\frac{\partial u_1}{\partial x_1} \right)^2 + \left(\frac{\partial u_2}{\partial x_2} \right)^2 - \frac{\partial u_1}{\partial x_1} \frac{\partial u_2}{\partial x_2} + \left(\frac{\partial u_1}{\partial x_2} \right)^2 + \left(\frac{\partial u_2}{\partial x_1} \right)^2 + \frac{\partial u_1}{\partial x_2} \frac{\partial u_2}{\partial x_1} \right] \right. \\ &\quad \left. + \sum_{k,l=1}^2 \left(\frac{\partial u_k}{\partial x_l} \frac{\partial \tilde{u}_3}{\partial x_3} \frac{\partial u_k}{\partial x_l} + u_k \frac{\partial \tilde{u}_3}{\partial x_3} \frac{\partial^2 u_k}{\partial x_l \partial x_3} - u_k \frac{\partial \tilde{u}_3}{\partial x_l} \frac{\partial^2 u_k}{\partial x_l \partial x_3} \right) \right. \\ &\quad \left. - \tilde{u}_3 \left[\sum_{j,l=1}^2 \left[\frac{\partial}{\partial x_j} \left(\frac{\partial u_j}{\partial x_l} \frac{\partial u_3}{\partial x_l} \right) + \frac{\partial}{\partial x_l} \left(\frac{\partial u_j}{\partial x_3} \frac{\partial u_j}{\partial x_l} \right) \right] - \sum_{l=1}^2 \frac{\partial}{\partial x_l} \left[\left(\frac{\partial u_1}{\partial x_1} + \frac{\partial u_2}{\partial x_2} \right) \frac{\partial u_3}{\partial x_l} \right] \right] \right. \\ &\quad \left. + \sum_{j,l=1}^2 \frac{\partial \tilde{u}_3}{\partial x_j} \frac{\partial u_j}{\partial x_l} \frac{\partial u_3}{\partial x_l} - \sum_{l=1}^2 \left(\frac{\partial u_1}{\partial x_1} + \frac{\partial u_2}{\partial x_2} \right) \frac{\partial \tilde{u}_3}{\partial x_l} \frac{\partial u_3}{\partial x_l} \right\} dx_1 \, dx_2 \, dx_3. \end{aligned}$$

Then, from the above and the Cauchy–Schwarz inequality we obtain

$$\begin{aligned} \frac{1}{2} \frac{d\|\nabla_h u\|_2^2}{dt} + \nu\|\Delta_h u\|_2^2 + \nu\|\nabla_h u_z\|_2^2 &\leq F\|\Delta_h u\|_2 + C \int_{\Omega} |\tilde{u}_3| |\nabla u| |\nabla_h \nabla u| \, dx_1 \, dx_2 \, dx_3 \\ &\quad + C \int_{\Omega} |\nabla \tilde{u}_3| |\nabla_h u|^2 \, dx_1 \, dx_2 \, dx_3 + C \int_{\Omega} |u| |\nabla \tilde{u}_3| |\nabla_h \nabla u| \, dx_1 \, dx_2 \, dx_3. \end{aligned}$$

where F is given in (21). By applying Lemma 1 we obtain

$$\begin{aligned}
 & \int_{\Omega} |\bar{u}_3| |\nabla u| |\nabla_h \nabla u| dx_1 dx_2 dx_3 \\
 & \leq C \|\bar{u}_3\|_2^{1/2} (\|\bar{u}_3\|_2 + \|\nabla_h \bar{u}_3\|_2)^{1/2} \|\nabla u\|_2^{1/2} (\|\nabla u\|_2 + \|\nabla_h \nabla u\|_2)^{1/2} \|\nabla_h \nabla u\|_2 \\
 & \leq C \|\bar{u}_3\|_2 \|\nabla u\|_2 \|\nabla_h \nabla u\|_2 + C \|\bar{u}_3\|_2^{1/2} \|\nabla_h \bar{u}_3\|_2^{1/2} \|\nabla u\|_2^{1/2} \|\nabla_h \nabla u\|_2^{1/2} \|\nabla_h \nabla u\|_2 \\
 & \leq C \|u\|_2 \|\nabla u\|_2 \|\nabla_h \nabla u\|_2 + C \|u\|_2^{1/2} \|\nabla_h u\|_2^{1/2} \|\nabla u\|_2^{1/2} \|\nabla_h \nabla u\|_2^{3/2}. \tag{24}
 \end{aligned}$$

By Hölder inequality, we reach

$$\begin{aligned}
 \int_{\Omega} |\nabla \tilde{u}_3| |\nabla_h u|^2 dx_1 dx_2 dx_3 & \leq C \|\nabla \tilde{u}_3\|_2 \|\nabla_h u\|_4^2 \\
 & \leq C \|\nabla \tilde{u}_3\|_2 \|\nabla_h u\|_2^{1/2} (\|\nabla_h u\|_2 + \|\nabla_h \nabla u\|_2)^{3/2}. \tag{25}
 \end{aligned}$$

And also by Hölder inequality, we obtain

$$\begin{aligned}
 & \int_{\Omega} |u| |\nabla \tilde{u}_3| |\nabla_h \nabla u| dx_1 dx_2 dx_3 \\
 & \leq C \int_{\mathbb{R}^2} \{ \|u(x_1, x_2, \cdot)\|_{\infty} \|\nabla \tilde{u}_3(x_1, x_2, \cdot)\|_2 \|\nabla_h \nabla u(x_1, x_2, \cdot)\|_2 \} dx_1 dx_2 \\
 & \leq C \int_{\mathbb{R}^2} \left\{ \|u(x_1, x_2, \cdot)\|_2^{1/2} \left\| \frac{\partial u}{\partial x_3}(x_1, x_2, \cdot) \right\|_2^{1/2} \right. \\
 & \quad \left. \|\nabla \tilde{u}_3(x_1, x_2, \cdot)\|_2 \|\nabla_h \nabla u(x_1, x_2, \cdot)\|_2 \right\} dx_1 dx_2 \\
 & \leq C \left\{ \int_{\mathbb{R}^2} \|u(x_1, x_2, \cdot)\|_2^4 dx_1 dx_2 \right\}^{1/8} \left\{ \int_{\mathbb{R}^2} \left\| \frac{\partial u}{\partial x_3}(x_1, x_2, \cdot) \right\|_2^4 dx_1 dx_2 \right\}^{1/8} \\
 & \quad \times \left\{ \int_{\mathbb{R}^2} \|\nabla \tilde{u}_3(x_1, x_2, \cdot)\|_2^4 dx_1 dx_2 \right\}^{1/4} \|\nabla_h \nabla u\|_2 \\
 & \leq C \|u\|_2^{1/4} \|\nabla_h u\|_2^{1/4} \left\| \frac{\partial u}{\partial x_3} \right\|_2^{1/4} \left\| \frac{\partial \nabla_h u}{\partial x_3} \right\|_2^{1/4} \|\nabla \tilde{u}_3\|_2^{1/2} \|\nabla_h \nabla \tilde{u}_3\|_2^{1/2} \|\nabla_h \nabla u\|_2. \tag{26}
 \end{aligned}$$

By (24)–(26) and Young's inequality we get

$$\begin{aligned}
 & \frac{d \|\nabla_h u\|_2^2}{dt} + \nu \|\nabla_h \nabla u\|_2^2 \leq CF^2 + C \|u\|_2^2 \|\nabla u\|_2^2 \\
 & \quad + C \left(\|u\|_2^{1/2} \|\nabla u\|_2^2 + \|\nabla \tilde{u}_3\|_2^4 + \|u\|_2^2 \left\| \frac{\partial u}{\partial x_3} \right\|_2^2 \|\nabla \tilde{u}_3\|_2^4 \right) \|\nabla_h u\|_2^2,
 \end{aligned}$$

where F is given in (21). Thanks to Gronwall inequality, we obtain, for all $t \in [0, T_*)$,

$$\|\nabla_h u(t)\|_2^2 + \nu \int_0^t \|\nabla_h \nabla u(s)\|_2^2 ds \leq K_2,$$

where

$$K_2 = e^{CK_1^2 + C(T+K_1^2)\max_{0 \leq s \leq T} \|\nabla \tilde{u}_3(s)\|_2^4} [\|u_0\|_{H^1(\Omega)}^2 + F^2 + CK_1^2]. \tag{27}$$

Recall that $\|u\|_V^2 = \int u \cdot Au \, dx_1 \, dx_2 \, dx_3$. It is well known that $\|u\|_V^2$ is equivalent to $\|\nabla u\|_2^2$ (see, e.g., Constantin and Foias, 1988). Taking the inner product of the equation (8) with Au in H , we get

$$\begin{aligned} \frac{1}{2} \frac{d\|u\|_V^2}{dt} + \nu \|Au\|_2^2 &= \int_{\Omega} (f - B(u, u)) \cdot Au \, dx_1 \, dx_2 \, dx_3 \\ &\leq F \|Au\|_2 + C \|u\|_6 \|\nabla u\|_3 \|Au\| \\ &\leq F \|Au\|_2 + C \|\nabla_h u\|_2^{2/3} \left\| \frac{\partial u}{\partial x_3} \right\|_2^{1/3} \|\nabla u\|_2^{1/2} \|Au\|^{3/2}. \end{aligned}$$

Here, we used (cf., e.g., Galdi, 1994, p. 33)

$$\|u\|_6 \leq C \left\| \frac{\partial u}{\partial x_1} \right\|_2^{1/3} \left\| \frac{\partial u}{\partial x_2} \right\|_2^{1/3} \left\| \frac{\partial u}{\partial x_3} \right\|_2^{1/3}.$$

By Young’s inequality we obtain

$$\frac{d\|u\|_V^2}{dt} + \nu \|Au\|_2^2 \leq CF^2 + C \|\nabla_h u\|_2^{8/3} \left\| \frac{\partial u}{\partial x_3} \right\|_2^{4/3} \|u\|_V^2.$$

By Gronwall inequality and (27), we obtain, for all $t \in [0, T_*)$,

$$\|u(t)\|_V^2 + \nu \int_0^t \|Au(s)\|_2^2 \, ds \leq K,$$

where

$$K = e^{CK_2^{4/3} K_1^{2/3}} [\|u_0\|_{H^1(\Omega)}^2 + F^2]. \tag{28}$$

Therefore,

$$\limsup_{t \rightarrow T_*^-} \|u(t)\|_V^2 \leq K,$$

which leads to a contradiction that $[0, T_*)$ is the maximal interval of existence, and this completes the proof. \square

Acknowledgments

E.S.T. would like to thank the Bernoulli Center of the École Polytechnique Fédéral de Lausanne for the kind hospitality where this work was completed. This work was supported in part by the NSF grant no. DMS-0504619, the BSF grant no. 2004271, the ISF grant no. 120/06, and by the MAOF Fellowship of the Israeli Council of Higher Education.

References

- Adams, R. A. (1975). *Sobolev Spaces*. New York: Academic Press.
- Berselli, L. C. (2002). On a regularity criterion for the solutions to the 3D Navier–Stokes equations. *Differential Integral Equations* 15:1129–1137.
- Berselli, L. C., Galdi, G. P. (2002). Regularity criteria involving the pressure for the weak solutions to the Navier–Stokes equations. *Proc. Amer. Math. Soc.* 130:3585–3595.
- Cao, C., Titi, E. S. (2003). Global well-posedness and finite dimensional global attractor for a 3-D planetary geostrophic viscous model. *Comm. Pure Appl. Math.* 56:198–233.
- Cao, C., Titi, E. S. (2007). Global well-posedness of the three-dimensional viscous primitive equations of large scale ocean and atmosphere dynamics. To appear in *Annals of Mathematics* 165.
- Cao, C., Titi, E. S. (2007). Pressure regularity criterion for the three-dimensional Navier–Stokes equations in infinite channel. *Annals Mathematics* 166:245–267.
- Chae, D., Lee, J. (2001). Regularity criterion in terms of pressure for the Navier–Stokes equations. *Nonlinear Anal, Ser. A: Theory Methods* 46:727–735.
- Constantin, P., Foias, C. (1988). *Navier–Stokes Equations*. The University of Chicago Press.
- Galdi, G. P. (1994). *An Introduction to the Mathematical Theory of the Navier–Stokes Equations*. Vol. I & II. Springer-Verlag.
- He, C. (2002). New sufficient conditions for regularity of solutions to the Navier–Stokes equations. *Adv. Math. Sci. Appl.* 12:535–548.
- Kato, T. (1984). Strong L^p solutions of the Navier–Stokes equation in R^m , with applications to weak solutions. *Math. Z.* 187:471–480.
- Kukavica, I., Ziane, M. (2006). One component regularity for the Navier–Stokes equation. *Nonlinearity* 19:453–470.
- Ladyzhenskaya, O. A. (1969). *The Mathematical Theory of Viscous Incompressible Flow*. [English transl.], New York, London: Gordon and Breach.
- Ladyzhenskaya, O. A. (1985). *The Boundary Value Problems of Mathematical Physics*. Springer-Verlag.
- Moise, I. (2006). Regularity of Navier–Stokes equations. *Differential Integral Equations* 19:31–50.
- Pokorný, M. (2003). On the result of He concerning the smoothness of solutions to the Navier–Stokes equations. *Electron. J. Differential Equations* 11:1–8.
- Prodi, G. (1959). Un teorema di unicità per le equazioni di Navier–Stokes. [A uniqueness theorem for the Navier–Stokes equations.] *Ann. Mat. Pura Appl.* 48:173–182.
- Serrin, J. (1962). On the interior regularity of weak solutions of the Navier–Stokes equations. *Arch. Rational Mech. Anal.* 9:187–191.
- Serrin, J. (1963). The initial value problem for the Navier–Stokes equations. In: Langer, R. E., ed. *Nonlinear Problems*. Madison: Univ. of Wisconsin Press, pp. 69–98.
- Sohr, H. (2001). *The Navier–Stokes Equations: An Elementary Functional Analytical Approach*. Birkhäuser Advanced Texts. Basel: Birkhäuser Verlag.
- Temam, R. (1983). *Navier-Stokes Equations and Nonlinear Functional Analysis*. CBMS Regional Conference series, No. 41. SIAM, Philadelphia.
- Temam, R. (2001). *Navier-Stokes Equations, Theory and Numerical Analysis*. 3rd revised ed. North-Holland.
- Zhou, Y. (2002). A new regularity criterion for the Navier–Stokes equations in terms of the gradient of one velocity component. *Methods Appl. Anal.* 9:563–578.