

$H(\text{div})$ -conforming finite element tensors with constraints

Long Chen^a, Xuehai Huang^{b,*}

^a Department of Mathematics, University of California at Irvine, Irvine, CA 92697, USA

^b School of Mathematics, Shanghai University of Finance and Economics, Shanghai, 200433, China

ARTICLE INFO

MSC:

65N30

65N12

58J10

15A69

Keywords:

$H(\text{div})$ -conforming finite element

Differential form

Tensors with linear constraints

Geometric decomposition

Bubble space

ABSTRACT

A unified construction of $H(\text{div})$ -conforming finite element tensors, including vector element, symmetric matrix element, traceless matrix element, and, in general, tensors with linear constraints, is developed in this work. It is based on the geometric decomposition of Lagrange elements into bubble functions on each sub-simplex. Each tensor at a sub-simplex is further decomposed into tangential and normal components. The tangential component forms the bubble function space, while the normal component characterizes the trace. Some degrees of freedom can be redistributed to $(n-1)$ -dimensional faces. The developed finite element spaces are $H(\text{div})$ -conforming and satisfy the discrete inf-sup condition. Intrinsic bases of the constraint tensor space are also established.

1. Introduction

Hilbert complexes play a fundamental role in the theoretical analysis and the design of stable numerical methods for partial differential equations [1–5]. Recently, in [6], Arnold and Hu have developed a systematic approach to derive new Hilbert complexes from well-understood differential complexes, such as the de Rham complex. In space \mathbb{R}^n , for $0 \leq k \leq n$, let $\text{Alt}^{k,n-1} := \text{Alt}^k \otimes \text{Alt}^{n-1}$ be the tensor product of alternating multilinear functional spaces, H^s be the standard Sobolev space with real index s , and κ_k be the Koszul operator for the de Rham complex. Below, we rotate the right end column of the Bernstein–Gelfand–Gelfand (BGG) diagram in [6] and switch the ordering of the index in [6] to match the row action of the operator div .

$$\begin{array}{ccccccc}
 H^{q-n} \otimes \text{Alt}^{n,n-1} & \xrightarrow{\kappa_n} & H^{q-n+1} \otimes \text{Alt}^{n-1,n-1} & \xrightarrow{\kappa_{n-1}} & \dots & \xrightarrow{\kappa_2} & H^{q-1} \otimes \text{Alt}^{1,n-1} & \xrightarrow{\kappa_1} & H^q \otimes \text{Alt}^{0,n-1} \\
 \downarrow \text{div} & \searrow \text{id} & \downarrow \text{div} & \searrow \text{skw} & & \searrow \text{div} & \downarrow \text{div} & \searrow \text{tr} & \downarrow \text{div} \\
 H^{q-n-1} \otimes \text{Alt}^{n,n} & \xrightarrow{\kappa_n} & H^{q-n} \otimes \text{Alt}^{n-1,n} & \xrightarrow{\kappa_{n-1}} & \dots & \xrightarrow{\kappa_2} & H^{q-2} \otimes \text{Alt}^{1,n} & \xrightarrow{\kappa_1} & H^{q-1} \otimes \text{Alt}^{0,n}
 \end{array}$$

The algebraic operator $s^{k,n-1} : \text{Alt}^{k,n-1} \rightarrow \text{Alt}^{k-1,n}$, for $1 \leq k \leq n$ along the \searrow direction, is defined as

$$s^{k,n-1} \omega(w_1, \dots, w_{k-1})(v_1, \dots, v_n) := \sum_{i=1}^n (-1)^{i-1} \omega(v_i, w_1, \dots, w_{k-1})(v_1, \dots, \widehat{v}_i, \dots, v_n) \quad \forall v_1, \dots, v_n, w_1, \dots, w_{k-1} \in \mathbb{R}^n.$$

The tensor space \mathbb{X} is defined as

$$\mathbb{X} := \ker(s^{k,n-1}) \cap \text{Alt}^{k,n-1}, \quad 1 \leq k \leq n-1.$$

* Corresponding author.

E-mail addresses: chenlong@math.uci.edu (L. Chen), huang.xuehai@sufe.edu.cn (X. Huang).

A tensor $\omega \in \text{Alt}^{k,n-1}$ can be represented by a matrix $\mathbf{A} = (a_{\sigma,i}) \in \mathbb{R}^{\binom{n}{k} \times n}$, which is called a proxy matrix, and then \mathbb{X} is a subset of matrices satisfying certain linear constraints. For simplicity, we will use matrix \mathbf{A} to represent an element in \mathbb{X} . Given a domain $\Omega \subset \mathbb{R}^n$, define $H(\text{div}, \Omega; \mathbb{X})$ space as

$$H(\text{div}, \Omega; \mathbb{X}) = \{ \mathbf{A} \in L^2(\Omega; \mathbb{X}) : \text{div } \mathbf{A} \in L^2(\Omega; \mathbb{R}^{\binom{n}{k}}) \},$$

where the divergence operator is applied to each row of \mathbf{A} . Notable examples are $H(\text{div}; \mathbb{S})$ with the symmetric matrix \mathbb{S} , which plays an important role in the discretization of the elasticity equation in the mixed form [7–9], and $H(\text{div}; \mathbb{T})$ with the traceless matrix \mathbb{T} , which is used in the mixed form of the linearized Einstein-Bianchi system [10–12].

Given a simplicial mesh \mathcal{T}_h of Ω , we shall develop a systematic construction of $H(\text{div})$ -conforming finite element subspaces of $H(\text{div}, \Omega; \mathbb{X})$ space. Take a proxy matrix \mathbf{A} for an elementwise smooth tensor $\omega \in L^2(\Omega; \mathbb{X})$. $H(\text{div})$ -conformity means $\mathbf{A} \mathbf{n}_F$ is continuous across each $(n-1)$ -dimensional face F of \mathcal{T}_h . Namely, each row of \mathbf{A} is an $H(\text{div})$ -conforming vector function.

It is the constraint $s^{k,n-1}(\mathbf{A}) = 0$ that makes the finite element construction difficult, as the constraint and the normal continuity should be satisfied simultaneously. For example, the symmetry of the tensors makes it a challenging problem to construct $H(\text{div}; \mathbb{S})$ -conforming finite elements. Arnold and Winther [9] constructed such an element in two dimensions, and later it was extended to higher dimensions in [13–15]. Hu and Zhang [16,17] constructed $H(\text{div}; \mathbb{S})$ -conforming finite elements with full polynomial spaces with matched order of approximation.

The approach we shall use is the geometric decomposition, which leads to explicit bases for finite elements. The geometric decomposition is an important tool for finite element analysis. For example, it is used in [18] to construct a local and bounded co-chain projection to the discrete de Rham complexes. The finite element system in [19] also originates from the geometric decomposition. The geometric decomposition of standard finite element de Rham complexes is well-studied in [1,20,21], and in [22] for nodal finite element de Rham complexes. Recently geometric decomposition has been extended to smooth finite elements and smooth finite element de Rham and Stokes complexes [23,24].

We shall integrate the geometric decomposition of the Lagrange element with tangential-normal (t - n) decompositions $\mathbb{R}^n = \mathcal{T}^f \oplus \mathcal{N}^f$ on subsimplices to produce geometric decompositions of $H(\text{div})$ -conforming finite element vectors and tensors, exhibiting normal continuity across all $(n-1)$ -dimensional faces. In a similar way, hierarchical geometric decomposition of $H(\text{div})$ -conforming finite elements in two and three dimensions is discussed in [25–27]. A geometric decomposition of $H(\text{div})$ -conforming finite element vectors with a different tangential-normal decomposition in three dimensions is also shown in [22]. While these prior studies offer similar insights, our methodology introduces a novel level of generality. A significant aspect of our contribution is the expansion of geometric decomposition techniques to effectively manage tensors subjected to specific constraints.

To satisfy the constraint while still keeping normal continuity, the crucial step is to get a t - n decomposition of \mathbb{X} with respect to the second component in $\text{Alt}^{k,n-1}$, i.e.,

$$\mathbb{X} = \mathcal{T}^f(\mathbb{X}) \oplus \mathcal{N}^f(\mathbb{X}),$$

where $\mathcal{T}^f(\mathbb{X}) = (\text{Alt}^k \otimes \mathcal{T}^f) \cap \ker(s^{k,n-1})$ and $\mathcal{N}^f(\mathbb{X}) = \pi_{\mathbb{X}}(\text{Alt}^k \otimes \mathcal{N}^f)$ with an oblique (non-orthogonal) projection operator $\pi_{\mathbb{X}} : \text{Alt}^k \otimes \mathcal{N}^f \rightarrow \mathbb{X}$ so that one constraint is used only once either in $\mathcal{T}^f(\mathbb{X})$ or $\mathcal{N}^f(\mathbb{X})$.

This induces the geometric decomposition

$$\mathbb{P}_r(T; \mathbb{X}) = \bigoplus_{\ell=0}^n \bigoplus_{f \in \Delta_\ell(T)} b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes [\mathcal{T}^f(\mathbb{X}) \oplus \mathcal{N}^f(\mathbb{X})]. \quad (1)$$

The tangential component will contribute to the polynomial bubble space

$$\mathbb{B}_r(\text{div}, T; \mathbb{X}) := \bigoplus_{\ell=1}^n \bigoplus_{f \in \Delta_\ell(T)} [b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{T}^f(\mathbb{X})],$$

and the normal component $b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{N}^f(\mathbb{X})$ to the trace.

As a direct result of decomposition (1), the following degrees of freedom (DoFs)

$$\omega(v_i), \quad i = 0, \dots, n, \omega \in \mathbb{X}, \quad (2a)$$

$$(\omega, \eta)_f, \quad \eta \in \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{N}^f(\mathbb{X}), \quad f \in \Delta_\ell(\mathcal{T}_h), \quad \ell = 1, \dots, n-1, \quad (2b)$$

$$(\omega, \eta)_T, \quad \eta \in \mathbb{B}_r(\text{div}, T; \mathbb{X}), \quad T \in \mathcal{T}_h, \quad (2c)$$

will determine a space $V_h \subset H(\text{div}, \Omega; \mathbb{X})$. Here we single out the vertex DoFs to emphasize the finite element function is continuous on vertices.

Discrete inf-sup condition will be established with requirement $r \geq n+1$ and with modification of DoFs for $r \geq k+1$ for $1 \leq k \leq n-2$. Variants can be constructed by further tuning DoFs (2), which will recover the existing $H(\text{div}; \mathbb{S})$ elements [16,17,28] and $H(\text{div}; \mathbb{T})$ elements [11,12].

The geometric decomposition approach in this paper is not easy to extend to the case $\mathbb{X} = \mathbb{S} \cap \mathbb{T}$, which requires special care and super-smoothness of DoFs; see the recent work [29].

The rest of this paper is organized as follows. Section 2 covers foundational concepts, including simplex, barycentric coordinates, Bernstein polynomials, t - n bases, and a geometric decomposition of Lagrange elements. Sections 3 and 4 explore the geometric decompositions of vector face elements and matrix face elements, respectively. Section 5 focuses on the constraint tensor space \mathbb{X} and its bases. The geometric decomposition of $H(\text{div})$ -conforming tensors is developed in Section 6. As the language of differential form is abstract, in the first few sections we present the results using vector and matrix language and then move to the differential forms in Sections 5 and 6.

2. Notation and background

We summarize the most important notation and integer indices in the beginning:

- \mathbb{R}^n : n is the dimension of the ambient Euclidean space and $n \geq 2$;
- \mathbb{P}_r : r is the degree of the polynomial and $r \geq 0$;
- Λ^k : k is the order of the differential form and $0 \leq k \leq n$;
- $\Delta_\ell(T)$: ℓ is the dimension of a sub-simplex $f \in \Delta_\ell(T)$ and $0 \leq \ell \leq n$.

2.1. Simplex and sub-simplices

Let $T \subset \mathbb{R}^n$ be an n -dimensional simplex with vertices v_0, v_1, \dots, v_n in general position. Following [20], we let $\Delta(T)$ denote all the subsimplices of T , while $\Delta_\ell(T)$ denotes the set of subsimplices of dimension ℓ , for $0 \leq \ell \leq n$.

For a sub-simplex $f \in \Delta_\ell(T)$, we will overload the notation f for both the geometric simplex and the algebraic set of indices. Namely on one hand $f = \{f(0), \dots, f(\ell)\} \subseteq \{0, 1, \dots, n\}$, and on the other hand

$$f = \text{Convex}(v_{f(0)}, \dots, v_{f(\ell)}) \in \Delta_\ell(T)$$

is the ℓ -dimensional simplex spanned by the vertices $v_{f(0)}, \dots, v_{f(\ell)}$. If $f \in \Delta_\ell(T)$, for $\ell = 0, \dots, n-1$, then $f^* \in \Delta_{n-\ell-1}(T)$ denotes the sub-simplex of T opposite to f . Algebraically treating f as a subset of $\{0, 1, \dots, n\}$, $f^* \subseteq \{0, 1, \dots, n\}$ so that $f \cup f^* = \{0, 1, \dots, n\}$, i.e., f^* is the complement of set f . Geometrically,

$$f^* = \text{Convex}(v_{f^*(1)}, \dots, v_{f^*(n-\ell)}) \in \Delta_{n-\ell-1}(T)$$

is the $(n-\ell-1)$ -dimensional simplex spanned by vertices not in f . We refer to [24, Fig. 2] for an illustration of f and f^* .

Denote by F_i the $(n-1)$ -dimensional face opposite to vertex v_i , i.e., $F_i = \{i\}^*$. Here capital F is reserved for an $(n-1)$ -dimensional face of T . For lower dimensional sub-simplices, we sometimes use more conventional notation. For example, the vertex will be denoted by v_i and the edge formed by v_i and v_j will be denoted by e_{ij} .

2.2. Tangential-normal (t - n) bases

For an ℓ -dimensional sub-simplex $f \in \Delta_\ell(T)$, choose ℓ linearly independent tangential vectors $\{t_1^f, \dots, t_\ell^f\}$ of f and $n-\ell$ linearly independent normal vectors $\{n_1^f, \dots, n_{n-\ell}^f\}$ of f . The set of n vectors $\{t_1^f, \dots, t_\ell^f, n_1^f, \dots, n_{n-\ell}^f\}$ forms a basis of \mathbb{R}^n . Notice that for $\ell = 0$, i.e., at vertices, there are no tangential vectors, and for $\ell = n$, there are no normal vectors. Define the tangent plane and normal plane of f as

$$\mathcal{T}^f := \text{span}\{t_i^f, i = 1, \dots, \ell\}, \quad \mathcal{N}^f := \text{span}\{n_i^f, i = 1, \dots, n-\ell\}.$$

All vectors are normalized but $\{t_i^f\}$ or $\{n_i^f\}$ may not form an orthonormal basis.

Inside the subspace \mathcal{T}^f , we can find a basis $\{\hat{t}_1^f, \dots, \hat{t}_\ell^f\}$ dual to $\{t_1^f, \dots, t_\ell^f\}$, i.e., $\hat{t}_i \in \mathcal{T}^f$ and $(\hat{t}_i, t_j) = \delta_{i,j}$ for $i, j = 1, \dots, \ell$. Similarly we have a basis $\{\hat{n}_1^f, \dots, \hat{n}_{n-\ell}^f\}$ of \mathcal{N}^f and $(\hat{n}_i, n_j) = \delta_{i,j}$ for $i, j = 1, \dots, n-\ell$. As $\mathcal{T}^f \perp \mathcal{N}^f$, the basis $\{\hat{t}_1^f, \dots, \hat{t}_\ell^f, \hat{n}_1^f, \dots, \hat{n}_{n-\ell}^f\}$ is also dual to $\{t_1^f, \dots, t_\ell^f, n_1^f, \dots, n_{n-\ell}^f\}$.

Given a sub-simplex $f \in \Delta_\ell(T)$, we now present two bases for its normal plane \mathcal{N}^f constructed in [30].

Recall that we label F_i as the $(n-1)$ -dimensional face opposite to the i th vertex. Then $f \subseteq F_i$ for $i \in f^*$. One basis is composed by unit normal vectors of all such $(n-1)$ -dimensional faces:

$$\{n_{F_i}, i \in f^*\},$$

and will be called the *face normal basis*.

We now give its dual basis in \mathcal{N}^f . For $f \in \Delta_\ell(T)$, $\ell = 0, 1, \dots, n-1$ and $i \in f^*$, let $f \cup \{i\}$ denotes the $(\ell+1)$ -dimensional face in $\Delta_{\ell+1}(T)$ with vertices $\{i, f(0), \dots, f(\ell)\}$. Let $n_{f \cup \{i\}}^f$ be a unit normal vector of f but tangential to $f \cup \{i\}$. The basis

$$\{n_{f \cup \{i\}}^f, i \in f^*\}$$

will be called the *tangential normal basis*.

Lemma 2.1. For $f \in \Delta_\ell(T)$, the rescaled tangential normal basis $\{n_{f \cup \{i\}}^f / (n_{f \cup \{i\}}^f \cdot n_{F_i}), i \in f^*\}$ of \mathcal{N}^f is dual to the face normal basis $\{n_{F_i}, i \in f^*\}$.

Proof. Clearly $n_{f \cup \{i\}}^f, n_{F_i} \in \mathcal{N}^f$ for $i \in f^*$. It suffices to prove

$$n_{f \cup \{i\}}^f \cdot n_{F_j} = 0 \quad \text{for } i, j \in f^*, i \neq j,$$

which follows from the fact $f \cup \{i\} \subseteq F_j$ and $n_{f \cup \{i\}}^f \in \mathcal{T}^{f \cup \{i\}}$. \square




Fig. 1. Face normal basis and tangential normal basis of a vertex and an edge in a tetrahedron.

Example 2.2. An important example is $f \in \Delta_0(T)$, i.e., f is a vertex. Without loss of generality, let $f = \{0\}$. Then $n_{f \cup \{i\}}^f$ is a unit normal vector of edge $\{0, i\}$: t_{0i} or t_{i0} depending on the orientation. Its dual basis is $\{n_{F_i}/(n_{F_i} \cdot t_{0i}), i = 1, \dots, n\}$. See Fig. 1(a).

Example 2.3. Let $f = \{0, 1\}$ be an edge of a tetrahedron. Then we have two bases of the normal plane \mathcal{N}^f : $\{n_{F_2}, n_{F_3}\}$ and $\{n_{\{0,1,2\}}^{\{0,1\}}, n_{\{0,1,3\}}^{\{0,1\}}\}$. They are dual to each other with an appropriate rescaling. See Fig. 1(b).

2.3. Barycentric coordinates and Bernstein polynomials

For a domain $\Omega \subseteq \mathbb{R}^n$ and integer $r \geq 0$, $\mathbb{P}_r(\Omega)$ denotes the space of real valued polynomials defined on Ω of degree less than or equal to r . For simplicity, we let $\mathbb{P}_r = \mathbb{P}_r(\mathbb{R}^n)$. Hence, if n -dimensional domain Ω has nonempty interior, then $\dim \mathbb{P}_r(\Omega) = \dim \mathbb{P}_r = \binom{r+n}{n}$. When $\Omega = \{v\}$ is a point, $\mathbb{P}_r(v) = \mathbb{R}$ for all $r \geq 0$. And we set $\mathbb{P}_r(\Omega) = \{0\}$ when $r < 0$. Let $\mathbb{H}_r(\Omega)$ denote the space of real valued polynomials defined on Ω of degree r .

For n -dimensional simplex T , we denote by $\lambda_0, \lambda_1, \dots, \lambda_n$ the barycentric coordinate functions with respect to T . That is $\lambda_i \in \mathbb{P}_1(T)$ and $\lambda_i(v_j) = \delta_{ij}$, $0 \leq i, j \leq n$, where δ_{ij} is the Kronecker delta function. The functions $\{\lambda_i, i = 0, 1, \dots, n\}$ form a basis for $\mathbb{P}_1(T)$, $\sum_{i=0}^n \lambda_i(x) = 1$, and $0 \leq \lambda_i(x) \leq 1, i = 0, 1, \dots, n$, for $x \in T$. The sub-simplices of T correspond to the zero sets of the barycentric coordinates. Indeed $f = \{x \in T \mid \lambda_i(x) = 0, i \in f^*\}$ for $f \in \Delta_\ell(T)$.

We will use the multi-index notation $\alpha \in \mathbb{N}^n$, meaning $\alpha = (\alpha_1, \dots, \alpha_n)$ with integer $\alpha_i \geq 0$. We define $x^\alpha = x_1^{\alpha_1} \dots x_n^{\alpha_n}$, and $|\alpha| := \sum_{i=1}^n \alpha_i$. We will also use the set $\mathbb{N}^{0:n}$ of multi-indices $\alpha = (\alpha_0, \dots, \alpha_n)$, with $\lambda^\alpha := \lambda_0^{\alpha_0} \dots \lambda_n^{\alpha_n}$ for $\alpha \in \mathbb{N}^{0:n}$.

We introduce the simplicial lattice [23,24], which is also known as the principal lattice [31]. A simplicial lattice of degree r and dimension n is a multi-index set of $n+1$ components and with fixed length r , i.e.,

$$\mathbb{T}_r^n = \{\alpha = (\alpha_0, \alpha_1, \dots, \alpha_n) \in \mathbb{N}^{0:n} \mid \alpha_0 + \alpha_1 + \dots + \alpha_n = r\}.$$

An element $\alpha \in \mathbb{T}_r^n$ is called a node of the lattice. The Bernstein representation of polynomial of degree r on a simplex T is

$$\mathbb{P}_r(T) := \text{span}\{\lambda^\alpha = \lambda_0^{\alpha_0} \lambda_1^{\alpha_1} \dots \lambda_n^{\alpha_n}, \alpha \in \mathbb{T}_r^n\}.$$

In the Bernstein form, for an $f \in \Delta_\ell(T)$,

$$\mathbb{P}_r(f) = \text{span}\{\lambda_f^\alpha = \lambda_{f(0)}^{\alpha_0} \lambda_{f(1)}^{\alpha_1} \dots \lambda_{f(\ell)}^{\alpha_\ell}, \alpha \in \mathbb{T}_r^\ell\}.$$

Through the natural extension defined by the barycentric coordinate, $\mathbb{P}_r(f) \subseteq \mathbb{P}_r(T)$. The bubble polynomial of f is a polynomial of degree $\ell+1$:

$$b_f := \lambda_f = \lambda_{f(0)} \lambda_{f(1)} \dots \lambda_{f(\ell)} \in \mathbb{P}_{\ell+1}(f).$$

We have the following property of the bubble polynomial b_f .

Lemma 2.4. Let $f, e \in \Delta(T)$. If $f \not\subseteq e$, then $b_f|_e = 0$.

Proof. As $f = (f \cap e^*) \cup (f \cap e)$ and $f \not\subseteq e$, we conclude $f \cap e^* \neq \emptyset$. So b_f contains λ_i for some $i \in e^*$ and consequently $b_f|_e = 0$. \square

In particular, b_f vanishes at all sub-simplices other than f with dimensions $\leq \dim f$, and higher dimensional sub-simplices not containing f .

2.4. Geometric decomposition of Lagrange elements

We begin with the geometric decomposition of the Lagrange element. The following geometric decomposition of Lagrange element is given in [20] without proofs. As it is the foundation of other geometric decompositions in later sections, we present it using our notation and provide a detailed proof. We refer to [20, Fig. 2.1] for an illustration of this geometric decomposition. For the 0-dimensional face, i.e., a vertex v , we understand that $\int_v u \, ds = u(v)$ for $1 \in \mathbb{P}_r(v) = \mathbb{R}$.

Theorem 2.5 (Geometric Decomposition of Lagrange Element, (2.6) in [20]). *For the polynomial space $\mathbb{P}_r(T)$ with $r \geq 1$ on an n -dimensional simplex T , we have the following decomposition*

$$\mathbb{P}_r(T) = \bigoplus_{\ell=0}^n \bigoplus_{f \in \Delta_\ell(T)} b_f \mathbb{P}_{r-(\ell+1)}(f). \quad (3)$$

And the function $u \in \mathbb{P}_r(T)$ is uniquely determined by degrees of freedom (DoFs)

$$\int_f u p \, ds, \quad p \in \mathbb{P}_{r-(\ell+1)}(f), f \in \Delta_\ell(T), \ell = 0, 1, \dots, n. \quad (4)$$

Proof. We first prove the decomposition (3). Each component $b_f \mathbb{P}_{r-(\ell+1)}(f) \subset \mathbb{P}_r(T)$ and the sum is direct due to the property of b_f , cf. Lemma 2.4. Then count the dimensions and use the combinatorial identity

$$\sum_{\ell=0}^n \binom{n+1}{\ell+1} \binom{r-1}{r-\ell-1} = \binom{n+r}{r},$$

which can be proved by looking at the coefficient of x^r in $(1+x)^{n+1}(1+x)^{r-1} = (1+x)^{n+r}$.

To prove the unisolvence, we choose a basis $\{\phi_i\}$ of $\mathbb{P}_r(T)$ by the decomposition (3) and denote DoFs (4) as $\{N_i\}$. By construction, the dimension of $\{\phi_i\}$ matches the number of DoFs $\{N_i\}$. The DoF-Basis matrix $(N_i(\phi_j))$ is thus square and block lower triangular in the sense that for $\phi_f \in b_f \mathbb{P}_{r-(\ell+1)}(f)$,

$$\int_e \phi_f p \, ds = 0, \quad e \in \Delta_m(T) \text{ with } m \leq \ell \text{ and } e \neq f, p \in \mathbb{P}_{r-\dim e-1}(e)$$

due to the property of b_f established in Lemma 2.4. Each diagonal block matrix is the Gram matrix

$$\int_f p q b_f \, dx_f, \quad p, q \in \mathbb{P}_{r-(\ell+1)}(f),$$

in the measure $b_f \, dx_f$ and thus symmetric and positive definite. In particular, it is invertible. So the unisolvence follows from the invertibility of this lower triangular matrix; see below for an illustration.

$$\begin{array}{c|cccccc} N_f \setminus \phi_f & 0 & 1 & \dots & n-1 & n \\ \hline 0 & \boxed{} & 0 & \dots & 0 & 0 \\ 1 & \boxed{} & \boxed{} & \dots & 0 & 0 \\ \vdots & \vdots & \vdots & \ddots & \vdots & \vdots \\ n-1 & \boxed{} & \boxed{} & \dots & \boxed{} & 0 \\ n & \boxed{} & \boxed{} & \dots & \boxed{} & \boxed{} \end{array} \quad (5)$$

□

Remark 2.6. It is important to note that $\mathbb{P}_{r-(\ell+1)}(f) = \{0\}$ when $r < \ell + 1$. As a result, the last non-zero term in (3) corresponds to $\ell \leq \min\{r-1, n\}$. This implies that the degree of the polynomial dictates the dimension of the sub-simplex in decomposition (3). For instance, with quadratic polynomials, the summation includes only edge bubbles and excludes face bubbles and higher dimensions. Despite this, the full summation notation $\bigoplus_{\ell=0}^n$ is retained for simplicity, with the implicit understanding that the range of non-zero sub-spaces will automatically truncate the limits.

Let $\{\mathcal{T}_h\}$ be a family of partitions of Ω into nonoverlapping simplices with $h_T := \text{diam}(T)$ and $h := \max_{T \in \mathcal{T}_h} h_T$. The mesh \mathcal{T}_h is conforming in the sense that the intersection of any two simplices is either empty or a common lower sub-simplex. Let $\Delta_\ell(\mathcal{T}_h)$ be the set of all ℓ -dimensional sub-simplices of the partition \mathcal{T}_h for $\ell = 0, 1, \dots, n$. The Lagrange finite element space

$$S_h^r := \{v \in C(\Omega) : v|_T \in \mathbb{P}_r(T), \forall T \in \mathcal{T}_h, \text{ DoFs (4) are single-valued}\},$$

has the geometric decomposition

$$S_h^r = \bigoplus_{\ell=0}^n \bigoplus_{f \in \Delta_\ell(\mathcal{T}_h)} b_f \mathbb{P}_{r-(\ell+1)}(f).$$

Here we extend the polynomial $b_f \mathbb{P}_{r-(\ell+1)}(f)$ to each element T containing f by the Bernstein form in the barycentric coordinate and thus it is a piecewise polynomial function and continuous in Ω . Consequently $S_h^r \subset H^1(\Omega)$ and the dimension of

S_h^r is

$$\dim S_h^r = \sum_{\ell=0}^n |\Delta_\ell(\mathcal{T}_h)| \binom{r-1}{\ell},$$

where $|\Delta_\ell(\mathcal{T}_h)|$ is the cardinality, i.e., the number of ℓ -dimensional simplices in \mathcal{T}_h .

The geometric decomposition of the vector Lagrange elements is a straightforward generalization:

$$\mathbb{P}_r(T; \mathbb{R}^n) = \bigoplus_{\ell=0}^n \bigoplus_{f \in \Delta_\ell(T)} [b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathbb{R}^n]. \quad (6)$$

In (6), a fixed orthonormal basis of \mathbb{R}^n is implicitly assumed in which the vector is expanded. It is usually the Cartesian coordinate describing the domain Ω .

3. Geometric decompositions of vector face elements

In this section we consider $H(\text{div})$ -conforming vector finite elements with local shape function space $\mathbb{P}_r(T; \mathbb{R}^n)$. Define $H(\text{div}, \Omega) := \{v \in L^2(\Omega; \mathbb{R}^n) : \text{div } v \in L^2(\Omega)\}$. For a subdomain $K \subseteq \Omega$, the trace operator for the div operator is

$$\text{tr}_K^{\text{div}} v = v \cdot n|_{\partial K} \quad \text{for } v \in C(K),$$

where n denotes the outwards unit normal vector of ∂K . Given a triangulation \mathcal{T}_h and a piecewise smooth function u , it is well known that $u \in H(\text{div}, \Omega)$ if and only if $n_F \cdot u$ is continuous across all faces $F \in \Delta_{n-1}(\mathcal{T}_h)$, which can be ensured by having DoFs on faces. An $H(\text{div})$ -conforming finite element is thus also called a face element.

3.1. Examples in three dimensions

We first use three-dimensional examples to illustrate the main idea. Recall that the geometric decomposition of the vector Lagrange elements in three dimensions reads

$$\mathbb{P}_r(T; \mathbb{R}^3) = \bigoplus_{\ell=0}^3 \bigoplus_{f \in \Delta_\ell(T)} [b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathbb{R}^3]. \quad (7)$$

A fixed orthonormal basis $\{e_i\}_{i=1}^3$ of \mathbb{R}^3 is used in (7). See Fig. 2(a).

An $H(\text{div})$ function is a vector proxy of an $(n-1)$ -form; see Section 5. As a differential form, it is an intrinsic quantity independent of the choice of coordinates/frames. Based on this observation, we shall choose different frames at different sub-simplex $f \in \Delta_\ell(T)$.

For $f \in \Delta_\ell(T)$ with $\ell = 0, 1, 2, 3$, the tangent plane and normal plane of f are

$$\mathcal{T}^f = \text{span}\{t_i^f, i = 1, \dots, \ell\}, \quad \mathcal{N}^f = \text{span}\{n_i^f, i = 1, \dots, 3 - \ell\}.$$

Then \mathbb{R}^3 admits a tangential-normal (t - n) decomposition $\mathbb{R}^3 = \mathcal{T}^f \oplus \mathcal{N}^f$. Coupled with the bubble polynomials, we obtain a t - n decomposition of $\mathbb{P}_r(T; \mathbb{R}^3)$ as

$$\mathbb{P}_r(T; \mathbb{R}^3) = \bigoplus_{\ell=0}^3 \bigoplus_{f \in \Delta_\ell(T)} [\mathbb{B}_r \mathcal{T}^f \oplus \mathbb{B}_r \mathcal{N}^f],$$

where

$$\mathbb{B}_r \mathcal{T}^f = b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{T}^f, \quad \mathbb{B}_r \mathcal{N}^f = b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{N}^f.$$

Notice that for a vertex $v \in \Delta_0(T)$, $\mathbb{B}_r \mathcal{T}^v = \{0\}$, and $\mathbb{B}_r \mathcal{N}^v = \lambda_v \mathbb{R}^3$ as $b_v = \lambda_v$, $\mathbb{P}_{r-1}(v) = \mathbb{R}$, and $\mathcal{N}^v = \mathbb{R}^3$.

Define the polynomial bubble space $\mathbb{B}_r(\text{div}, T) := \ker(\text{tr}_T^{\text{div}}) \cap \mathbb{P}_r(T; \mathbb{R}^3)$. The tangential component will form the div bubble space: for $r \geq 2$, it holds that

$$\mathbb{B}_r(\text{div}, T) = \bigoplus_{\ell=1}^3 \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_r \mathcal{T}^f.$$

Verification of $\mathbb{B}_r \mathcal{T}^f \subseteq \mathbb{B}_r(\text{div}, T)$ is straightforward. For face F not containing f , $b_f|_F = 0$. For face F containing f , $\text{tr}_F^{\text{div}} u|_F = u \cdot n_F = 0$ as $t_i^f \cdot n_F = 0$.

The normal component will contribute to the trace. We can derive the second family of Nédélec element [32,33], which generalizes the Brezzi–Douglas–Marini (BDM) element [34] in two dimensions, from a special t - n basis. For face $F \in \Delta_2(T)$, choose two linearly independent tangential vectors t_1^F, t_2^F of F and one normal vector n_F of F . For edge $e \in \Delta_1(T)$, choose a tangential vector t^e of e and the face normal basis vectors $\{n_{F_1}, n_{F_2}\}$, where F_1, F_2 are two faces containing e . For vertex $v \in \Delta_0(T)$, choose $\{n_{F_1}, n_{F_2}, n_{F_3}\}$, where F_1, F_2, F_3 are three faces containing vertex v . For tetrahedron T , we keep the canonical basis $\{e_i\}_{i=1}^3$, which is considered as tangential vectors of T . See Fig. 2(b).

We then group normal components $\{\mathbb{B}_r \mathcal{N}^f, f \in \Delta(T)\}$ face by face. On each face F , again by the geometric decomposition of Lagrange element, the normal components at different sub-simplices $f \in \Delta(F)$ will determine the scalar function $u \cdot n_F$. In Fig. 2(c), we show such face-wise redistribution for a quadratic face element.




Fig. 2. Classical face elements can be obtained by a special t - n decomposition and face-wise redistribution of normal components.

Example 3.1 (Nédélec Element/BDM Element). Taking $\mathcal{N}^f := \text{span}\{n_{F_i}, f \subset F_i\}$, the following DoFs

$$(v \cdot n_{F_i})|_{F_i}(v), \quad F_i \in \Delta_2(T), v \in \Delta_0(F_i), \quad (8a)$$

$$\int_e (v \cdot n_{F_i})|_{F_i} p \, ds, \quad F_i \in \Delta_2(T), e \in \Delta_1(F_i), p \in \mathbb{P}_{r-2}(e), \quad (8b)$$

$$\int_{F_i} (v \cdot n_{F_i}) p \, ds, \quad F_i \in \Delta_2(T), p \in \mathbb{P}_{r-3}(F_i), \quad (8c)$$

$$\int_T v \cdot p \, dx, \quad p \in \mathbb{B}_r(\text{div}, T) \quad (8d)$$

define the Nédélec element/BDM element. DoFs (8a)–(8c) are all located on face F_i , and uniquely determine $(v \cdot n_{F_i})|_{F_i} \in \mathbb{P}_r(F_i)$. Hence DoFs (8a)–(8c) can be combined to one DoF

$$\int_{F_i} (v \cdot n_{F_i}) p \, ds, \quad F_i \in \Delta_2(T), p \in \mathbb{P}_r(F_i).$$

The interior DoF (8d) can be further replaced by $p \in (\mathbb{P}_{r-2}(T; \mathbb{R}^n) \oplus \mathbb{H}_{r-2}(T; \mathbb{K})x)$ with \mathbb{K} being the skew-symmetric matrix space; see [28]. Therefore DoFs (8) induce the Nédélec/BDM element; see Fig. 3(a).

Different $H(\text{div})$ -conforming finite elements can be obtained by different t - n basis.

Example 3.2 (Stenberg Element). Taking $\{n_i^v\}_{i=1}^3 = \{e_i\}_{i=1}^3$ and $\{n_i^e\}_{i=1}^2 = \{n_{F_i}\}_{i=1}^2$ as two face F_i sharing e , the following DoFs

$$\begin{aligned} v(v), \quad v \in \Delta_0(T), \\ \int_e (v \cdot n_{F_i})|_{F_i} p \, ds, \quad F_i \in \Delta_2(T), e \in \Delta_1(F_i), p \in \mathbb{P}_{r-2}(e), \\ \int_F v \cdot n_F p \, ds, \quad F \in \Delta_2(T), p \in \mathbb{P}_{r-3}(F), \\ \int_T v \cdot p \, dx, \quad p \in \mathbb{B}_r(\text{div}, T) \end{aligned}$$

induce the Stenberg element [35], which is continuous at vertices; see Fig. 3(b).

Example 3.3 (Christiansen–Hu–Hu Element). Taking $\{n_i^v\}_{i=1}^3 = \{e_i\}_{i=1}^3$ and $\{n_i^e\}_{i=1}^2$ depending only on e , the following DoFs

$$\begin{aligned} v(v), \quad v \in \Delta_0(T), \\ \int_e (v \cdot n_i^e) p \, ds, \quad e \in \Delta_1(T), p \in \mathbb{P}_{r-2}(e), i = 1, 2, \\ \int_F (v \cdot n_F) p \, ds, \quad F \in \Delta_2(T), p \in \mathbb{P}_{r-3}(F), \\ \int_T v \cdot p \, dx, \quad p \in \mathbb{B}_r(\text{div}, T) \end{aligned}$$

define the Christiansen–Hu–Hu (CHH) face element [22, Section 3.5], which has extra continuity at vertices and on the normal planes of edges; see Fig. 3(c).




Fig. 3. Different $H(\text{div})$ -conforming finite elements can be obtained by different t - n decompositions.

3.2. Face elements in arbitrary dimension

We now present and prove the result in arbitrary $n \geq 2$ dimensions. Again the main idea is using different and in general non-orthonormal t - n decomposition adapted to different sub-simplices.

3.2.1. Bubble spaces

For an ℓ -dimensional sub-simplex $f \in \Delta_\ell(T)$, choose ℓ linearly independent tangential vectors $\{t_1^f, \dots, t_\ell^f\}$ of f and $n - \ell$ linearly independent normal vectors $\{n_1^f, \dots, n_{n-\ell}^f\}$ of f . The set of n vectors $\{t_1^f, \dots, t_\ell^f, n_1^f, \dots, n_{n-\ell}^f\}$ forms a basis of \mathbb{R}^n . Notice that for $\ell = 0$, i.e., at vertices, there are no tangential vectors and for $\ell = n$, there are no normal vectors. Define the tangent plane and normal plane of f as

$$\mathcal{T}^f := \text{span}\{t_i^f, i = 1, \dots, \ell\}, \quad \mathcal{N}^f := \text{span}\{n_i^f, i = 1, \dots, n - \ell\}.$$

All vectors are normalized but $\{t_i^f\}$ or $\{n_i^f\}$ may not be an orthonormal basis. Coupled with the bubble polynomial, for $r \geq 1$, define

$$\mathbb{B}_r \mathcal{T}^f = b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{T}^f, \quad \mathbb{B}_r \mathcal{N}^f = b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{N}^f.$$

Lemma 3.4. *The shape function space $\mathbb{P}_r(T; \mathbb{R}^n)$ has a geometric decomposition*

$$\mathbb{P}_r(T; \mathbb{R}^n) = \bigoplus_{\ell=0}^n \bigoplus_{f \in \Delta_\ell(T)} [\mathbb{B}_r \mathcal{T}^f \oplus \mathbb{B}_r \mathcal{N}^f]. \quad (9)$$

A function $u \in \mathbb{P}_r(T; \mathbb{R}^n)$ is uniquely determined by the DoFs: for all $f \in \Delta_\ell(T)$, $\ell = 0, 1, \dots, n$,

$$\int_f (u \cdot t_i^f) p \, ds, \quad i = 1, \dots, \ell, p \in \mathbb{P}_{r-(\ell+1)}(f), \quad (10a)$$

$$\int_f (u \cdot n_j^f) p \, ds, \quad j = 1, \dots, n - \ell, p \in \mathbb{P}_{r-(\ell+1)}(f). \quad (10b)$$

Proof. Since $\{t_1^f, \dots, t_\ell^f, n_1^f, \dots, n_{n-\ell}^f\}$ forms a basis of \mathbb{R}^n , DoFs (10a)–(10b) are equivalent to

$$\int_f u \cdot p \, ds, \quad p \in \mathbb{P}_{r-(\ell+1)}(f; \mathbb{R}^n).$$

Then the unisolvence follows from Theorem 2.5 for the Lagrange element. \square

Next we use $\mathbb{B}_r \mathcal{T}^f$ or $\mathbb{B}_r \mathcal{N}^f$ to characterize the kernel or range of the trace operator, respectively. Define the polynomial bubble space

$$\mathbb{B}_r(\text{div}, T) := \ker(\text{tr}^{\text{div}}) \cap \mathbb{P}_r(T; \mathbb{R}^n).$$

It is obvious that $\mathbb{B}_0(\text{div}, T) = \{\mathbf{0}\}$ and $\mathbb{B}_1(\text{div}, T) = \{\mathbf{0}\}$.

Lemma 3.5. *For $r \geq 2$, it holds that*

$$\mathbb{B}_r(\text{div}, T) = \bigoplus_{\ell=1}^n \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_r \mathcal{T}^f, \quad (11)$$

and

$$\text{tr}^{\text{div}} : \bigoplus_{\ell=0}^{n-1} \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_r \mathcal{N}^f \rightarrow \text{tr}^{\text{div}} \mathbb{P}_r(T; \mathbb{R}^n) \quad (12)$$

is a bijection. Consequently

$$\dim \mathbb{B}_r(\text{div}, T) = \sum_{\ell=1}^n \binom{n+1}{\ell+1} \binom{r-1}{\ell} \binom{\ell}{1}.$$

Proof. Verification of

$$\bigoplus_{\ell=1}^n \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_r \mathcal{T}^f \subseteq \mathbb{B}_r(\text{div}, T)$$

is straightforward. For face F not containing f , $b_f|_F = 0$. For face F containing f , $\text{tr}^{\text{div}} \mathbf{u}|_F = \mathbf{u} \cdot \mathbf{n}_F = 0$ as $\mathbf{t}_i^f \cdot \mathbf{n}_F = 0$ for $\mathbf{u} \in \mathbb{B}_r \mathcal{T}^f$. Then apply the trace operator to the decomposition (9) and use $\text{tr}^{\text{div}}(\mathbb{B}_r \mathcal{T}^f) = 0$ to obtain

$$\text{tr}^{\text{div}} \left(\bigoplus_{\ell=0}^{n-1} \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_r \mathcal{N}^f \right) = \text{tr}^{\text{div}} \mathbb{P}_r(T; \mathbb{R}^n).$$

So the map tr^{div} in (12) is onto.

We then prove it is also injective. For an $f \in \Delta_\ell(T)$, recall that $\{\mathbf{n}_{F_i}, i \in f^*\}$ is the face normal basis of \mathcal{N}^f and $\{\hat{\mathbf{n}}_{F_i}, i \in f^*\}$ is its dual basis of \mathcal{N}^f . We expand $\mathbf{u} \in \bigoplus_{\ell=0}^{n-1} \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_r \mathcal{N}^f$ in the dual basis, i.e., $\mathbf{u} = \sum_{\ell=0}^{n-1} \sum_{f \in \Delta_\ell(T)} \sum_{i \in f^*} b_f p_f^i \hat{\mathbf{n}}_{F_i}$ with $p_f^i \in \mathbb{P}_{r-(\ell+1)}(f)$. We will prove if $\text{tr}^{\text{div}} \mathbf{u} = 0$, then $\mathbf{u} = 0$.

To do so, we consider the operator $N_i^f(\mathbf{u}) = (\mathbf{u} \cdot \mathbf{n}_{F_i})|_f$. Condition $\text{tr}^{\text{div}} \mathbf{u} = 0$ implies $N_i^f(\mathbf{u}) = 0$ for all $f \in \Delta(T)$ and $i \in f^*$. By the choice of the basis of \mathcal{N}^f and the property of bubble polynomial b_f , the corresponding N - ϕ matrix is block lower triangular with diagonal matrices in the diagonal block; see (5) for an illustration. Therefore $N_i^f(\mathbf{u}) = 0$ implies $\mathbf{u} = 0$. More specifically, we have $N_i^v(\mathbf{u}) = p_v^i = 0$ for $v \in \Delta_0(T)$ and $i \in v^*$. After that, we apply $N_i^f(\mathbf{u})$ to conclude $p_f^i = 0$ for $f \in \Delta_1(T)$. Using this forward substitution argument for the lower triangular matrix, we conclude all coefficient polynomials $p_f^i = 0$.

Once we have proved the map tr in (12) is bijective, we conclude (11) from the decomposition (9). \square

With this characterization, tangential DoFs in (10a) can be merged as one $\mathbb{B}_r(\text{div}, T)$.

3.2.2. Brezzi–Douglas–Marini element

Given an $f \in \Delta_\ell(T)$, we choose $\{\mathbf{n}_F, f \subseteq F \in \Delta_{n-1}(T)\}$ as the basis for its normal plane \mathcal{N}^f .

Lemma 3.6 (Local BDM Element). *The shape function space $\mathbb{P}_r(T; \mathbb{R}^n)$ is uniquely determined by the DoFs*

$$\int_F \mathbf{v} \cdot \mathbf{n}_F p \, ds, \quad p \in \mathbb{P}_r(F), F \in \Delta_{n-1}(T), \quad (13a)$$

$$\int_T \mathbf{v} \cdot \mathbf{p} \, dx, \quad \mathbf{p} \in \mathbb{B}_r(\text{div}, T). \quad (13b)$$

Proof. By the geometric decomposition of $\mathbb{P}_r(F)$ element, (13a) can be decomposed into

$$\int_f (\mathbf{v} \cdot \mathbf{n}_F)|_F p \, ds, \quad F \in \Delta_{n-1}(T), f \in \Delta_\ell(F), p \in \mathbb{P}_{r-(\ell+1)}(f), \ell = 0, 1, \dots, n-1. \quad (14)$$

We switch the ordering of f and F to

$$\int_f (\mathbf{v} \cdot \mathbf{n}_F)|_F p \, ds, \quad f \in \Delta_\ell(T), F \in \Delta_{n-1}(T), f \subseteq F, p \in \mathbb{P}_{r-(\ell+1)}(f), \ell = 0, 1, \dots, n-1,$$

which is exactly (10b) according to our choice of normal basis.

Therefore given a $\mathbf{v} \in \mathbb{P}_r(T; \mathbb{R}^n)$, if (13a) vanishes, then $\text{tr} \mathbf{v} = 0$ and consequently $\mathbf{v} \in \mathbb{B}_r(\text{div}, T)$. Finally the vanishing DoF (13b) implies $\mathbf{v} = 0$. \square

One benefit of using the decomposed version (14) instead of the merged one (13a) is that the well documented Lagrange basis functions can be used in the implementation of the face element; see [30].

We call the change from (14) to (13a) the facewise redistribution of normal DoFs. Namely by using the face normal basis, we can redistribute the DoFs on f to each $(n-1)$ -dimensional face F containing f .

To glue local finite elements to form an $H(\text{div})$ -conforming finite element, we need to enforce continuity of $\mathbf{v} \cdot \mathbf{n}_F$ by choosing \mathbf{n}_F depending only on F not element T .

Lemma 3.7 (BDM Space). *For each $F \in \Delta_{n-1}(\mathcal{T}_h)$, choose a normal vector \mathbf{n}_F . For the shape function space $\mathbb{P}_r(T; \mathbb{R}^n)$, the following DoFs*

$$\int_F \mathbf{v} \cdot \mathbf{n}_F p \, ds, \quad p \in \mathbb{P}_r(F), F \in \Delta_{n-1}(\mathcal{T}_h), \quad (15a)$$

$$\int_T \mathbf{v} \cdot \mathbf{p} \, dx, \quad \mathbf{p} \in \mathbb{B}_r(\text{div}, T), T \in \mathcal{T}_h, \quad (15b)$$

define an $H(\text{div})$ -conforming space

$$V_{\text{BDM}} = \{\mathbf{v}_h \in H(\text{div}, \Omega) : \mathbf{v}_h|_T \in \mathbb{P}_r(T; \mathbb{R}^n), \forall T \in \mathcal{T}_h\}.$$

Proof. On each element T , DoFs (15) will determine a function in $\mathbb{P}_r(T; \mathbb{R}^n)$ by Lemma 3.6. DoF (15a) will determine the trace $\mathbf{v} \cdot \mathbf{n}_F$ on F independent of the element containing F , and thus the function is $H(\text{div}, \Omega)$ -conforming. \square

We have the geometric decomposition of the global BDM element space

$$V_{\text{BDM}} = \bigoplus_{F \in \Delta_{n-1}(\mathcal{T}_h)} \bigoplus_{\ell=0}^{n-1} \bigoplus_{f \in \Delta_\ell(F)} \mathbb{B}_{r, \mathcal{N}^f}(F, \Omega) \oplus \bigoplus_{T \in \mathcal{T}_h} \mathbb{B}_r(\text{div}, T),$$

and

$$\dim V_{\text{BDM}} = |\Delta_{n-1}(\mathcal{T}_h)| \binom{n+r-1}{r} + |\Delta_n(\mathcal{T}_h)| \sum_{\ell=1}^n \binom{n+1}{\ell+1} \binom{r-1}{\ell}.$$

Here

$$\mathbb{B}_{r, \mathcal{N}^f}(F, \Omega) := \{v_h \in H(\text{div}, \Omega) : v_h|_T \in \mathbb{B}_{r, \mathcal{N}^f} \text{ for } T \in \mathcal{T}_h, F \subseteq T, v_h|_{T'} = 0 \text{ for } T' \in \mathcal{T}_h, F \not\subseteq T'\}.$$

3.2.3. Stenberg's type element

We can construct an $H(\text{div})$ -conforming element with more continuity on the normal plane of lower dimensional sub-simplices by choosing a global basis of \mathcal{N}^f .

In the t - n decomposition, if a basis vector t_i^f or n_i^f depends only on f not on element T containing f , we call it *global* and otherwise $t_i^f(T)$ or $n_i^f(T)$ is local and the corresponding DoFs are different for different T containing f . For a global basis vector, the corresponding DoF (10a) or (10b) only depends on f and thus imposes continuity in that direction. In the extreme case, if all t - n basis vectors are global, we obtain the Lagrange element.

The following is a generalization of Stenberg element by imposing more continuity on the normal plane for sub-simplices with dimension from 0 to m for some $-1 \leq m \leq n-2$. When $m = 0$, it is the original Stenberg's element [35], i.e., only continuous at vertices. When $m = n-2$, it is Christiansen–Hu–Hu element constructed in [22]. When $m = -1$, no DoF (16a) exists and thus reduces to the BDM element.

Lemma 3.8 (Stenberg Type Element). *Let $-1 \leq m \leq n-2$. For each $f \in \Delta_\ell(\mathcal{T}_h)$ with $\ell \leq m$, we choose $n-\ell$ normal vectors $\{n_1^f, \dots, n_{n-\ell}^f\}$. Then the DoFs*

$$\int_f v \cdot n_i^f p \, ds, \quad p \in \mathbb{P}_{r-(\ell+1)}(f), \quad f \in \Delta_\ell(\mathcal{T}_h), i = 1, \dots, n-\ell, \ell = 0, \dots, m, \quad (16a)$$

$$\int_f (v \cdot n_F)|_F p \, ds, \quad p \in \mathbb{P}_{r-(\ell+1)}(f), F \in \Delta_{n-1}(\mathcal{T}_h), f \in \Delta_\ell(F), \ell = m+1, \dots, n-1, \quad (16b)$$

$$\int_T v \cdot p \, dx, \quad p \in \mathbb{B}_r(\text{div}, T), \quad T \in \mathcal{T}_h, \quad (16c)$$

will determine a space $V_{\text{Stenberg}}^r \subset H(\text{div}, \Omega)$.

Proof. For $T \in \mathcal{T}_h$ and $f \in \Delta_\ell(T)$, both $\{n_1^f, \dots, n_{n-\ell}^f\}$ and $\{n_F, F \in \Delta_{n-1}(\mathcal{T}), f \subset F\}$ are bases of the normal plane \mathcal{N}^f . Then the number of DoFs (16) restricted to T equals to the number of DoFs (13). DoF (16a) determines DoF (15a) for $\ell = 0, \dots, m$. Thus we conclude the result from Lemma 3.7. \square

We have the geometric decomposition of the global Stenberg element space

$$V_{\text{Stenberg}}^r = \bigoplus_{\ell=0}^m \bigoplus_{f \in \Delta_\ell(\mathcal{T}_h)} \mathbb{B}_{r, \mathcal{N}^f}(\Omega) \oplus \bigoplus_{F \in \Delta_{n-1}(\mathcal{T}_h)} \bigoplus_{\ell=m+1}^{n-1} \bigoplus_{f \in \Delta_\ell(F)} \mathbb{B}_{r, \mathcal{N}^f}(F, \Omega) \oplus \bigoplus_{T \in \mathcal{T}_h} \mathbb{B}_r(\text{div}, T),$$

and

$$\dim V_{\text{Stenberg}}^r = \sum_{\ell=0}^m |\Delta_\ell(\mathcal{T}_h)| (n-\ell) \binom{r-1}{\ell} + |\Delta_{n-1}(\mathcal{T}_h)| \sum_{\ell=m+1}^{n-1} \binom{n}{\ell+1} \binom{r-1}{\ell} + |\Delta_n(\mathcal{T}_h)| \sum_{\ell=1}^n \binom{n+1}{\ell+1} \binom{r-1}{\ell} \binom{\ell}{1}.$$

Here

$$\mathbb{B}_{r, \mathcal{N}^f}(\Omega) := \{v_h \in H(\text{div}, \Omega) : v_h|_T \in \mathbb{B}_{r, \mathcal{N}^f} \text{ for } T \in \mathcal{T}_h, f \subseteq T, v_h|_{T'} = 0 \text{ for } T' \in \mathcal{T}_h, f \not\subseteq T'\}.$$

Clearly $V_{\text{Stenberg}}^r \subseteq V_{\text{BDM}}$, and $\dim V_{\text{Stenberg}}^r < \dim V_{\text{BDM}}$ for $0 \leq m \leq n-2$.

We introduce an n -dimensional smoothness vector $\mathbf{r} = (r_0, r_1, \dots, r_{n-1})^\top \in \mathbb{R}^n$ to characterize the smoothness of the finite element space at sub-simplices of dimension $\ell = 0, 1, \dots, n-1$. For the space V_{Stenberg}^r defined by DoFs (16), the smoothness vector is given by

$$\mathbf{r} = (0, -\frac{1}{n-1}, \dots, -\frac{m}{n-1}, -1, \dots, -1)^\top.$$

For an ℓ -dimensional face $f \in \Delta_\ell(\mathcal{T}_h)$, a smoothness parameter of $-\frac{\ell}{n-1}$ means the vector is discontinuous only on the ℓ -dimensional tangential space and continuous on the normal plane \mathcal{N}^f . The value -1 indicates that the DoF is redistributed to faces, and the vector is discontinuous on the tangential space of $(n-1)$ -dimensional faces. The same smoothness vector will determine the same global finite element space, although DoFs may be slightly different; see the modification of DoFs in Corollary 3.11.

Here are the smoothness vectors for the examples in Section 3.1: Nédélec element/BDM element: $\mathbf{r} = -1$ with $m = -1$, Stenberg element: $\mathbf{r} = (0, -1, -1)^\top$ with $m = 0$, and CHH element: $\mathbf{r} = (0, -1/2, -1)^\top$ with $m = 1$. These vectors give the smoothness properties of the respective finite element spaces at vertices, edges, and faces.

3.3. Discrete inf-sup condition

In the unsolvence of vector Lagrange elements, cf. the proof of [Lemma 3.4](#), any basis of \mathbb{R}^n at f is allowed. The t - n basis is used for two purposes: 1. the $H(\text{div})$ -conformity; 2. the discrete div stability.

If the $H(\text{div})$ -conformity is the only concern, we can simply choose the Lagrange element. Another and more important consideration is from the div stability. At the continuous level, we have the div stability, i.e., $\text{div} : H(\text{div}, \Omega) \rightarrow L^2(\Omega)$ is surjective, which has a continuous right inverse. A regular right inverse in $H^1(\Omega)$ also exists [\[36\]](#).

By the Euler's formula for homogeneous degree polynomials $\mathbb{H}_{r-1}(T)$, i.e. $\text{div}(xq) = (r-1+n)q$ for any $q \in \mathbb{H}_{r-1}(T)$, clearly we have $\text{div } \mathbb{P}_r(T; \mathbb{R}^n) = \mathbb{P}_{r-1}(T)$. Hence the discrete div stability in one element always holds. We discuss the global version. Let

$$\begin{aligned} V_h &:= \{v_h \in H(\text{div}, \Omega) : v_h|_T \in \mathbb{P}_r(T; \mathbb{R}^n) \text{ for each } T \in \mathcal{T}_h\}, \\ Q_h &:= \{q_h \in L^2(\Omega) : q_h|_T \in \mathbb{P}_{r-1}(T) \text{ for each } T \in \mathcal{T}_h\}. \end{aligned} \quad (17)$$

The discrete div stability refers to $\text{div } V_h = Q_h$ and the operator div has a continuous right inverse.

We will use the L^2 -inner product $(\cdot, \cdot)_T$ and define the orthogonal complement of a subspace $V \subset L^2(T)$ as V^\perp , i.e.,

$$V^\perp = \{u \in L^2(T) : (u, v)_T = 0 \quad \forall v \in V\}.$$

We first give the following characterization of the range of the div operator on the bubble polynomial space.

Lemma 3.9 (Lemma 3.6 in [\[28\]](#)). *It holds*

$$\text{div } \mathbb{B}_r(\text{div}, T) = \mathbb{P}_{r-1}(T) \cap \mathbb{R}^\perp, \quad r \geq 0.$$

Proof. When $r = 0, 1$, both sides are zero. Therefore we focus on $r \geq 2$.

The inclusion $\text{div}(\mathbb{B}_r(\text{div}, T)) \subseteq (\mathbb{P}_{r-1}(T) \cap \mathbb{R}^\perp)$ is proved through integration by parts

$$(\text{div } v, p)_T = -(v, \text{grad } p)_T = 0, \quad p \in \ker(\text{grad}) = \mathbb{R}.$$

If $\text{div}(\mathbb{B}_r(\text{div}, T)) \neq \mathbb{P}_{r-1}(T) \cap \mathbb{R}^\perp$, there exists a $p \in \mathbb{P}_{r-1}(T) \cap \mathbb{R}^\perp$ and $p \perp \text{div}(\mathbb{B}_r(\text{div}, T))$, which is equivalent to $\text{grad } p \perp \mathbb{B}_r(\text{div}, T)$. Expand the vector $\text{grad } p$ in the basis $\{n_i, i = 1, \dots, n\}$ as $\text{grad } p = \sum_{i=1}^n q_i n_i$ with $q_i \in \mathbb{P}_{r-2}(T)$. Then set $v_p = \sum_{i=1}^n |\nabla \lambda_i| q_i \lambda_0 \lambda_i t_{i,0} \in \mathbb{B}_r(\text{div}, T)$, where $t_{i,0} := v_0 - v_i$. We have

$$(\text{grad } p, v_p)_T = \sum_{i=1}^n \int_T q_i^2 \lambda_0 \lambda_i dx = 0,$$

which implies $q_i = 0$ for $i = 1, \dots, n$, i.e., $\text{grad } p = 0$ and $p = 0$ as $p \in \mathbb{P}_{r-1}(T) \cap \mathbb{R}^\perp$. \square

Next we verify the discrete div stability.

Proposition 3.10. *Let $r \geq 1$ and V_{BDM} be the BDM space defined in [Lemma 3.7](#) and Q_h defined by (17). It holds the discrete inf-sup condition*

$$\|q_h\|_0 \lesssim \sup_{v_h \in V_{\text{BDM}}} \frac{(\text{div } v_h, q_h)}{\|v_h\|_0 + \|\text{div } v_h\|_0} \quad \forall q_h \in Q_h. \quad (18)$$

Proof. By Theorem 1.1 in [\[36\]](#), there exists $v \in H^1(\Omega; \mathbb{R}^n)$ such that

$$\text{div } v = q_h, \quad \|v\|_1 \lesssim \|q_h\|_0. \quad (19)$$

Let $\tilde{v}_h \in V_{\text{BDM}}$ satisfy

$$\int_F \tilde{v}_h \cdot n_F ds = \int_F v \cdot n_F ds, \quad F \in \Delta_{n-1}(\mathcal{T}_h),$$

and the rest DoFs vanish. By the scaling argument,

$$\|\tilde{v}_h\|_0 + \|\text{div } \tilde{v}_h\|_0 \lesssim \|v\|_1 \lesssim \|q_h\|_0. \quad (20)$$

Clearly we have $\text{div}(\tilde{v}_h - v)|_T \in \mathbb{P}_{r-1}(T) \cap \mathbb{R}^\perp$ for each $T \in \mathcal{T}_h$. By [Lemma 3.9](#), there exists $b_h \in L^2(\Omega; \mathbb{R}^n)$ such that $b_h|_T \in \mathbb{B}_r(\text{div}, T)$ for each $T \in \mathcal{T}_h$, and

$$\text{div } b_h = \text{div}(v - \tilde{v}_h), \quad \|b_h\|_{0,T} \lesssim h_T \|\text{div}(\tilde{v}_h - v)\|_{0,T}. \quad (21)$$

Take $v_h = \tilde{v}_h + b_h \in V_{\text{BDM}}$. By (19) and (21), it holds

$$\text{div } v_h = \text{div } \tilde{v}_h + \text{div } b_h = \text{div } v = q_h. \quad (22)$$

It follows from (20) and (21) that

$$\|v_h\|_0 + \|\text{div } v_h\|_0 = \|v_h\|_0 + \|q_h\|_0 \leq \|\tilde{v}_h\|_0 + \|b_h\|_0 + \|q_h\|_0 \lesssim \|\tilde{v}_h\|_0 + h \|\text{div } \tilde{v}_h\|_0 + \|q_h\|_0 \lesssim \|q_h\|_0. \quad (23)$$

Combining (22)–(23) yields (18). \square

For the Stenberg-type element, the continuity at normal planes introduces some constraint and makes the discrete inf-sup condition harder. As all bubble functions are treated locally, $\operatorname{div} \mathbb{B}_r(\operatorname{div}, T) = \mathbb{P}_{r-1}(T) \cap \mathbb{R}^\perp$ still holds. We only need to show $\cup_{T \in \mathcal{T}_h} \mathbb{P}_0(T)$ is in the range of $\operatorname{div} V_h$, which requires the face average $\int_F \mathbf{v} \cdot \mathbf{n}_F \, ds$ in DoF.

We propose the following modification to have a better discrete inf-sup condition. Denote by

$$\mathbb{B}_{r,m+1}(F) := \left[\bigoplus_{\ell=m+1}^{n-1} \bigoplus_{f \in \Delta_\ell(F)} b_f \mathbb{P}_{r-(\ell+1)}(f) \right], \quad -1 \leq m \leq n-2.$$

Corollary 3.11. *With the same setting in Lemma 3.8 and further assume $r \geq m+2$ for $-1 \leq m \leq n-2$. Replacing DoF (16b) by*

$$\int_F \mathbf{v} \cdot \mathbf{n}_F \, q \, ds, \quad F \in \Delta_{n-1}(\mathcal{T}_h), \, q \in (\mathbb{B}_{r,m+1}(F) \cap \mathbb{P}_0^\perp(F)) \oplus \mathbb{P}_0(F), \quad (24)$$

will define the same finite element space.

Proof. When $-1 \leq m \leq n-2$ and $r \geq m+2$, $\dim \mathbb{B}_{r,m+1}(F) \geq |\Delta_{m+1}(F)| \geq \dim \mathbb{P}_0(F)$. Therefore the number of DoFs remains the same as DoF (16b).

Vanishing DoFs (16a) will imply $\mathbf{v} \cdot \mathbf{n}_F|_F \in \mathbb{B}_{r,m+1}(F)$, which can be determined by (24). So the unsolvence follows.

As the change will not affect the continuity, it will define the same finite element space. \square

Now $\int_F \mathbf{v} \cdot \mathbf{n}_F \, ds$ is in DoF (24). The proof of the following result is identical to that of Proposition 3.10.

Proposition 3.12. *Let $-1 \leq m \leq n-2$ and $r \geq m+2$. Let V_m^r be the $H(\operatorname{div})$ -conforming finite element defined in Lemma 3.8. The following discrete inf-sup condition holds with a constant independent of h*

$$\|q_h\|_0 \lesssim \sup_{\mathbf{v}_h \in V_m^r} \frac{(\operatorname{div} \mathbf{v}_h, q_h)}{\|\mathbf{v}_h\|_0 + \|\operatorname{div} \mathbf{v}_h\|_0} \quad \forall q_h \in Q_h. \quad (25)$$

4. Geometric decompositions of matrix face elements

In this section, we generalize the geometric decomposition of $H(\operatorname{div})$ -conforming vector finite elements to two $H(\operatorname{div})$ -conforming matrix finite elements: the traceless matrix \mathbb{T} and the symmetric matrix \mathbb{S} .

4.1. Traceless matrix elements

We consider the $H(\operatorname{div}, \Omega; \mathbb{T})$ -conforming finite element spaces, where $\mathbb{T} \in \mathbb{R}^{n \times n}$ is the set of square matrices with vanishing trace, i.e., the sum of the diagonal is zero.

We start from the tensor product of the Lagrange element and \mathbb{T} :

$$\mathbb{P}_r(T; \mathbb{T}) = \bigoplus_{\ell=0}^n \bigoplus_{f \in \Delta_\ell(T)} [b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathbb{T}].$$

That is each component of the matrix function is a Lagrange element of degree r and thus is continuous. To be $H(\operatorname{div})$ -conforming, however, normal continuity is sufficient.

To impose the normal continuity of a traceless matrix function, the key is a t - n decomposition at each sub-simplex. Here the t - n decomposition is with respect to the second component in the tensor product form $\mathbf{u} \otimes \mathbf{v}$ of representing a matrix. Given a sub-simplex $f \in \Delta_\ell(T)$, choose a t - n basis $\{\mathbf{t}_i^f, \mathbf{n}_j^f\}_{i=1, \dots, \ell}^{j=1, \dots, n-\ell}$ and decompose $\mathbb{R}^n = \mathcal{T}^f \oplus^\perp \mathcal{N}^f$. All basis vectors are normalized but may not be mutually orthogonal. By the tensor product the $n \times n$ matrix space \mathbb{M} has the following decomposition

$$\mathbb{M} = (\mathbb{R}^n \otimes \mathcal{T}^f) \oplus (\mathbb{R}^n \otimes \mathcal{N}^f). \quad (26)$$

For a matrix $\mathbf{A} \in \mathbb{M}$, $\operatorname{tr}_F^{\operatorname{div}}(\mathbf{A}) = \mathbf{A} \mathbf{n}_F$ and thus $\operatorname{tr}_F^{\operatorname{div}}(\mathbb{R}^n \otimes \mathcal{T}^f) = 0$. The normal component $\mathbb{R}^n \otimes \mathcal{N}^f$ will contribute to the normal trace.

We then modify (26) for \mathbb{M} to impose the traceless constraint while not changing $\operatorname{tr}^{\operatorname{div}}$. When computing the trace of a matrix, we use $\operatorname{trace}(\mathbf{u} \otimes \mathbf{v}) = \mathbf{v} \cdot \mathbf{u}$. We pick up the element $\mathbf{t}_1^f \otimes \mathbf{t}_1^f \in \mathbb{R}^n \otimes \mathcal{T}^f$ and use it to modify the basis in $(\mathbb{R}^n \otimes \mathcal{T}^f) \oplus (\mathbb{R}^n \otimes \mathcal{N}^f)$ to get the following t - n decomposition on $f \in \Delta_\ell(T)$ for $\ell \geq 1$:

$$\begin{aligned} \mathcal{T}^f(\mathbb{T}) &:= \operatorname{span}\{\mathbf{n}_j^f \otimes \mathbf{t}_i^f, 1 \leq i \leq \ell, 1 \leq j \leq n-\ell\} \oplus \operatorname{span}\{\mathbf{t}_i^f \otimes \mathbf{t}_j^f - (\mathbf{t}_i^f \cdot \mathbf{t}_j^f) \mathbf{t}_1^f \otimes \mathbf{t}_1^f, 1 \leq i, j \leq \ell\}, \\ \mathcal{N}^f(\mathbb{T}) &:= \operatorname{span}\{\mathbf{t}_i^f \otimes \mathbf{n}_j^f, 1 \leq i \leq \ell, 1 \leq j \leq n-\ell\} \oplus \operatorname{span}\{\mathbf{n}_i^f \otimes \mathbf{n}_j^f - (\mathbf{n}_i^f \cdot \mathbf{n}_j^f) \mathbf{t}_1^f \otimes \mathbf{t}_1^f, 1 \leq i, j \leq n-\ell\}. \end{aligned}$$

By counting the dimensions, it is easy to show the direct decomposition

$$\mathbb{T} = \mathcal{T}^f(\mathbb{T}) \oplus \mathcal{N}^f(\mathbb{T}), \quad f \in \Delta_\ell(T), \ell = 1, \dots, n.$$

For $\ell = 0$, i.e., at vertex $v \in \Delta_0(T)$, we understand $\mathcal{T}^v(\mathbb{T}) = \{0\}$ and $\mathcal{N}^v(\mathbb{T}) = \mathbb{T}$. For $\ell = n$, $\mathcal{T}^T(\mathbb{T}) = \mathbb{T}$ and $\mathcal{N}^T(\mathbb{T}) = \{0\}$. Coupled with the bubble polynomials, we define

$$\mathbb{B}_r \mathcal{T}^f(\mathbb{T}) := b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{T}^f(\mathbb{T}), \quad \mathbb{B}_r \mathcal{N}^f(\mathbb{T}) := b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{N}^f(\mathbb{T}).$$

Lemma 4.1. *The shape function space $\mathbb{P}_r(T; \mathbb{T})$ has a geometric decomposition*

$$\mathbb{P}_r(T; \mathbb{T}) = \bigoplus_{\ell=0}^n \bigoplus_{f \in \Delta_\ell(T)} [\mathbb{B}_r \mathcal{T}^f(\mathbb{T}) \oplus \mathbb{B}_r \mathcal{N}^f(\mathbb{T})].$$

A function $\mathbf{A} \in \mathbb{P}_r(T; \mathbb{T})$ is uniquely determined by the DoFs

$$\int_f \mathbf{A} : \mathbf{q} \, ds, \quad \mathbf{q} \in \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{T}^f(\mathbb{T}), f \in \Delta_\ell(T), \ell = 1, \dots, n, \quad (27a)$$

$$\int_f \mathbf{A} : \mathbf{q} \, ds, \quad \mathbf{q} \in \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{N}^f(\mathbb{T}), f \in \Delta_\ell(T), \ell = 0, \dots, n-1. \quad (27b)$$

Proof. Since $\mathbb{T} = \mathcal{T}^f(\mathbb{T}) \oplus \mathcal{N}^f(\mathbb{T})$, DoFs (27a)–(27b) are equivalent to

$$\int_f \mathbf{A} : \mathbf{q} \, ds, \quad \mathbf{q} \in \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathbb{T}, f \in \Delta_\ell(T), \ell = 0, \dots, n.$$

Then the unisolvence follows from the unisolvence of the Lagrange element. \square

Define the bubble polynomial space

$$\mathbb{B}_r(\text{div}, T; \mathbb{T}) := \mathbb{P}_r(T; \mathbb{T}) \cap \ker(\text{tr}^{\text{div}}).$$

Follow the same proof of Lemma 3.5, we have the characterization of the bubble space.

Lemma 4.2. *For $r \geq 2$, it holds that*

$$\mathbb{B}_r(\text{div}, T; \mathbb{T}) = \bigoplus_{\ell=1}^n \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_r \mathcal{T}^f(\mathbb{T}),$$

and

$$\text{tr}^{\text{div}} : \bigoplus_{\ell=0}^{n-1} \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_r \mathcal{N}^f(\mathbb{T}) \rightarrow \text{tr}^{\text{div}} \mathbb{P}_r(T; \mathbb{T})$$

is a bijection.

Similar to the generalized Stenberg element, we can redistribute some normal DoFs onto the $(n-1)$ -dimensional faces to obtain the following $H(\text{div}; \mathbb{T})$ element.

Theorem 4.3 ($H(\text{div}; \mathbb{T})$ -Conforming Finite Elements). *Let $0 \leq m \leq n-2$. For each $f \in \Delta_\ell(\mathcal{T}_h)$ with $\ell \leq m$, we choose $n-\ell$ normal vectors $\{\mathbf{n}_1^f, \dots, \mathbf{n}_{n-\ell}^f\}$. For each $F \in \Delta_{n-1}(\mathcal{T}_h)$, choose a normal vector \mathbf{n}_F . Then the DoFs*

$$\mathbf{A}(\mathbf{v}), \quad \mathbf{v} \in \Delta_0(\mathcal{T}_h), \mathbf{A} \in \mathbb{T}, \quad (28a)$$

$$\int_f (\mathbf{A} \mathbf{n}_i^f) \cdot \mathbf{q} \, ds, \quad f \in \Delta_\ell(\mathcal{T}_h), \mathbf{q} \in \mathbb{P}_{r-(\ell+1)}(f; \mathbb{R}^n), i = 1, \dots, n-\ell, \ell = 1, \dots, m, \quad (28b)$$

$$\int_f (\mathbf{A} \mathbf{n}_F)|_F \cdot \mathbf{q} \, ds, \quad F \in \Delta_{n-1}(\mathcal{T}_h), f \in \Delta_\ell(F), \mathbf{q} \in \mathbb{P}_{r-(\ell+1)}(f; \mathbb{R}^n), \ell = m+1, \dots, n-1, \quad (28c)$$

$$\int_T \mathbf{A} : \mathbf{q} \, dx, \quad T \in \mathcal{T}_h, \mathbf{q} \in \mathbb{B}_r(\text{div}, T; \mathbb{T}), \quad (28d)$$

will determine a space $V^r(\mathbb{T}) \subset H(\text{div}, \Omega; \mathbb{T})$, where

$$V^r(\mathbb{T}) := \{\mathbf{A} \in L^2(\Omega; \mathbb{T}) : \mathbf{A}|_T \in \mathbb{P}_r(T; \mathbb{T}) \quad \forall T \in \mathcal{T}_h, \text{DoFs (28a)–(28b) are single-valued across } f \in \Delta_\ell(\mathcal{T}_h) \text{ for } \ell = 0, \dots, m, \\ \text{DoF (28c) is single-valued across } F \in \Delta_{n-1}(\mathcal{T}_h)\}.$$

Proof. For $T \in \mathcal{T}_h$ and $f \in \Delta_\ell(T)$, both $\{\mathbf{n}_1^f, \dots, \mathbf{n}_{n-\ell}^f\}$ and $\{\mathbf{n}_F, F \in \Delta_{n-1}(\mathcal{T}), f \subset F\}$ are basis of the normal plane \mathcal{N}^f . Then DoFs (28) restricted to T are equivalent to DoFs (27), thus uniquely determine $\mathbb{P}_r(T; \mathbb{T})$.

For $F \in \Delta_{n-1}(\mathcal{T}_h)$, DoFs (28a)–(28b) restricted to F will determine

$$\int_f (\mathbf{A} \mathbf{n}_F)|_F \cdot \mathbf{q} \, ds, \quad f \in \Delta_\ell(F), \mathbf{q} \in \mathbb{P}_{r-(\ell+1)}(f; \mathbb{R}^n), \ell = 0, \dots, m.$$

Then by the unisolvence of Lagrange element in Theorem 2.5, $(\mathbf{A} \mathbf{n}_F)|_F$ is determined by DoFs (28a)–(28c) restricted to F . Therefore $V^r(\mathbb{T}) \subset H(\text{div}, \Omega; \mathbb{T})$. \square

To have a better discrete div stability, we modify the face DoFs.

Corollary 4.4. *With the same setting in Theorem 4.3 and further assume $r \geq m+2$ for $0 \leq m \leq n-3$, and $r \geq n+1$ for $m = n-2$. Replacing DoF (28c) by*

$$\int_F (\mathbf{A} \mathbf{n}_F) \cdot \mathbf{q} \, ds, \quad F \in \Delta_{n-1}(\mathcal{T}_h), \mathbf{q} \in [(\mathbb{B}_{r,m+1}(F) \cap \mathbb{P}_1^\perp(F)) \oplus \mathbb{P}_1(F)] \otimes \mathbb{R}^n, \quad (29)$$

will define the same finite element space.

Proof. When $m = n - 2$, we require $r \geq n + 1$ so that $r - (m + 1 + 1) \geq 1$ and $\dim \mathbb{B}_{r,m+1}(F) \geq \dim \mathbb{P}_1(F)$. When $0 \leq m \leq n - 3$ and $r \geq m + 2$, $\dim \mathbb{B}_{r,m+1}(F) \geq |\Delta_{m+1}(F)| \geq \dim \mathbb{P}_1(F)$. Therefore the number of DoFs remains the same as DoF (28c).

Vanishing DoFs (28a)–(28b) will imply $\mathbf{A} \mathbf{n}_F|_F \in \mathbb{B}_{r,m+1}(F) \otimes \mathbb{R}^n$, which can be determined by (29). So the unisolvence follows. As the change will not affect the continuity, it will define the same finite element space. \square

DoF (29) is more friendly for verifying the discrete inf-sup condition and (28c) is better for the uni-solvence and implementation.

When $m = 0$, it is the generalization of Stenberg element for vector functions to traceless tensor functions. Almost all DoFs are redistributed to face F except at the vertex, where the traceless constraint is imposed. The case $n = 3, r \geq 2$ is the $H(\text{div}; \mathbb{T})$ element constructed in [11].

Remark 4.5. Comparing with the vector face element, cf. Lemma 3.8, m starts from 0 not -1 . Namely the $H(\text{div}; \mathbb{T})$ -element should be continuous at vertices. We argue that the continuity at vertices is also necessary. Take a vertex in $\Delta_0(T)$, for example v_0 . Then $\mathbf{A} \mathbf{n}_{F_i}(v_0)$ is determined by the vector $\mathbf{A} \mathbf{n}_{F_i}|_{F_i} \in \mathbb{R}^n$ for $i = 1, \dots, n$. If it is continuous on each face but not on vertices, the number of elements in $\mathbf{A} \mathbf{n}_{F_i}(v_0)$ is n for each face. Running i from 1 to n , $\mathbf{A}(v_0)$ is determined by n^2 conditions, which is more than $\dim \mathbb{T} = n^2 - 1$.

4.2. Discrete div stability for traceless tensors

We first show the div stability of the bubble space $\mathbb{B}_r(\text{div}, T; \mathbb{T})$. Denote by $\mathbf{t}_{i,j}$ the edge vector from v_i to v_j . By computing the constant directional derivative $\mathbf{t}_{i,j} \cdot \nabla \lambda_\ell$ by values on the two vertices, we have

$$\mathbf{t}_{i,j} \cdot \nabla \lambda_\ell = \delta_{j\ell} - \delta_{i\ell} = \begin{cases} 1, & \text{if } \ell = j, \\ -1, & \text{if } \ell = i, \\ 0, & \text{if } \ell \neq i, j. \end{cases}$$

Lemma 4.6. The set of traceless tensors $\{\nabla \lambda_i \otimes \mathbf{t}_{i+1,j}\}_{j \in \{0, \dots, n\} \setminus \{i, i+1\}}^{i=0, \dots, n}$ is a basis of the traceless tensor space \mathbb{T} . Its dual basis is $\{\mathbf{t}_{j,i} \otimes \nabla \lambda_j + \frac{1}{n} I\}_{j \in \{0, \dots, n\} \setminus \{i, i+1\}}^{i=0, \dots, n}$. All indices are modulo n .

Proof. It suffices to prove

$$(\mathbf{t}_{\ell,k} \otimes \nabla \lambda_\ell) : (\nabla \lambda_i \otimes \mathbf{t}_{i+1,j}) = \delta_{ik} \delta_{j\ell}$$

for $0 \leq i, k \leq n, j \in \{0, \dots, n\} \setminus \{i, i+1\}$, and $\ell \in \{0, \dots, n\} \setminus \{k, k+1\}$.

When $i = k$, by $\ell \neq i, i+1$, it follows

$$(\mathbf{t}_{\ell,k} \otimes \nabla \lambda_\ell) : (\nabla \lambda_i \otimes \mathbf{t}_{i+1,j}) = \nabla \lambda_\ell \cdot \mathbf{t}_{i+1,j} = \delta_{j\ell}.$$

When $i = \ell$, by $i \neq j$, it follows

$$(\mathbf{t}_{\ell,k} \otimes \nabla \lambda_\ell) : (\nabla \lambda_i \otimes \mathbf{t}_{i+1,j}) = 0.$$

When $i \neq k, \ell$, clearly $(\mathbf{t}_{\ell,k} \otimes \nabla \lambda_\ell) : (\nabla \lambda_i \otimes \mathbf{t}_{i+1,j}) = 0$. \square

Let $\text{RT} = \{a\mathbf{x} + b : a \in \mathbb{R}, b \in \mathbb{R}^n\}$. For a matrix \mathbf{A} , define $\text{dev } \mathbf{A} = \mathbf{A} - \frac{1}{n} \text{trace}(\mathbf{A})I$ as the projection of \mathbf{A} to \mathbb{T} . It is straight forward to verify $\ker(\text{dev grad}) = \text{RT}$. Again let RT^\perp be the L^2 -orthogonal complement in $L^2(T; \mathbb{R}^n)$.

Lemma 4.7. For each $T \in \mathcal{T}_h$, it holds

$$\text{div } \mathbb{B}_r(\text{div}, T; \mathbb{T}) = \mathbb{P}_{r-1}(T; \mathbb{R}^n) \cap \text{RT}^\perp. \quad (30)$$

Proof. It follows from the integration by parts that

$$\text{div } \mathbb{B}_r(\text{div}, T; \mathbb{T}) \subseteq (\mathbb{P}_{r-1}(T; \mathbb{R}^n) \cap \text{RT}^\perp).$$

We claim the equality holds. If $\text{div } \mathbb{B}_r(\text{div}, T; \mathbb{T}) \subset (\mathbb{P}_{r-1}(T; \mathbb{R}^n) \cap \text{RT}^\perp)$, then there exists $\mathbf{u} \in \mathbb{P}_{r-1}(T; \mathbb{R}^n) \cap \text{RT}^\perp$ satisfying the orthogonality condition $(\mathbf{u}, \text{div } \mathbf{A})_T = 0$ for any $\mathbf{A} \in \mathbb{B}_r(\text{div}, T; \mathbb{T})$. Equivalently

$$(\text{dev grad } \mathbf{u}, \mathbf{A})_T = 0 \quad \forall \mathbf{A} \in \mathbb{B}_r(\text{div}, T; \mathbb{T}).$$


By expressing $\text{dev grad } \mathbf{u} = \sum_{i=0}^n \sum_{j \in \{0, \dots, n\} \setminus \{i, i+1\}} q_{ij} (\mathbf{t}_{j,i} \otimes \nabla \lambda_j + \frac{1}{n} I)$ with $q_{ij} \in \mathbb{P}_{r-2}(T)$, we choose

$$\mathbf{A} = \sum_{i=0}^n \sum_{j \in \{0, \dots, n\} \setminus \{i, i+1\}} \lambda_{i+1} \lambda_j q_{ij} \nabla \lambda_i \otimes \mathbf{t}_{i+1,j} \in \mathbb{B}_r(\text{div}, T; \mathbb{T}).$$

Then we have

$$\sum_{i=0}^n \sum_{j \in \{0, \dots, n\} \setminus \{i, i+1\}} (\lambda_{i+1} \lambda_j q_{ij}, q_{ij})_T = 0.$$

Therefore $q_{ij} = 0$ for all i and j . Thus $\mathbf{u} = 0$. \square



(a) t - n decomposition at a vertex. (b) t - n decomposition on an edge. (c) t - n decomposition on a face. (d) t - n decomposition of a tetrahedron. No normal component.

Fig. 4. Several t - n decompositions for \mathbb{S} in \mathbb{R}^3 . Blocks with the same symbol (circle, square, or diamond) are in the same constraint sequence and the white block is used as the pair index. Color of the block represents: Green: free rows and free blocks; Blue: all free indices not in free rows; Red: bubble functions. Blue or green blocks are free indices in $\mathcal{N}^f(\mathbb{S})$. All white blocks are pair indices and the corresponding coefficients are determined by the free variables through the constraints. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

We mention that characterization (30) in three dimensions is firstly proved in [12].

Proposition 4.8 (Discrete Inf-Sup Condition for $H(\text{div}; \mathbb{T})$ -Conforming Finite Elements). *Let $0 \leq m \leq n-2$. Let $r \geq m+2$ for $0 \leq m \leq n-3$, and $r \geq n+1$ for $m = n-2$. Let $V^r(\mathbb{T})$ be defined in Theorem 4.3. It holds the discrete inf-sup condition*

$$\|v_h\|_0 \lesssim \sup_{A_h \in V^r(\mathbb{T})} \frac{(\text{div } A_h, v_h)}{\|A_h\|_0 + \|\text{div } A_h\|_0} \quad \forall v_h \in Q_h, \quad (31)$$

where $Q_h := \{v_h \in L^2(\Omega; \mathbb{R}^n) : v_h|_T \in \mathbb{P}_{r-1}(T; \mathbb{R}^n) \text{ for each } T \in \mathcal{T}_h\}$.

Proof. First there exists $A \in H^1(\Omega; \mathbb{T})$ such that

$$\text{div } A = v_h, \quad \|A\|_1 \lesssim \|v_h\|_0. \quad (32)$$

Thanks to DoFs (28) and (29), we let $\tilde{A}_h \in V^r(\mathbb{T})$ satisfy

$$\int_F (\tilde{A}_h n_F) \cdot q \, ds = \int_F (A n_F) \cdot q \, ds, \quad q \in \mathbb{P}_1(F; \mathbb{R}^n), F \in \Delta_{n-1}(\mathcal{T}_h),$$

and other DoFs vanish. By the scaling argument,

$$\|\tilde{A}_h\|_0 + \|\text{div } \tilde{A}_h\|_0 \lesssim \|A\|_1 \lesssim \|v_h\|_0. \quad (33)$$

Then through integration by parts we have $\text{div}(\tilde{A}_h - A)|_T \in \mathbb{P}_{r-1}(T; \mathbb{R}^n) \cap \text{RT}^\perp$ for each $T \in \mathcal{T}_h$. By Lemma 4.7, there exists $b_h \in L^2(\Omega; \mathbb{T})$ such that $b_h|_T \in \mathbb{B}_r(\text{div}, T; \mathbb{T})$ for each $T \in \mathcal{T}_h$, and

$$\text{div } b_h = \text{div}(A - \tilde{A}_h), \quad \|b_h\|_{0,T} \lesssim h_T \|\text{div}(\tilde{A}_h - A)\|_{0,T}. \quad (34)$$

Take $A_h = \tilde{A}_h + b_h \in V^r(\mathbb{T})$. By (32) and (34), it holds

$$\text{div } A_h = \text{div } \tilde{A}_h + \text{div } b_h = \text{div } A = v_h. \quad (35)$$

It follows from (33) and (34) that

$$\|A_h\|_0 + \|\text{div } A_h\|_0 = \|A_h\|_0 + \|v_h\|_0 \leq \|\tilde{A}_h\|_0 + \|b_h\|_0 + \|v_h\|_0 \lesssim \|\tilde{A}_h\|_0 + h \|\text{div } \tilde{A}_h\|_0 + \|v_h\|_0 \lesssim \|v_h\|_0. \quad (36)$$

Combining (35)–(36) yields (31). \square

4.3. Symmetric matrix elements

We start from the tensor product of the Lagrange element and the symmetric matrix \mathbb{S} :

$$\mathbb{P}_r(T; \mathbb{S}) = \bigoplus_{\ell=0}^n \bigoplus_{f \in \Delta_\ell(T)} [b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathbb{S}]. \quad (37)$$

The construction process is similar to the traceless case in which t - n decompositions of \mathbb{S} at sub-simplices are the key. Additional complication arises as the $n(n-1)/2$ symmetry constraints are more complicated than only 1 traceless constraint.

Let $\{v_1, \dots, v_n\}$ be a basis of \mathbb{R}^n . Then $\mathbb{M} = \{\sum_{i,j=1}^n a_{ij} v_i \otimes v_j, a_{ij} \in \mathbb{R}\}$ is the space of $n \times n$ -matrices. An element $A \in \mathbb{M}$ can be identified with the coefficient matrix (a_{ij}) and will be still denoted by A . For a better explanation, we illustrate an $n \times n$ -matrix by $n \times n$ blocks; see Fig. 4 for the case $n = 3$. Block (i, j) corresponds to the basis function $v_i \otimes v_j$. We will identify $n(n+1)/2$ blocks and modify corresponding basis function to form a basis of \mathbb{S} with consideration of the normal continuity.

Given an $f \in \Delta_\ell(T)$, choose a t - n basis $\{t_i^f, n_j^f\}_{i=1, \dots, \ell}^{j=1, \dots, n-\ell}$ and decompose $\mathbb{R}^n = \mathcal{T}^f \oplus \mathcal{N}^f$. All basis vectors are normalized. We have the matrix decomposition

$$\mathbb{M} = (\mathbb{R}^n \otimes \mathcal{T}^f) \oplus (\mathbb{R}^n \otimes \mathcal{N}^f).$$

We modify each component to impose the symmetric constraint. For the tangential component, we simply take

$$\mathcal{T}^f(\mathbb{S}) := (\mathbb{R}^n \otimes \mathcal{T}^f) \cap \mathbb{S}.$$

Using the tensor product with the scalar bubble polynomial, we can construct

$$\mathbb{B}_r \mathcal{T}^f(\mathbb{S}) := b_r \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{T}^f(\mathbb{S}) \subset \ker(\text{tr}^{\text{div}}) \cap \mathbb{P}_r(T; \mathbb{S}).$$

We provide a specific example for $n = 3, \ell = 2$, i.e., a t - n decomposition on a face F of a tetrahedron. Choose a t - n basis $\{t_1^F, t_2^F, n_F\}$. An element in $\mathbb{R}^n \otimes \mathcal{T}^F$ has the matrix representation $\begin{pmatrix} * & * & 0 \\ * & * & 0 \\ * & * & 0 \end{pmatrix}$, where $*$ represents a generic value. The symmetric

constraint implies an element in $\mathcal{T}^F(\mathbb{S})$ has the form $\begin{pmatrix} * & * & 0 \\ * & * & 0 \\ 0 & 0 & 0 \end{pmatrix}$. The value a_{21} is left blank since due to the symmetric constraint it is equal to a_{12} . We call $(1, 2)$ a free index while $(2, 1)$ is the pair index of $(1, 2)$.

We then move to the normal component. A naive definition of $\mathcal{N}^f(\mathbb{S})$ would be $(\mathbb{R}^n \otimes \mathcal{N}^f) \cap \mathbb{S}$. Unfortunately in general for three subspaces of a vector space:

$$(A \cap C) \oplus (B \cap C) \subseteq (A \oplus B) \cap C.$$

And the equality may not hold. We continue the example with $n = 3, \ell = 2$. An element in $\mathbb{R}^n \otimes \mathcal{N}^F$ has the matrix form $\begin{pmatrix} 0 & 0 & * \\ 0 & 0 & * \\ 0 & 0 & * \end{pmatrix}$

w.r.t the basis $\{t_1^F, t_2^F, n_F\}$. An element in $(\mathbb{R}^n \otimes \mathcal{N}^F) \cap \mathbb{S}$ has the form $\begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & * \end{pmatrix}$. Then

$$\dim((\mathbb{R}^n \otimes \mathcal{T}^F) \cap \mathbb{S}) + \dim((\mathbb{R}^n \otimes \mathcal{N}^F) \cap \mathbb{S}) = 3 + 1 = 4 < 6 = \dim \mathbb{S}.$$

The discrepancy is due to the fact that the constraints $a_{13} = a_{31}$ and $a_{23} = a_{32}$ have been used duplicately in both $(\mathbb{R}^n \otimes \mathcal{T}^F) \cap \mathbb{S}$ and $(\mathbb{R}^n \otimes \mathcal{N}^F) \cap \mathbb{S}$. We need to make sure one constraint is used only once either in $\mathcal{T}^f(\mathbb{S})$ or $\mathcal{N}^f(\mathbb{S})$.

To this end, for each constraint, we set the index (j, i) with $i < j$ as the pair index and call (i, j) a free index. The pair index value a_{ji} is determined by the free variable a_{ij} through the constraint $a_{ji} = a_{ij}$.

We introduce the concept: *normal constraints*. We call the constraint $a_{ij} = a_{ji}, i \neq j$, a normal constraint if both (i, j) and (j, i) are in the normal component, i.e., $i, j \geq \ell + 1$ w.r.t the t - n basis $\{t_i^f, n_j^f\}_{i=1, \dots, \ell}^{j=1, \dots, n-\ell}$. For example, in Fig. 4(b), for an edge $e \in \Delta_1(T)$, there are $(n-1)(n-2)/2$ normal constraints. As the constraint involves two entries, no normal constraints for $F \in \Delta_{n-1}(T)$; see Fig. 4(c). The normal constraints will be imposed inside the normal components.

For non-normal constraint, in the pair index (j, i) , i is inside the tangential component. We can use it to change the basis without affecting the normal trace. As an example, consider the circle block in Fig. 4(b)–(c), the basis function is changing from $t \otimes n$ to $\text{sym}(t \otimes n) = (n \otimes t + t \otimes n)/2$ and $2 \text{sym}(t \otimes n)n_F = (t \otimes n)n_F$. By doing this way, we ensure each constraint is used only once and the normal trace remains unchanged.

In summary, for $f \in \Delta_\ell(T)$, $\ell = 0, \dots, n$, define

$$\begin{aligned} \mathcal{T}^f(\mathbb{S}) &:= \text{span}\{\text{sym}(t_i^f \otimes t_j^f), 1 \leq i \leq j \leq \ell\}, \\ \mathcal{N}^f(\mathbb{S}) &:= \text{span}\{\text{sym}(t_i^f \otimes n_j^f), 1 \leq i \leq \ell, 1 \leq j \leq n - \ell\} \oplus \text{span}\{\text{sym}(n_i^f \otimes n_j^f), 1 \leq i \leq j \leq n - \ell\}. \end{aligned}$$

By counting the dimensions, it is easy to show the direct decomposition

$$\mathbb{S} = \mathcal{T}^f(\mathbb{S}) \oplus \mathcal{N}^f(\mathbb{S}).$$

Then tensor product with the scalar bubble function to obtain

$$\mathbb{B}_r \mathcal{T}^f(\mathbb{S}) = b_r \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{T}^f(\mathbb{S}), \quad \mathbb{B}_r \mathcal{N}^f(\mathbb{S}) = b_r \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{N}^f(\mathbb{S}).$$

With the t - n decomposition, (37) can be rewritten as

$$\mathbb{P}_r(T; \mathbb{S}) = \bigoplus_{\ell=0}^n \bigoplus_{f \in \Delta_\ell(T)} [\mathbb{B}_r \mathcal{T}^f(\mathbb{S}) \oplus \mathbb{B}_r \mathcal{N}^f(\mathbb{S})].$$

Define the polynomial bubble space

$$\mathbb{B}_r(\text{div}, T; \mathbb{S}) := \ker(\text{tr}^{\text{div}}) \cap \mathbb{P}_r(T; \mathbb{S}).$$

Again the tangential component $\mathbb{B}_r \mathcal{T}^f(\mathbb{S})$ will contribute to the bubble space. Namely for $r \geq 2$, it holds that

$$\mathbb{B}_r(\text{div}, T; \mathbb{S}) = \bigoplus_{\ell=1}^n \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_r \mathcal{T}^f(\mathbb{S}). \quad (38)$$

Characterization of $\mathbb{B}_r(\text{div}, T; \mathbb{S})$ in (38) is new and different with the one given in [16,17].

The normal component $\mathbb{B}_r \mathcal{N}^f(\mathbb{S})$ will determine the trace of the div operator. Notice that due to the constraint, not all n^2 components of the matrix are included when defining $\mathbb{B}_r \mathcal{T}^f(\mathbb{S})$ and $\mathbb{B}_r \mathcal{N}^f(\mathbb{S})$. In Fig. 4, function values in all white blocks are determined by the corresponding free variables through the constraints.

Unlike the traceless case, not all normal DoFs can be redistributed to faces since the normal constraint should be imposed on $\mathcal{N}^f \otimes \mathcal{N}^f$ with a global normal basis $\{\mathbf{n}_i^f\}$, which can be thought of as the super-smoothness induced by the constraint. For example, for symmetric matrix \mathbf{A} , the restriction of \mathbf{A} to the normal plane of f , which is a symmetric matrix of smaller size $(n - \ell) \times (n - \ell)$, should be continuous due to DoF (39b). The tangential-normal component can be redistributed to face F . Therefore, in DoFs (39), (39c) is posed globally and (39d) is facewisely.

Theorem 4.9 (*H(div; \mathbb{S})-Conforming Finite Elements*). Let $0 \leq m \leq n - 2$. For each $f \in \Delta_\ell(\mathcal{T}_h)$, we choose a global t -n basis $\{t_1^f, \dots, t_\ell^f, \mathbf{n}_1^f, \dots, \mathbf{n}_{n-\ell}^f\}$. Then the DoFs

$$\mathbf{A}(\mathbf{v}), \quad \mathbf{v} \in \Delta_0(\mathcal{T}_h), \mathbf{A} \in \mathbb{S}, \quad (39a)$$

$$\int_f ((\mathbf{n}_i^f)^T \mathbf{A} \mathbf{n}_j^f) q \, ds, \quad f \in \Delta_\ell(\mathcal{T}_h), q \in \mathbb{P}_{r-(\ell+1)}(f), 1 \leq i \leq j \leq n - \ell, \ell = 1, \dots, n - 1, \quad (39b)$$

$$\int_f ((t_i^f)^T \mathbf{A} \mathbf{n}_j^f) q \, ds, \quad f \in \Delta_\ell(\mathcal{T}_h), q \in \mathbb{P}_{r-(\ell+1)}(f), i = 1, \dots, \ell, j = 1, \dots, n - \ell, \ell = 1, \dots, m, \quad (39c)$$

$$\int_f ((t_i^f)^T \mathbf{A} \mathbf{n}_F) |_F q \, ds, \quad F \in \Delta_{n-1}(\mathcal{T}_h), f \in \Delta_\ell(F), q \in \mathbb{P}_{r-(\ell+1)}(f), i = 1, \dots, \ell, \ell = m + 1, \dots, n - 1, \quad (39d)$$

$$\int_T \mathbf{A} : \mathbf{q} \, dx, \quad T \in \mathcal{T}_h, \mathbf{q} \in \mathbb{B}_r(\text{div}, T; \mathbb{S}), \quad (39e)$$

will determine a space $V^r(\mathbb{S}) \subset H(\text{div}, \Omega; \mathbb{S})$, where

$$V^r(\mathbb{S}) := \{\mathbf{A} \in L^2(\Omega; \mathbb{S}) : \mathbf{A}|_T \in \mathbb{P}_r(T; \mathbb{S}) \quad \forall T \in \mathcal{T}_h,$$

DoFs (39a)–(39b) are single-valued across $f \in \Delta_\ell(\mathcal{T}_h)$ for $\ell = 0, \dots, n - 1$,

DoF (39c) is single-valued across $f \in \Delta_\ell(\mathcal{T}_h)$ for $\ell = 1, \dots, m$,

DoF (39d) is single-valued across $F \in \Delta_{n-1}(\mathcal{T}_h)\}$.

Proof. For $T \in \mathcal{T}_h$ and $f \in \Delta_\ell(T)$, both $\{\mathbf{n}_1^f, \dots, \mathbf{n}_{n-\ell}^f\}$ and $\{\mathbf{n}_F, F \in \Delta_{n-1}(\mathcal{T}_h), f \subset F\}$ are basis of the normal plane \mathcal{N}^f . DoFs (39a)–(39d) restricted to T will determine normal component $\mathbb{B}_r \mathcal{N}^f(\mathbb{S})$ and (39e) for the tangential component $\mathbb{B}_r \mathcal{T}^f(\mathbb{S})$, thus uniquely determine $\mathbb{P}_r(T; \mathbb{S})$.

For $F \in \Delta_{n-1}(\mathcal{T}_h)$, DoFs (39a)–(39d) restricted to F will determine

$$\int_f (\mathbf{A} \mathbf{n}_F) |_F \cdot \mathbf{q} \, ds, \quad f \in \Delta_\ell(F), \mathbf{q} \in \mathbb{P}_{r-(\ell+1)}(f; \mathbb{R}^n), \ell = 0, \dots, n - 1,$$

and thanks to the unisolvence of Lagrange element in Theorem 2.5 will determine $(\mathbf{A} \mathbf{n}_F) |_F$. Therefore $V^r(\mathbb{S}) \subset H(\text{div}, \Omega; \mathbb{S})$. \square

When $m = n - 2$, it is the Hu-Zhang element [16,17]. When $m = 0$, DoF (39d) can be further merged to one and lead to the modification in [28, Lemma 4.5]

$$\int_F (\Pi_F \mathbf{A} \mathbf{n}_F) \cdot \mathbf{q} \, ds, \quad F \in \Delta_{n-1}(\mathcal{T}_h), \mathbf{q} \in \text{ND}_{r-2}(F), \quad (40)$$

where $\text{ND}_{r-2}(F) := \{\mathbf{q} \in \mathbb{P}_{r-1}(F; \mathbb{R}^{n-1}) : \mathbf{q} \cdot \mathbf{x} \in \mathbb{P}_{r-1}(F)\}$ and Π_F is the projection of a vector to the plane \mathcal{T}^F .

4.4. Discrete div stability for symmetric tensors

For each $T \in \mathcal{T}_h$, the range of the div operator on the bubble space of symmetric tensors [16,17] is

$$\text{div } \mathbb{B}_r(\text{div}, T; \mathbb{S}) = \mathbb{P}_{r-1}(T; \mathbb{R}^n) \cap \text{RM}^\perp, \quad (41)$$

where $\text{RM} = \{\mathbf{N} \mathbf{x} + \mathbf{b} : \mathbf{N} \in \mathbb{K}, \mathbf{b} \in \mathbb{R}^n\}$ and RM^\perp is the L^2 -orthogonal complement in $L^2(T; \mathbb{R}^n)$. It can be proved similar to Lemma 4.7 and an abstract version will be proved in Lemma 6.9.

Applying the same argument as in Proposition 4.8, we derive the div stability of space $V^r(\mathbb{S})$ from (41).

Proposition 4.10. Let $r \geq n + 1$ and $V^r(\mathbb{S})$ be defined in Theorem 4.9. It holds the discrete inf-sup condition

$$\|\mathbf{v}_h\|_0 \lesssim \sup_{\mathbf{A}_h \in V^r(\mathbb{S})} \frac{(\text{div } \mathbf{A}_h, \mathbf{v}_h)}{\|\mathbf{A}_h\|_0 + \|\text{div } \mathbf{A}_h\|_0} \quad \forall \mathbf{v}_h \in Q_h.$$

Notice that due to the extra normal continuity, we cannot modify the face DoFs to relax the degree requirement $r \geq n + 1$ to $r \geq 2$ as we have done for the traceless element in Proposition 4.8. Thanks to DoF (40), the tangential-normal component contains $\mathbb{P}_1(F)$ for $r \geq 2$. Lower order $H(\text{div}; \mathbb{S})$ -conforming finite elements are designed by enriching the symmetric quadratic polynomial space with only $(n + 1)$ -order normal-normal face bubbles in [37], which have only $n(n + 1)^2$ DoFs for the reduced one.

5. Constraint tensor spaces

In this section we shall introduce the constraint tensor space \mathbb{X} as a kernel space, and discover bases of \mathbb{X} . We first recall some background on differential forms, then give concrete formulae on the algebraic operator $s^{k,n-1}$ and define the constraint tensor space \mathbb{X} as the kernel of $s^{k,n-1}$. Lastly we present two bases of \mathbb{X} .

5.1. Background on differential forms

5.1.1. Increasing sequence

We mainly follow the notation set in [20] but with some simplification. For non-negative integers a, b, l, m , with $0 \leq b - a \leq m - l$, define the set of increasing sequences as

$$\Sigma(a : b, l : m) := \{\sigma : \{a, \dots, b\} \rightarrow \{l, \dots, m\} \mid \sigma(a) < \sigma(a+1) < \dots < \sigma(b)\}.$$

We will overload the notation σ as its range, i.e., for $\sigma \in \Sigma(a : b, l : m)$, we use the same notation σ to refer to the set $\{\sigma(i) \mid i = a, \dots, b\}$. The set $\Sigma(0 : k, 0 : n)$ will be mainly used for the description of sub-simplices, and $\Sigma(1 : k, 1 : n)$ for k differential forms in \mathbb{R}^n . For $\sigma \in \Sigma(0 : k, 0 : n)$, $f_\sigma \in \Delta_k(T)$ is the sub-simplex formed by vertices with index $\{\sigma(0), \dots, \sigma(k)\}$. On the other hand, for $f \in \Delta_k(T)$, the index of its vertices can be sorted in ascending order to get an increasing sequence σ_f .

For $\sigma \in \Sigma(0 : k, 0 : n)$, denote by $\sigma^* \in \Sigma(1 : n - k, 0 : n)$ the complementary map characterized by

$$\sigma \cup \sigma^* = \{0, 1, \dots, n\}.$$

For $\sigma \in \Sigma(1 : k, 1 : n)$, its complementary map $\sigma^c \in \Sigma(1 : n - k, 1 : n)$ satisfies

$$\sigma \cup \sigma^c = \{1, \dots, n\}.$$

For the unique element in $\Sigma(a : b, a : b)$, we simply write it as $[a : b]$.

We follow [38] to introduce notation on the addition and subtraction of increasing sequences. Let $\sigma \in \Sigma(a : b, l : m)$. If $q \in [l : m] \setminus \sigma$, then we write $\sigma + q = q + \sigma$ for the unique element of $\Sigma(a : b + 1, l : m)$ with image $\sigma \cup \{q\}$. In that case, we also write $\epsilon(q, \sigma)$ for the signum of the permutation that orders the sequence $[q, \sigma(a), \dots, \sigma(b)]$ in the ascending order. For $q \in \sigma$, $\sigma - q$ is the unique element in $\Sigma(a : b - 1, l : m)$ s.t. $(\sigma - q) + q = \sigma$.

5.1.2. Differential forms

We consider an n -dimensional domain $\Omega \subset \mathbb{R}^n$. Usually we choose a Cartesian coordinate and describe a point $x = (x_1, \dots, x_n) \in \Omega$ in this coordinate. We also use \mathbb{R}^n to denote the n -dimensional linear vector space, which can be identified with the space of points by identifying a point x with the vector $x = o\vec{x}$. We use ∂_{x_i} as the unit vector from the origin o to point $(0, \dots, 1, \dots, 0)$, which is considered as an element in the tangent space $T_o\Omega$. Its dual basis of $(\mathbb{R}^n)^*$ is denoted by $\{dx_i\}_{i=1}^n$, i.e., $dx_i(\partial_{x_j}) = \delta_{i,j}$. We use the standard inner product of vectors to make \mathbb{R}^n a Hilbert space, which introduces an inner product on $(\mathbb{R}^n)^*$: $\langle dx_i, dx_j \rangle = \delta_{i,j}$, $i, j = 1, \dots, n$. We shall reserve notation $\{dx_i\}_{i=1}^n$ for the orthonormal basis induced by the ambient orthonormal coordinate of \mathbb{R}^n .

A generic basis will be denoted by $\{dy_i\}_{i=1}^n$, which may not be orthonormal. We can find another basis $\{d\hat{y}_i\}_{i=1}^n$ dual to $\{dy_i\}_{i=1}^n$ in the sense that $\langle d\hat{y}_i, dy_j \rangle = \delta_{i,j}$. Indeed let $M = (\langle dy_i, dy_j \rangle)_{i,j=1}^n$. Then $(d\hat{y}_1, \dots, d\hat{y}_n)^T = M^{-1}(dy_1, \dots, dy_n)^T$. When $\{dy_i\}_{i=1}^n$ is orthonormal, $d\hat{y}_i = dy_i$ for $i = 1, \dots, n$ as M is identity.

For a vector space V , we define the space of exterior k -forms as the alternating multilinear functional space on $V^k := \underbrace{V \times \dots \times V}_k$

and denote it by $\text{Alt}^k(V)$ or simply Alt^k if V is clear in the context. By definition, $\text{Alt}^k \subset (V^k)^*$. The best way to study a k -form is through the action on k vectors in V .

Let $\omega \in \text{Alt}^p$ and $\eta \in \text{Alt}^q$, we define the wedge product $\omega \wedge \eta \in \text{Alt}^{p+q}$:

$$(\omega \wedge \eta)(v_1, \dots, v_{p+q}) = \sum_{\sigma} \text{sign}(\sigma) \omega(v_{\sigma(1)}, \dots, v_{\sigma(p)}) \eta(v_{\sigma(p+1)}, \dots, v_{\sigma(p+q)}),$$

where the sum is over all permutations σ of $\{1, \dots, p + q\}$, for which $\sigma(1) < \sigma(2) < \dots < \sigma(p)$, $\sigma(p + 1) < \sigma(p + 2) < \dots < \sigma(p + q)$, and $\text{sign}(\sigma)$ is the signature of the permutation σ . We have the determinant formula on the wedge product. For $\omega_i \in V^*$, $v_i \in V$, $i = 1, \dots, p$,

$$(\omega_1 \wedge \dots \wedge \omega_p)(v_1, \dots, v_p) = \det(\omega_i(v_j))_{i,j=1,\dots,p}.$$

For a smooth manifold Ω , a k th order differential form is a section of the tangent bundle $\cup_{x \in \Omega} \text{Alt}^k(T_x\Omega)$, where $T_x\Omega$ is the tangent space at x . The linear space formed by all k th differential forms is denoted by $\Lambda^k(\Omega)$, or simply Λ^k . As Ω is a domain in \mathbb{R}^n , given any point x in the interior of Ω , the tangent space $T_x\Omega$ is isomorphism to $T_o\Omega$ by shifting the origin to x . That is we can use one basis $\{dy_i\}$ for all $\text{Alt}^1(T_x\Omega)$, $x \in \Omega$.

For $\sigma \in \Sigma(1 : k, 1 : n)$, we extend the multi-index notation to write $dy_\sigma \in \text{Alt}^k$:

$$dy_\sigma := dy_{\sigma(1)} \wedge \cdots \wedge dy_{\sigma(k)}.$$

An element $\omega \in \Lambda^k(\Omega)$ thus has a representation

$$\omega = \sum_{\sigma \in \Sigma(1:k, 1:n)} a_\sigma(x) dy_\sigma, \quad x \in \Omega. \quad (42)$$

For a manifold, the basis $\{dy_i\}_{i=1}^n$ is defined on a local chart while in (42), as Ω is flat, a global coordinate is used.

Using (42), we define the exterior derivative $d : \Lambda^k(\Omega) \rightarrow \Lambda^{k+1}(\Omega)$ as: for $\omega = \sum_{\sigma \in \Sigma(1:k, 1:n)} a_\sigma(x) dy_\sigma$, define $d\omega \in \Lambda^{k+1}(\Omega)$ by

$$d\omega = \sum_{\sigma \in \Sigma(1:k, 1:n)} \sum_{i \notin \sigma} \partial_{y_i} a_\sigma dy_i \wedge dy_\sigma = \sum_{\sigma \in \Sigma(1:k, 1:n)} \sum_{i \notin \sigma} \left(\partial_{y_i} a_\sigma \right) \epsilon(i, \sigma) dy_{i+\sigma}.$$

It can be verified that this definition of $d\omega$ is independent of the choice of bases.

The Hodge star for the ambient orthonormal basis $\{dx_i\}_{i=1}^n$ is defined as

$$\star dx_i = (-1)^{i-1} dx_{i^c}, \quad \star dx_{i^c} = (-1)^{n-i} dx_i,$$

which satisfy

$$dx_i \wedge \star dx_i = dx, \quad dx_{i^c} \wedge \star dx_{i^c} = dx,$$

with the volume $dx := dx_1 \wedge dx_2 \wedge \cdots \wedge dx_n$. By definition, $\star \star \omega = (-1)^{n-1} \omega$ for $\omega \in \text{Alt}^1$ or Alt^{n-1} .

We extend the definition to a generic coordinate and define

$$\star dy_i := (-1)^{i-1} dy_{i^c} \quad \text{satisfying} \quad dy_i \wedge \star dy_j = \delta_{i,j} dy.$$

5.1.3. Inner product

An intrinsic definition of an inner product on Alt^k is

$$\langle \omega, \eta \rangle := \sum_{\sigma \in \Sigma(1:k, 1:n)} \omega(e_{\sigma(1)}, \dots, e_{\sigma(k)}) \eta(e_{\sigma(1)}, \dots, e_{\sigma(k)}),$$

where (e_1, \dots, e_n) is any orthonormal basis of \mathbb{R}^n . Then by definition

$$\omega \wedge \star \eta = \langle \omega, \eta \rangle dx, \quad \omega, \eta \in \text{Alt}^k.$$

Recall that $\{dx_i\}_{i=1}^n$ is an orthonormal basis of Alt^1 , i.e., $\langle dx_i, dx_j \rangle = \delta_{i,j}$, $i, j = 1, \dots, n$. It is naturally extended to an orthonormal basis $\{dx_\sigma, \sigma \in \Sigma(1 : k, 1 : n)\}$ of Alt^k , i.e.

$$\langle dx_\sigma, dx_\eta \rangle = \delta_{\sigma, \eta}, \quad \sigma, \eta \in \Sigma(1 : k, 1 : n).$$

The duality of $\{d\hat{y}_i\}_{i=1}^n$ and $\{dy_i\}_{i=1}^n$ are also extended to the k -forms

$$\langle d\hat{y}_\sigma, dy_\eta \rangle = \delta_{\sigma, \eta}, \quad \sigma, \eta \in \Sigma(1 : k, 1 : n). \quad (43)$$

But $\{dy_i\}_{i=1}^n$ may not be orthogonal.

For $\omega, \eta \in \Lambda^k(\Omega)$, a further integral over the domain is included, i.e.,

$$(\omega, \eta)_\Omega = \int_\Omega \langle \omega, \eta \rangle dx, \quad \omega, \eta \in \Lambda^k(\Omega).$$

For a sub-manifold f of Ω , the volume form dx will induce the one for f and denoted by dx_f . Define

$$(\omega, \eta)_f = \int_f \langle \omega, \eta \rangle dx_f.$$

For $\omega, \eta \in \Lambda^k(\Omega)$ with expression

$$\omega = \sum_{\sigma \in \Sigma(1:k, 1:n)} a_\sigma(x) dx_\sigma, \quad \text{and} \quad \eta = \sum_{\sigma \in \Sigma(1:k, 1:n)} b_\sigma(x) dx_\sigma,$$

it is easy to prove that

$$(\omega, \eta)_\Omega = \sum_{\sigma \in \Sigma(1:k, 1:n)} \int_\Omega a_\sigma(x) b_\sigma(x) dx, \quad (44)$$

and (44) is invariant when changing to another orthonormal basis. For non-orthonormal basis, transformation will enter the formulae of the inner product.

Denote by $\mathbb{P}_r \Lambda^k(\Omega)$ the space with polynomial coefficients, and $L^2 \Lambda^k(\Omega)$ is the space with square-integrable coefficient functions. The space $H \Lambda^k(\Omega) := \{\omega \in L^2 \Lambda^k(\Omega) : d\omega \in L^2 \Lambda^{k+1}(\Omega)\}$. When $k = n-1$, $H \Lambda^{n-1}(\Omega)$ is isomorphism to $H(\text{div}, \Omega) := \{v \in L^2(\Omega; \mathbb{R}^n) : \text{div } v \in L^2(\Omega)\}$.

5.1.4. Proxy vectors of differential forms

Representation (42) enables us to identify a differential form with a vector function:

$$\omega \leftrightarrow (a_\sigma)_{\sigma \in \Sigma(1:k,1:n)},$$

and (a_σ) is called a vector proxy of ω . Be aware that, by definition, the differential form is coordinate independent while a vector proxy depends on the coordinate.

For a 1-form $\omega = \sum_{i=1}^n u_i dx_i \in \Lambda^1$, define

$$\text{Prox}_1(\omega) = \mathbf{u} = (u_1, u_2, \dots, u_n)^T.$$

For a vector $\mathbf{t} = (t_1, t_2, \dots, t_n)^T$ representing the tangent vector $\sum_{i=1}^n t_i \partial_{x_i}$, the action is

$$\omega(\mathbf{t}) = \sum_{i,j=1}^n u_i t_j \langle dx_i, \partial_{x_j} \rangle = \mathbf{u} \cdot \mathbf{t}.$$

For $\omega \in \Lambda^{n-1}$, we can write $\omega = \sum_{i=1}^n u_i \star dx_i$, which induces an isomorphism

$$\text{Prox}_{n-1} : \omega \rightarrow \mathbf{u} = (u_1, u_2, \dots, u_n)^T.$$

By definition, we have

$$\begin{aligned} \text{Prox}_1(\star \omega) &= (-1)^{n-1} \text{Prox}_{n-1}(\omega), & \omega &\in \Lambda^{n-1}, \\ \text{Prox}_{n-1}(\star \omega) &= \text{Prox}_1(\omega), & \omega &\in \Lambda^1. \end{aligned}$$

Notice that the proxy vectors are defined using an orthonormal basis. Using the proxy vectors, we can change the wedge product to the inner product of vectors

$$\begin{aligned} \langle \omega, \eta \rangle &= (\text{Prox}_1 \omega, \text{Prox}_{n-1}(\star \eta)), & \omega, \eta &\in \Lambda^1, \\ \langle \omega, \eta \rangle &= (\text{Prox}_k \omega, \text{Prox}_k \eta), & \omega &\in \Lambda^k, \eta \in \Lambda^k, k = 1 \text{ or } n-1, \\ \omega \wedge \eta &= (\text{Prox}_1 \omega, \text{Prox}_{n-1} \eta) dx, & \omega &\in \Lambda^1, \eta \in \Lambda^{n-1}. \end{aligned}$$

For $\omega \in \Lambda^{n-1}$, the representation of $d\omega$ using the proxy vector is $\text{div } \mathbf{u}$, i.e.,

$$d\omega = (\text{div } \mathbf{u}) dx, \quad \text{with } \mathbf{u} = \text{Prox}_{n-1} \omega.$$

The Prox operator is a bijection. More precisely, given a vector $\mathbf{u} = (u_1, \dots, u_n)^T$ represented in the ambient coordinate, let $\omega = \sum_{i=1}^n u_i dx_i \in \Lambda^1$ and $\star \omega = \sum_{i=1}^n u_i \star dx_i \in \Lambda^{n-1}$. Then $\text{Prox}_1(\omega) = \text{Prox}_{n-1}(\star \omega) = \mathbf{u}$. To resemble the notation of differential forms, we introduce notation

$$d\mathbf{u} := \text{Prox}_1^{-1}(\mathbf{u}) = \sum_{i=1}^n u_i dx_i, \quad \star d\mathbf{u} := \text{Prox}_{n-1}^{-1}(\mathbf{u}) = \sum_{i=1}^n u_i \star dx_i.$$

Here in $d\mathbf{u}$, d is understood as a dual operator mapping a tangent vector \mathbf{u} to a co-tangent vector $d\mathbf{u} \in \Lambda^1$, and the symbol d is not associated to any differentiation. A textbook notation of $d\mathbf{u}$ is ${}^b\mathbf{u}$ and $\star d\mathbf{u}$ is ${}^{b_{n-1}}\mathbf{u}$.

Denote the proxy vector of $\{dy_i\}_{i=1}^n$ by $\{\mathbf{v}_i\}_{i=1}^n$ and $\{d\hat{y}_i\}_{i=1}^n$ by $\{\hat{\mathbf{v}}_i\}_{i=1}^n$. Then $\{\hat{\mathbf{v}}_i\}_{i=1}^n$ is dual to $\{\mathbf{v}_i\}_{i=1}^n$ in the sense that $\langle \hat{\mathbf{v}}_i, \mathbf{v}_j \rangle = \delta_{i,j}$. Treat \mathbf{v}_i as a column vector and form the matrix $V = (\mathbf{v}_1, \dots, \mathbf{v}_n)$ and $\hat{V} = (\hat{\mathbf{v}}_1, \dots, \hat{\mathbf{v}}_n)$. The gram matrix is $M = V^T V$. Then we have the relation $\hat{V} = V M^{-1}$.

From

$$\begin{aligned} (\mathbf{v}_i, \text{Prox}_{n-1}(\star dy_j)) dx &= dy_i \wedge \star dy_j = \delta_{i,j} dy = \delta_{i,j} \det(V) dx, \\ (\hat{\mathbf{v}}_i, \text{Prox}_{n-1}(\star d\hat{y}_j)) dx &= d\hat{y}_i \wedge \star d\hat{y}_j = \delta_{i,j} d\hat{y} = \delta_{i,j} \det(\hat{V}) dx, \end{aligned}$$

we get

$$\begin{aligned} \text{Prox}_{n-1}(\star dy_i) &= \det(V) \text{Prox}_1(d\hat{y}_i) = \det(V) \hat{\mathbf{v}}_i, \\ \text{Prox}_{n-1}(\star d\hat{y}_i) &= \det(\hat{V}) \text{Prox}_1(dy_i) = \det(\hat{V}) \mathbf{v}_i, \end{aligned}$$

where the scaling $\det(V)$ or $\det(\hat{V})$ is due to the non-orthogonality. In view of proxy vectors, \star is like a kind of dual operator mapping a vector to its dual vector.

5.1.5. Trace operator

For $F \in \Delta_{n-1}(T)$, let the trace operator $\text{tr}_F^{\text{div}} : \Lambda^{n-1}(T) \rightarrow \Lambda^{n-1}(F)$ be the pullback of the inclusion $F \hookrightarrow T$. That is for any tangent vectors v_1, \dots, v_{n-1} of F , we also treat them as tangent vectors of T and define

$$\text{tr}_F^{\text{div}} \omega(v_1, \dots, v_{n-1}) := \omega(v_1, \dots, v_{n-1}), \quad \omega \in \Lambda^{n-1}(T).$$

We denote by $\text{tr}^{\text{div}} : \Lambda^{n-1}(T) \rightarrow \cup_{F \in \partial T} \Lambda^{n-1}(F)$ as $\text{tr}_F^{\text{div}} \omega|_F = \text{tr}_F^{\text{div}} \omega$.




Fig. 5. Sub-simplices and their tangential and normal vectors.

Let \mathbf{n}_F be the normal vector of F so that the orientation of F , which is given by the volume $\text{d}x_F \in \Lambda^{n-1}(F)$, and \mathbf{n}_F form a consistent orientation of the ambient orthonormal basis. Then

$$\text{tr}_F^{\text{div}} \omega = \mathbf{u} \cdot \mathbf{n}_F \text{d}x_F \quad \text{with } \mathbf{u} = \text{Prox}_{n-1} \omega. \quad (45)$$

On the other hand, for any $p \in L^2(F)$, we have

$$(\omega, p \star \text{d}\mathbf{n}_F)_F = \int_F \mathbf{u} \cdot \mathbf{n}_F p \text{d}x_F = \int_F p \text{tr}_F^{\text{div}} \omega.$$

Based on (45), we can discuss the trace operator in the more familiar vector function setting. The trace operator for space $H(\text{div}, T)$

$$\text{tr}^{\text{div}} : H(\text{div}, T) \rightarrow H^{-1/2}(\partial T)$$

is a continuous extension of $\text{tr}^{\text{div}} \mathbf{u} = \mathbf{u} \cdot \mathbf{n}|_{\partial T}$ defined on smooth functions.

5.1.6. Differential forms in the barycentric coordinates

As $\sum_{i=0}^n \lambda_i = 1$, $\sum_{i=0}^n \text{d}\lambda_i = 0$ and $\{\text{d}\lambda_0, \dots, \text{d}\lambda_n\}$ is not a basis of Alt^1 . Set a vertex as the origin, without loss of generality, say v_0 , then $\{\text{d}\lambda_1, \dots, \text{d}\lambda_n\}$ forms a basis of Alt^1 . In general, through the index σ , there is one-to-one correspondence between $\Delta_{k-1}(F_0)$ and $\text{Alt}^k(T)$. Namely for $\sigma \in \Sigma(1 : k, 1 : n)$, f_σ is a $(k-1)$ -dimensional simplex in $\Delta_{k-1}(F_0)$ with vertices $\{\sigma(1), \dots, \sigma(k)\}$ and $\text{d}\lambda_\sigma = \text{d}\lambda_{\sigma(1)} \wedge \dots \wedge \text{d}\lambda_{\sigma(k)} \in \text{Alt}^k$. We can also write as $\text{d}\lambda_f$ assuming the index of the vertices of f is sorted in the ascending order.

The 1-form $\text{d}\lambda_i$ has a vector representation $\nabla \lambda_i$, which is a scaled normal vector \mathbf{n}_{F_i} of face F_i . For a simplex $f \in \Delta_\ell(T)$, $\{\nabla \lambda_i, i \in f^*\}$ are $n-\ell$ normal vectors of f and can span the normal plane of f . The vector representations of $(n-1)$ -forms, for $i = 1, \dots, \ell$,

$$\text{d}\lambda_{[\sigma(0), \sigma(i)]^*} = \text{d}\lambda_{[0:n]-\sigma(0)-\sigma(i)} := \text{d}\lambda_0 \wedge \dots \wedge \widehat{\text{d}\lambda_{\sigma(0)}} \wedge \dots \wedge \widehat{\text{d}\lambda_{\sigma(i)}} \wedge \dots \wedge \text{d}\lambda_n,$$

are scaling of tangential vectors $\mathbf{t}_{\sigma(0)\sigma(i)}$ of f_σ and can span the tangent plane of f . This is illustrated in Fig. 5.

5.2. An algebraic operator

Define $\text{Alt}^{k,i}(\mathbb{R}^n) = \text{Alt}^k(\mathbb{R}^n) \otimes \text{Alt}^i(\mathbb{R}^n)$ for $k, i = 0, 1, \dots, n$. In particular $\text{Alt}^{k,n-1}(\mathbb{R}^n) = \text{Alt}^k(\mathbb{R}^n) \otimes \text{Alt}^{n-1}(\mathbb{R}^n) \cong \text{Alt}^k(\mathbb{R}^n) \otimes \mathbb{R}^n$. To simplify the notation, $\text{Alt}^{k,i}(\mathbb{R}^n)$ is abbreviated as $\text{Alt}^{k,i}$. In [6], the algebraic operator $s^{k,n-1} : \text{Alt}^{k,n-1} \rightarrow \text{Alt}^{k-1,n}$ is defined as

$$s^{k,n-1} \omega(w_1, \dots, w_{k-1})(v_1, \dots, v_n) := \sum_{i=1}^n (-1)^{i-1} \omega(v_i, w_1, \dots, w_{k-1})(v_1, \dots, \widehat{v_i}, \dots, v_n) \quad \forall v_1, \dots, v_n, w_1, \dots, w_{k-1} \in \mathbb{R}^n.$$

Recall that we have reserved $\{\text{d}x_i\}$ for a fixed orthonormal basis of $\text{Alt}^1(\mathbb{R}^n)$. We are going to derive more concrete forms of operator $s^{k,n-1}$ in a generic basis $\{\text{d}y_i\}$, which may not be orthonormal. We expand $\omega \in \text{Alt}^{k,n-1}$ in this basis as

$$\omega = \sum_{i=1}^n \sum_{\sigma \in \Sigma(1:k, 1:n)} a_{\sigma,i} \text{d}y_\sigma \otimes \text{d}y_i.$$

An element in $\text{Alt}^{k,n-1}$ can be identified as a matrix $\mathbf{A} = (a_{\sigma,i})$ of size $\binom{n}{k} \times n$ indexed by (σ, i) for $i = 1, 2, \dots, n$ and $\sigma \in \Sigma(1 : k, 1 : n)$.

Lemma 5.1. For $\omega = \sum_{i=1}^n \sum_{\sigma \in \Sigma(1:k, 1:n)} a_{\sigma,i} \text{d}y_\sigma \otimes \text{d}y_i$, we have

$$s^{k,n-1} \omega = \sum_{\tau \in \Sigma(1:k-1, 1:n)} \left(\sum_{i \in \tau^c} \epsilon(i, \tau) a_{i+\tau, i} \right) \text{d}y_\tau \otimes \text{d}y_i.$$




Fig. 6. Illustration of constraint sequences and operator $s^{k,n-1}$ for $k=1,3$ and $k=n-1$ for a simplex in \mathbb{R}^5 . For a $(k-2)$ -dimensional sub-simplex τ , $\sigma = i + \tau$ with $i \in \tau^c$ are all $(k-1)$ -dimensional sub-simplex containing τ . For each σ , we attach a vector of length n and consider its i th component, which is a representation of $dy_\sigma \otimes dy_i$. The constraint sequence of τ will be formed by all (σ, i) surrounding τ .

Proof. Let $\{\partial y_i\}$ be the basis of the tangent space dual to $\{dy_i\}$. That is $dy_i(\partial y_j) = \delta_{i,j}$ for $1 \leq i, j \leq n$. Given $\sigma \in \Sigma(1 : k, 1 : n)$, we use the notation ∂y_σ to denote k vectors $(\partial y_{\sigma(1)}, \dots, \partial y_{\sigma(k)})$. Then $dy_\sigma(\partial y_{\sigma'}) = \delta_{\sigma,\sigma'}$ for $\sigma, \sigma' \in \Sigma(1 : k, 1 : n)$. To get the coefficient of $s^{k,n-1}\omega$ for the component $dy_\tau \otimes dy_i$, $\tau \in \Sigma(1 : k-1, 1 : n)$, we check the action

$$\begin{aligned} s^{k,n-1}\omega(\partial y_{\tau(1)}, \dots, \partial y_{\tau(k-1)})(\partial y_1, \dots, \partial y_n) &= \sum_{i=1}^n (-1)^{i-1} \omega(\partial y_i, \partial y_{\tau(1)}, \dots, \partial y_{\tau(k-1)})(\partial y_1, \dots, \widehat{\partial y_i}, \dots, \partial y_n) \\ &= \sum_{i \in \tau^c} \omega(\epsilon(i, \tau) dy_{i+\tau})(-1)^{i-1} \partial y_{i^c} = \sum_{i \in \tau^c} \epsilon(i, \tau) a_{i+\tau, i}. \end{aligned}$$

If $i \in \tau$, then vectors $\partial y_i, \partial y_{\tau(1)}, \dots, \partial y_{\tau(k-1)}$ are linearly dependent and thus the term vanishes. So only $i \in \tau^c$ are left in the summation. \square

For a given $\tau \in \Sigma(1 : k-1, 1 : n)$, we call the sequence of index $\{(i_m + \tau, i_m), i_m \in \tau^c, m = 1, 2, \dots, n-k+1\}$, the *constraint sequence* of τ , which can be also written as $\{(\sigma_{i_m}, i_m), m = 1, 2, \dots, n-k+1, \sigma_{i_m} = i_m + \tau\}$. The length of the constraint sequence is $|\tau^c| = n-k+1$. Without loss of generality, we can sort as $i_1 < i_2 < \dots < i_{n-k+1}$. The first one $(i_1 + \tau, i_1)$ will be called the pair index of the constraint sequence.

We provide some visualization of the constraint sequence. The tensor product $dy_\sigma \otimes dy_i$ can be visualized as follows: for each sub-simplex f_σ , we attach a vector of length n . The sub-index i in dy_i corresponds to the i th component of this vector. See 5-edge stars in Fig. 6. We can associate the $(k-1)$ -form dy_τ with the sub-simplex f_τ of dimension $k-2$, then $\{f_{i+\tau}\}_{i \in \tau^c}$ corresponds to all $(k-1)$ -dimensional sub-simplices of F_0 (excluding $f_{0+\tau}$ as index 0 is not used in differential forms) using τ as a boundary face. See Fig. 6.

If we identify entries of the matrix proxy as nodes of a graph, a constraint sequence will define a path of nodes. See Fig. 7. Indices in different constraint sequences are different. Namely for $\tau \neq \tau'$, $(i + \tau, i) \neq (j + \tau', j)$ as either $i \neq j$ or $i + \tau \neq j + \tau'$. On the graph, different constraint sequences will correspond to disjoint paths.

Since only the value $a_{\sigma,i}$ on the constraint sequence will contribute to the image $s^{k,n-1}\omega$, we conclude that

$$s^{k,n-1}(dy_\sigma \otimes dy_i) = 0 \iff i \in \sigma^c.$$

For each row, i.e., for a fixed σ , there are k entries $(\sigma, i), i \in \sigma$ on k different constraint sequences and the rest $n-k$ entries are not in any constraint sequence.

We can identify $\text{Alt}^{k-1,n}$ as a vector in $\mathbb{R}^{\dim \text{Alt}^{k-1}}$. With the matrix and vector representations, the $s^{k,n-1}$ operator induces an operator from matrix A to a vector in $\mathbb{R}^{\dim \text{Alt}^{k-1}}$ and will be still denoted by $s^{k,n-1}$. We collect the coefficients of a constraint sequence and denote by $\mathbf{a}_\tau = (a_{i_m+\tau, i_m})_{m=1,2,\dots,n-k+1}$. Let $\epsilon_\tau = (\epsilon(i_m, \tau))_{m=1,2,\dots,n-k+1}$ be the corresponding sign vector. We can write the operator as

$$(s^{k,n-1}A)_\tau = \mathbf{a}_\tau \cdot \epsilon_\tau, \quad \tau \in \Sigma(1 : k-1, 1 : n).$$

That is the action is along each constraint sequence.

Lemma 5.2. For $k=1, \dots, n-1$, the operator $s^{k,n-1} : \text{Alt}^{k,n-1} \rightarrow \text{Alt}^{k-1,n}$ is onto. And $s^{n,n-1} : \text{Alt}^{n,n-1} \rightarrow \text{Alt}^{n-1,n}$ is a bijection and its proxy $s^{n,n-1}$ is the transpose operator.

Proof. First consider $k=1, \dots, n-1$. By the linearity, it suffices to prove for $\tau = [1, \dots, k-1]$, there exists $\omega = \sum_{i=1}^n \sum_{\sigma \in \Sigma(1:k,1:n)} a_{\sigma,i} dy_\sigma \otimes dy_i$ such that $s^{k,n-1}\omega = dy_\tau \otimes dy$.

For a given τ , we just pick up one $\sigma = i + \tau$ from its constraint sequence and set the coefficient be $\epsilon(i, \tau)$. More precisely, take $a_{[1,\dots,k],k} = \epsilon(k, [1 : k-1]) = (-1)^{k-1}$, and $a_{\sigma,i} = 0$ for the rest. Then

$$s^{k,n-1}\omega = \sum_{\tilde{\tau} \in \Sigma(1:k-1,1:n)} \left(\sum_{i \in \tilde{\tau}^c} \epsilon(i, \tilde{\tau}) a_{i+\tilde{\tau}, i} \right) dy_{\tilde{\tau}} \otimes dy = dy_\tau \otimes dy.$$




Fig. 7. Constraint sequences for $n = 4, k = 2$ with sign $\epsilon(i, \tau)$. A constraint sequence will define a path of nodes. Different constraint sequences will correspond to disjoint paths. The white circle denotes the pair index of each constraint sequence which is a non-free index. Other circles in yellow are free indices. For each row, there are k entries $(\sigma, i), i \in \sigma$ on k different constraint sequences and the rest $n - k$ entries are not in any constraint sequence. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Next consider $k = n$. For $\omega = \sum_{i=1}^n a_i dy_i \otimes \dots \otimes dy_i$, we have

$$s^{n,n-1}\omega = \sum_{\tau \in \Sigma(1:n-1,1:n)} \left(\sum_{i \in \tau^c} \epsilon(i, \tau) a_i \right) dy_\tau \otimes dy = \sum_{i=1}^n a_i \otimes dy_i \otimes dy.$$

Namely $s^{n,n-1}$ maps the row vector (a_1, \dots, a_n) to a column vector $(a_1, \dots, a_n)^T$. \square

If we identify the row and column vector by the transpose, we can also say $s^{n,n-1}$ is the identity operator.

5.3. Constraint tensor spaces

Now we are ready to introduce the tensor space

$$\mathbb{X} := \ker(s^{k,n-1}) \cap \text{Alt}^{k,n-1} = \{\omega \in \text{Alt}^{k,n-1} \mid s^{k,n-1}\omega = 0\}, \quad 1 \leq k \leq n-1.$$

When $k = n$, as $s^{n,n-1}$ is bijection, $\mathbb{X} = \{0\}$ is trivial. So throughout the rest of the paper, we will consider the non-trivial case $1 \leq k \leq n-1$. For a given basis $\{dy_i\}$, it will be more convenient to work on the matrix representation

$$\mathbb{X} := \left\{ \omega = \sum_{i=1}^n \sum_{\sigma \in \Sigma(1:k,1:n)} a_{\sigma,i} dy_\sigma \otimes \dots \otimes dy_i \mid \mathbf{A} = (a_{\sigma,i}) \in \mathbb{R}^{\binom{n}{k} \times n} : \sum_{i \in \tau^c} \epsilon(i, \tau) a_{i+\tau,i} = 0, \quad \forall \tau \in \Sigma(1:k-1,1:n) \right\}.$$

As $s^{k,n-1}$ is surjective,

$$\dim \mathbb{X} = \dim \text{Alt}^{k,n-1} - \dim \text{Alt}^{k-1,n} = n \binom{n}{k} - \binom{n}{k-1} = (n-k) \binom{n+1}{k}. \quad (46)$$

For the orthonormal basis $\{dx_i\}$, we introduce

$$H(\text{div}, \Omega; \mathbb{X}) := \left\{ \mathbf{A} = (a_{\sigma,i}) \in L^2(\Omega, \mathbb{X}) : \text{div } \mathbf{A} \in L^2(\Omega, \mathbb{R}^{\binom{n}{k}}) \right\}$$

with $\text{div } \mathbf{A} := \left(\sum_{i=1}^n \partial_{x_i} a_{\sigma,i} \right)$, i.e., the divergence operator is applied row-wise. Its differential form version is

$$H(d_{n-1}, \Omega; \mathbb{X}) := \left\{ \omega \in L^2 \Lambda^{k,n-1}(\Omega) : s^{k,n-1}\omega = 0, d_{n-1}\omega \in L^2 \Lambda^{k,n}(\Omega) \right\},$$

where the exterior derivative d_{n-1} is applied to the component Λ^{n-1} in $\Lambda^{k,n-1}$. In view of the matrix proxy \mathbf{A} in the orthonormal basis $\{dx_i\}$, the trace on face F

$$\text{tr}_F^{\text{div}} \mathbf{A} = \mathbf{A} \mathbf{n}_F$$

is a column vector of length $\binom{n}{k}$, and should be continuous on the $(n-1)$ -dimensional faces across simplices.

Example 5.3. Consider $k = n-1$. For $\omega = \sum_{i=1}^n \sum_{j=1}^n a_{i,j} \otimes dy_i \otimes \dots \otimes dy_j$, we have

$$s^{n-1,n-1}\omega = \sum_{\tau \in \Sigma(1:n-2,1:n)} \left(\sum_{i \in \tau^c} (-1)^{([1:n] \setminus [i+\tau]) - 1} \epsilon(i, \tau) a_{[1:n] \setminus [i+\tau], i} \right) dy_\tau \otimes dy$$

$$= \sum_{1 \leq i < j \leq n} (-1)^{i+j} (a_{j,i} - a_{i,j}) dy_{(i,j)^c} \otimes dy,$$

where $dy_{(i,j)^c} := dy_1 \wedge \cdots \wedge \widehat{dy}_i \wedge \cdots \wedge \widehat{dy}_j \wedge \cdots \wedge dy_n$. In terms of the matrix proxy, it holds

$$s^{n-1,n-1}(\mathbf{A}) = 2\text{vskw}(\mathbf{A}) \quad \text{with} \quad \mathbf{A} = (a_{i,j})_{n \times n} \in \mathbb{R}^{n \times n},$$

where operator $\text{vskw} : \mathbb{R}^{n \times n} \rightarrow \mathbb{R}^{n(n-1)/2}$ is defined by

$$(\text{vskw}(\mathbf{A}))_\sigma = \frac{1}{2}(-1)^{i+j} (a_{j,i} - a_{i,j}) \quad \text{with} \quad \sigma = [i, j] \in \Sigma(1 : 2, 1 : n).$$

Thus $\mathbb{X} = \mathbb{S}$ consists of all symmetric matrices.

Example 5.4. Consider $k = 1$. For $\omega = \sum_{i=1}^n \sum_{j=1}^n a_{j,i} dy_j \otimes dy_i$, we have

$$s^{1,n-1}\omega = \sum_{i=1}^n a_{i,i} dy.$$

In terms of the matrix proxy,

$$s^{1,n-1}(\mathbf{A}) = \text{trace}(\mathbf{A}).$$

Thus $\mathbb{X} = \mathbb{T}$ is the traceless matrix space.

5.4. Bases of the constraint tensor space

Recall that

$$\text{Alt}^{k,n-1} = \text{span}\{dy_\sigma \otimes dy_i, \sigma \in \Sigma(1 : k, 1 : n), i = 1, \dots, n\}, \quad 1 \leq k \leq n.$$

We shall modify the basis function $dy_\sigma \otimes dy_i$ to get a basis of $\mathbb{X} = \text{Alt}^{k,n-1} \cap \ker(s^{k,n-1})$.

Recall that we consider the non-trivial case: $1 \leq k \leq n-1$ so that the length of the constraint sequence $n-k+1$ is greater than or equal to 2. We shall define an oblique (non-orthogonal) projection operator $\pi_{\mathbb{X}}$ applied to $dy_\sigma \otimes dy_i$ for index pair (σ, i) .

When $i \in \sigma^c$, $s^{k,n-1}(dy_\sigma \otimes dy_i) = 0$, and thus we keep it unchanged, i.e.,

$$\pi_{\mathbb{X}}(dy_\sigma \otimes dy_i) = dy_\sigma \otimes dy_i, \quad i \in \sigma^c.$$

For each constraint sequence $(\sigma_{i_m}, i_m)_{m=1, \dots, n-k+1}$, set (σ_{i_1}, i_1) as the pair index and modify basis functions to, for $m = 1, \dots, n-k+1$,

$$\pi_{\mathbb{X}}(dy_{\sigma_{i_m}} \otimes dy_{i_m}) := dy_{\sigma_{i_m}} \otimes dy_{i_m} - \epsilon(i_m, \tau) \epsilon(i_1, \tau) dy_{\sigma_{i_1}} \otimes dy_{i_1}.$$

In terms of the coefficient vector, $\pi_{\mathbb{X}}$ will map the vector $\mathbf{a}_\tau = (0, \dots, 1, \dots, 0)$ to the vector

$$\tilde{\mathbf{a}}_\tau = (-\epsilon(i_m, \tau) \epsilon(i_1, \tau), 0, \dots, 1, \dots, 0),$$

so that the constraint $\tilde{\mathbf{a}}_\tau \cdot \mathbf{e}_\tau = 0$ is satisfied. By the linear combination, we get the mapping $\pi_{\mathbb{X}} : \text{Alt}^{k,n-1} \rightarrow \mathbb{X}$.

An index (σ, i) will be called a *free index* if $\pi_{\mathbb{X}}(dy_\sigma \otimes dy_i) \neq 0$. By definition, only $\pi_{\mathbb{X}}(dy_{\sigma_{i_1}} \otimes dy_{i_1}) = 0$, i.e., only the pair index of each constraint sequence is not free. Therefore the number of basis functions is reduced by one for each constraint sequence. In total, we remove $\binom{n}{k-1}$ basis functions of $\text{Alt}^{k,n-1}$ and obtain a basis of \mathbb{X} :

$$\mathbb{X} = \text{span}\{\pi_{\mathbb{X}}(dy_\sigma \otimes dy_i), (\sigma, i) \text{ is free}\}.$$

In Section 6, we will use $\pi_{\mathbb{X}}$ to define a t - n decomposition of \mathbb{X} and construct finite element subspaces of $H(\text{div}, \Omega; \mathbb{X})$.

Next we will present intrinsic bases of \mathbb{X} using the barycentric coordinate.

Lemma 5.5. For any $\sigma \in \Sigma(0 : n-k, 0 : n)$ and $i = 1, \dots, n-k$, it holds

$$d\lambda_{\sigma^*} \otimes d\lambda_{[\sigma(0), \sigma(i)]^*} \in \mathbb{X},$$

where

$$d\lambda_{[\sigma(0), \sigma(i)]^*} = d\lambda_{([0:n]-\sigma(0))-\sigma(i)} = d\lambda_0 \wedge \cdots \wedge \widehat{d\lambda}_{\sigma(0)} \wedge \cdots \wedge \widehat{d\lambda}_{\sigma(i)} \wedge \cdots \wedge d\lambda_n.$$

Proof. We treat $\sigma(0)$ as the origin. Let $y_i = \lambda_{\sigma(i)}$ for $i = 1, \dots, n-k$, and $y_{n-k+i} = \lambda_{\sigma^*(i)}$ for $i = 1, \dots, k$. Then $d\lambda_{\sigma^*} \otimes d\lambda_{[\sigma(0), \sigma(i)]^*} = (-1)^{i-1} (dy_{n-k+1} \wedge \cdots \wedge dy_n) \otimes dy_i \in \mathbb{X}$ for $i = 1, \dots, n-k$ as $i \in [n-k+1, \dots, n]^c$. \square

The vector proxy of $d\lambda_{[\sigma(0), \sigma(i)]^*}$ is a scaling of the edge vector $t_{\sigma(0)\sigma(i)}$, which is on the tangent plane of $f_\sigma \in \Delta_{n-k}(T)$. The k -form $d\lambda_{\sigma^*}$ is the volume of the normal plane of f_σ , i.e. \mathcal{N}^{f_σ} of dimension k . Their tensor product for all $f \in \Delta_{n-k}(T)$ forms a basis of \mathbb{X} .

The inner product of k -form $\langle \omega, \eta \rangle$ can be extended to $\text{Alt}^{k,n-1}$ by the tensor product. Define $P_{\mathbb{X}}$ as the orthogonal projection from $\text{Alt}^{k,n-1}$ to \mathbb{X} w.r.t. the inner product $\langle \cdot, \cdot \rangle$.

Lemma 5.6. *The set*

$$\left\{ d\lambda_{\sigma^*} \otimes d\lambda_{[\sigma(0), \sigma(i)]^*} \right\}_{i=1, \dots, n-k}^{\sigma \in \Sigma(0: n-k, 0: n)}$$

in \mathbb{X} is dual to the set

$$\left\{ P_{\mathbb{X}}(\star d\lambda_{\sigma-\sigma(i)} \otimes \star d\lambda_{\sigma(i)}) \right\}_{i=1, \dots, n-k}^{\sigma \in \Sigma(0: n-k, 0: n)}$$

in the sense that: for any $\sigma, \tau \in \Sigma(0: n-k, 0: n)$, and $i, j = 1, \dots, n-k$,

$$\langle d\lambda_{\sigma^*} \otimes d\lambda_{[\sigma(0), \sigma(i)]^*}, P_{\mathbb{X}}(\star d\lambda_{\tau-\tau(j)} \otimes \star d\lambda_{\tau(j)}) \rangle = \delta_{\sigma, \tau} \delta_{i, j}.$$

Proof. Since $d\lambda_{\sigma^*} \otimes d\lambda_{[\sigma(0), \sigma(i)]^*} \in \mathbb{X}$, by definition of $P_{\mathbb{X}}$ and the inner product $\langle \cdot, \cdot \rangle$, it suffices to prove that: for any $\sigma, \tau \in \Sigma(0: n-k, 0: n)$, and $i, j = 1, \dots, n-k$,

$$(d\lambda_{\sigma^*} \otimes d\lambda_{[\sigma(0), \sigma(i)]^*}) \wedge (d\lambda_{\tau-\tau(j)} \otimes d\lambda_{\tau(j)}) \neq 0$$

if and only if

$$\sigma = \tau \text{ and } i = j.$$

By definition,

$$(d\lambda_{\sigma^*} \otimes d\lambda_{[\sigma(0), \sigma(i)]^*}) \wedge (d\lambda_{\tau-\tau(j)} \otimes d\lambda_{\tau(j)}) = (d\lambda_{\sigma^*} \wedge d\lambda_{\tau-\tau(j)}) \otimes (d\lambda_{[\sigma(0), \sigma(i)]^*} \wedge d\lambda_{\tau(j)}).$$

Then $(d\lambda_{\sigma^*} \otimes d\lambda_{[\sigma(0), \sigma(i)]^*}) \wedge (d\lambda_{\tau-\tau(j)} \otimes d\lambda_{\tau(j)}) \neq 0$ is equivalent to

$$\tau(j) \in \{\sigma(0), \sigma(i)\} \quad \text{and} \quad \tau - \tau(j) \subset \sigma.$$

This indicates $\tau \subseteq \sigma$. We finish the proof by the fact τ and σ have the same length. \square

We are in the position to present intrinsic bases of \mathbb{X} using the barycentric coordinates.

Theorem 5.7 (Intrinsic Bases of \mathbb{X}). *The set*

$$\left\{ d\lambda_{\sigma^*} \otimes d\lambda_{[\sigma(0), \sigma(i)]^*} \right\}_{i=1, \dots, n-k}^{\sigma \in \Sigma(0: n-k, 0: n)}$$

is a basis of \mathbb{X} . Its dual basis is

$$\left\{ P_{\mathbb{X}}(\star d\lambda_{\sigma-\sigma(i)} \otimes \star d\lambda_{\sigma(i)}) \right\}_{i=1, \dots, n-k}^{\sigma \in \Sigma(0: n-k, 0: n)}. \quad (47)$$

Proof. The number of the set $\left\{ d\lambda_{\sigma^*} \otimes d\lambda_{[\sigma(0), \sigma(i)]^*} \right\}_{i=1, \dots, n-k}^{\sigma \in \Sigma(0: n-k, 0: n)}$ is $(n-k) \binom{n+1}{k}$, which equals to $\dim \mathbb{X}$, cf. (46). Hence it suffices to prove that they are linearly independent. Assume there exist $c_{\sigma, i} \in \mathbb{R}$ for each $\sigma \in \Sigma(0: n-k, 0: n)$ and $i = 1, \dots, n-k$ such that

$$\sum_{\sigma \in \Sigma(0: n-k, 0: n)} \sum_{i=1}^{n-k} c_{\sigma, i} d\lambda_{\sigma^*} \otimes d\lambda_{[\sigma(0), \sigma(i)]^*} = 0.$$

Then apply the wedge product with $d\lambda_{\tau-\tau(j)} \otimes d\lambda_{\tau(j)}$ for $\tau \in \Sigma(0: n-k, 0: n)$ and $0 \leq j \leq n-k$, due to Lemma 5.6, we obtain $c_{\tau, j} = 0$. As (τ, j) runs over the whole set $\Sigma(0: n-k, 0: n) \times \{1, \dots, n-k\}$, we conclude all $c_{\tau, j}$ vanishes. \square

Example 5.8. When $k = n-1$, f_{σ} is an edge and the vector proxy of $d\lambda_{\sigma^*}$ is a scaling of the tangent vector $t_{\sigma(0)\sigma(1)}$ of f_{σ} . A basis of \mathbb{X} is thus given by $\{d\lambda_{\sigma^*} \otimes d\lambda_{\sigma^*}\}_{\sigma \in \Sigma(0: 1, 0: n)}$, and the dual basis is $\{\text{sym}(\star d\lambda_{\sigma(0)} \otimes \star d\lambda_{\sigma(1)})\}_{\sigma \in \Sigma(0: 1, 0: n)}$. Equivalently, in terms of the vector proxy, a basis of \mathbb{S} is $\{t^e \otimes t^e\}_{e \in \Delta_1(T)}$ and the dual basis is $\{\text{sym}(\mathbf{n}_{F_i} \otimes \mathbf{n}_{F_j})\}_{e = \text{Convex}(v_i, v_j) \in \Delta_1(T)}$, which are crucial in designing the $H(\text{div}; \mathbb{S})$ element [16, 17, 39] and useful in the Regge calculus [40].

Example 5.9. When $k = 1$, f_{σ} is an $(n-1)$ -dimensional face F and the vector proxy of $d\lambda_{\sigma^*}$ is \mathbf{n}_F . In the matrix proxy, a basis of \mathbb{T} is $\{\mathbf{n}_F \otimes t_i^F\}_{i=1, \dots, n-1}^{F \in \Delta_{n-1}(T)}$, which is discovered in [12] and presented in Lemma 4.6.

5.5. Formulae on the projections

We will present an explicit formula on $P_{\mathbb{X}}$. Recall that the basis $\{\hat{d}y_i\}_{i=1}^n$ is dual to $\{dy_i\}_{i=1}^n$ in the sense that $\langle \hat{d}y_i, dy_j \rangle = \delta_{i, j}$ for $i, j = 1, \dots, n$. The duality also holds for corresponding bases of Alt^k ; see (43).

Lemma 5.10. *It holds*

$$\mathbb{X}^{\perp} = \text{span}\{\hat{e}_{\tau}^A, \tau \in \Sigma(1: k-1, 1: n)\}, \quad (48)$$

where

$$\hat{e}_{\tau}^A := \sum_{i \in \tau^c} \epsilon(i, \tau) d\hat{y}_{i+\tau} \otimes * d\hat{y}_i. \quad (49)$$

For $\omega \in \text{Alt}^{k,n-1}$ expanded in the basis $\omega = \sum_{\sigma,i} a_{\sigma,i} d\hat{y}_\sigma \otimes * d\hat{y}_i$, it holds

$$P_{\mathbb{X}^\perp} \omega = \sum_{\tau \in \Sigma(1:k-1,1:n)} \frac{\mathbf{a}_\tau \cdot \mathbf{e}_\tau}{n-k+1} \hat{\mathbf{e}}_\tau^A, \quad (50)$$

where $P_{\mathbb{X}^\perp} := I - P_{\mathbb{X}}$. Consequently for $\sigma \in \Sigma(1:k,1:n)$,

$$(P_{\mathbb{X}} \omega)_{\sigma,i} = \begin{cases} a_{\sigma,i}, & i \in \sigma^c, \\ a_{\sigma,i} - \frac{\mathbf{a}_\tau \cdot \mathbf{e}_\tau}{n-k+1} \epsilon(i, \tau), & i \in \sigma, \tau = \sigma - i. \end{cases} \quad (51)$$

Proof. Let $\eta = \sum_{i=1}^n \sum_{\sigma \in \Sigma(1:k,1:n)} a_{\sigma,i} dy_\sigma \otimes * dy_i \in \mathbb{X}$. Then

$$\langle \hat{\mathbf{e}}_\tau^A, \eta \rangle = \sum_{i \in \tau^c} \epsilon(i, \tau) a_{i+\tau,i} = 0, \quad \tau \in \Sigma(1:k-1,1:n).$$

That is $\hat{\mathbf{e}}_\tau^A \perp \mathbb{X}$. As any two constraint sequences are disjointed, $\{\hat{\mathbf{e}}_\tau^A, \tau \in \Sigma(1:k-1,1:n)\}$ is linear independent and (48) follows from the dimensions match.

Let $\mathbf{e}_\tau^A = \sum_{i \in \tau^c} \epsilon(i, \tau) dy_{i+\tau} \otimes * dy_i$, which also forms a basis of \mathbb{X}^\perp . Formula (50) holds by testing with \mathbf{e}_τ^A

$$\langle \omega, \mathbf{e}_\tau^A \rangle = \mathbf{a}_\tau \cdot \mathbf{e}_\tau, \quad \langle \hat{\mathbf{e}}_\tau^A, \mathbf{e}_\tau^A \rangle = n - k + 1.$$

Combining (50) and $P_{\mathbb{X}} = I - P_{\mathbb{X}^\perp}$ gives (51). \square

The constraint tensor spaces \mathbb{X} and \mathbb{X}^\perp are defined intrinsically using properties of differential forms, which is independent of choices of the basis. In the proof above, we use different bases $\{dy_i\}$ or $\{d\hat{y}_i\}$ for the ease of computing the projection.

6. Geometric decomposition of $H(\text{div})$ -conforming tensors with constraints

In this section, we generalize the geometric decomposition of the $H(\text{div})$ -conforming vector finite element to the $H(\text{div})$ -conforming tensor finite element. We decompose $\mathbb{P}_r(T; \mathbb{X})$ into a direct sum of the tangential bubble subspace and a normal subspace. Then we present DoFs and show the $H(\text{div})$ -conformity and the discrete inf-sup condition.

6.1. Decomposition of the constraint tensor space

We start from the tensor product of the Lagrange element with \mathbb{X} :

$$\mathbb{P}_r(T; \mathbb{X}) = \bigoplus_{\ell=0}^n \bigoplus_{f \in \Delta_\ell(T)} [b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathbb{X}].$$

For an ℓ -dimensional face $f \in \Delta_\ell(T)$, there is a matrix function $\mathbf{A}^f \in \mathbb{R}^{\binom{n}{\ell} \times n}$ satisfying the constraint $s^{k,n-1}(\mathbf{A}^f) = 0$. The vector $H(\text{div})$ element is $k = 0$ for which the matrix \mathbf{A} is degenerated to a vector of length n and no constraint is imposed. For $1 \leq k \leq n-1$, it is the constraint $s^{k,n-1}(\mathbf{A}^f) = 0$ that makes the finite element construction difficult as the constraint and the normal continuity should be satisfied simultaneously.

As before, for a face $f \in \Delta_\ell(T)$, we choose a t - n basis $\{t_1^f, \dots, t_\ell^f, \mathbf{n}_1^f, \dots, \mathbf{n}_{n-\ell}^f\}$, where the set of ℓ tangential vectors $\{t_1^f, \dots, t_\ell^f\}$ is a basis of the tangent plane \mathcal{T}^f of f and the set of $n - \ell$ normal vectors $\{\mathbf{n}_1^f, \dots, \mathbf{n}_{n-\ell}^f\}$ forms a basis of the normal plane \mathcal{N}^f of f . All basis vectors are normalized but may not be orthogonal. We write $i \in \mathcal{T}^f$ and $i \in \mathcal{N}^f$ to emphasize the range of the index.

Inside the subspace \mathcal{T}^f , we can find a basis $\{\hat{t}_1^f, \dots, \hat{t}_\ell^f\}$ dual to $\{t_1^f, \dots, t_\ell^f\}$, i.e., $\hat{t} \in \mathcal{T}^f$ and $(\hat{t}_i, t_j) = \delta_{i,j}$ for $i, j \in \mathcal{T}^f$. Similarly we have a basis $\{\hat{\mathbf{n}}_1^f, \dots, \hat{\mathbf{n}}_{n-\ell}^f\}$ of \mathcal{N}^f and $(\hat{\mathbf{n}}_i, \mathbf{n}_j) = \delta_{i,j}$ for $i, j \in \mathcal{N}^f$. As $\mathcal{T}^f \perp \mathcal{N}^f$, the basis $\{\hat{t}_1^f, \dots, \hat{t}_\ell^f, \hat{\mathbf{n}}_1^f, \dots, \hat{\mathbf{n}}_{n-\ell}^f\}$ is also dual to $\{t_1^f, \dots, t_\ell^f, \mathbf{n}_1^f, \dots, \mathbf{n}_{n-\ell}^f\}$. Let $V = (t_1^f, \dots, t_\ell^f, \mathbf{n}_1^f, \dots, \mathbf{n}_{n-\ell}^f)$ and $\hat{V} = (\hat{t}_1^f, \dots, \hat{t}_\ell^f, \hat{\mathbf{n}}_1^f, \dots, \hat{\mathbf{n}}_{n-\ell}^f)$.

We say the basis $\{dy_i^f\}$ is the basis of Alt^1 corresponding to a t - n basis if

$$\text{Prox}_1(dy_i^f) = \begin{cases} t_i^f & \text{for } i \in \mathcal{T}^f, \\ \mathbf{n}_{i-\ell}^f & \text{for } i \in \mathcal{N}^f. \end{cases}$$

Then its dual $\{d\hat{y}_i^f\}$ has the vector proxy

$$\text{Prox}_1(d\hat{y}_i^f) = \begin{cases} \hat{t}_i^f & \text{for } i \in \mathcal{T}^f, \\ \hat{\mathbf{n}}_{i-\ell}^f & \text{for } i \in \mathcal{N}^f. \end{cases}$$

We extend the domain of Prox and Prox^{-1} to subspaces. For example, $\text{Prox}_{n-1}^{-1} \mathcal{T}^f = \text{span}\{\text{Prox}_{n-1}^{-1}(\hat{t}_i^f), i = 1, \dots, \ell\} = \text{span}\{\text{Prox}_{n-1}^{-1}(t_i^f), i = 1, \dots, \ell\}$. Then

$$\begin{aligned} \text{Alt}^{k,n-1} &= \text{span}\{dy_\sigma^f \otimes * dy_i^f, \sigma \in \Sigma(1:k,1:n), i = 1, \dots, n\} \\ &= (\text{Alt}^k \otimes \text{Prox}_{n-1}^{-1} \mathcal{T}^f) \oplus (\text{Alt}^k \otimes \text{Prox}_{n-1}^{-1} \mathcal{N}^f). \end{aligned}$$

We introduce the concept *normal constraints*. A constraint with the constraint sequence $\{(\sigma_{i_m}, i_m)_{m=1, \dots, n-k+1}\}$ is called a normal constraint if all $i_m \in \mathcal{N}^f$. The normal constraints will be imposed inside the normal component. Recall that we sort the constraint sequence s.t. $i_1 < i_2 < \dots < i_{n-k+1}$ and set (σ_{i_1}, i_1) as the pair index. So for a non-normal constraint, the pair index (σ_{i_1}, i_1) , $i_1 \in \mathcal{T}^f$ is in the tangential component for $\dim f \geq 1$. Also recall that non-pair indices are free indices.

Define, for $f \in \Delta_\ell(T)$ with $0 \leq \ell \leq n$,

$$\mathcal{T}^f(\mathbb{X}) := \text{span}\{\pi_{\mathbb{X}}(\text{dy}_\sigma^f \otimes \text{dy}_i^f), (\sigma, i) \text{ is free}, i \in \mathcal{T}^f\},$$

$$\mathcal{N}^f(\mathbb{X}) := \text{span}\{\pi_{\mathbb{X}}(\text{dy}_\sigma^f \otimes \text{dy}_i^f), (\sigma, i) \text{ is free}, i \in \mathcal{N}^f\}.$$

For $\ell = 0$, i.e., at vertex $v \in \Delta_0(T)$, we understand $\mathcal{T}^v(\mathbb{X}) = \{0\}$ as no tangent plane and $\mathcal{N}^v(\mathbb{X}) = \mathbb{X}$ as $\mathcal{N}^v = \mathbb{R}^n$.

Lemma 6.1. *Given a t - n basis of a face $f \in \Delta_\ell(T)$, we have the following decomposition*

$$\mathbb{X} = \mathcal{T}^f(\mathbb{X}) \oplus \mathcal{N}^f(\mathbb{X}).$$

Their dimensions are

$$\dim \mathcal{T}^f(\mathbb{X}) = \ell \binom{n}{k} + \binom{n-\ell}{n-k+1} - \binom{n}{k-1},$$

$$\dim \mathcal{N}^f(\mathbb{X}) = (n-\ell) \binom{n}{k} - \binom{n-\ell}{n-k+1}.$$

Proof. By construction, the sum is direct. It suffices to count the dimension. The number of constraints is $\dim \text{Alt}^{k-1}$. By the proof of the surjectivity of $s^{k, n-1}$, all constraints are linearly independent.

Therefore

$$\dim \mathbb{X} = \dim \text{Alt}^{n-1} \times \dim \text{Alt}^k - \dim \text{Alt}^{k-1} = n \dim \text{Alt}^k - \dim \text{Alt}^{k-1}.$$

For each normal constraint, it will remove one index in \mathcal{N}^f . So

$$\dim \mathcal{N}^f(\mathbb{X}) = (n-\ell) \times \dim \text{Alt}^k - \# \text{ normal constraints}.$$

If a constraint is non-normal, then the pair index is in the tangential component. So

$$\dim \mathcal{T}^f(\mathbb{X}) = \ell \times \dim \text{Alt}^k - \# \text{ non-normal constraints}.$$

Sum these two and use the fact

$$\# \text{ normal constraints} + \# \text{ non-normal constraints} = \# \text{ all constraints} = \dim \text{Alt}^{k-1}$$

to conclude $\dim \mathbb{X} = \dim \mathcal{T}^f(\mathbb{X}) + \dim \mathcal{N}^f(\mathbb{X})$.

The number of the normal constraints is $\binom{n-\ell}{n-k+1}$ (among $n-\ell$ indices of the normal plane, choose $n-k+1$ to form the constraint sequence $\{i_m, m = 1, \dots, n-k+1\}$) and thus the number of the non-normal constraints is $\binom{n}{k-1} - \binom{n-\ell}{n-k+1}$. \square

6.2. Geometric decomposition of polynomial constraint tensors

Define the bubble polynomial space

$$\mathbb{B}_r(\text{div}, T; \mathbb{X}) := \mathbb{P}_r(T; \mathbb{X}) \cap \ker(\text{tr}^{\text{div}}).$$

There is no bubble polynomial for lower degree $r = 0, 1$.

Lemma 6.2. *We have $\mathbb{B}_0(\text{div}, T; \mathbb{X}) = \mathbb{B}_1(\text{div}, T; \mathbb{X}) = 0$.*

Proof. Take $\omega \in \mathbb{B}_r(\text{div}, T; \mathbb{X})$ with $r = 0, 1$. Since $\{\mathbf{n}_{F_0}, \dots, \mathbf{n}_{F_{j-1}}, \mathbf{n}_{F_{j+1}}, \dots, \mathbf{n}_{F_n}\}$ form a basis of \mathbb{R}^n , and $(\text{tr}_{F_i} \omega)(v_j) = 0$ for $0 \leq i \neq j \leq n$, we get $\omega(v_j) = 0$. Thus, $\omega = 0$. \square

The tangential component contributes to the bubble. The normal component will contribute to the normal trace. Coupled with the bubble polynomials, we define, for $f \in \Delta_\ell(T)$ with $0 \leq \ell \leq n$,

$$\mathbb{B}_r \mathcal{T}^f(\mathbb{X}) := b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{T}^f(\mathbb{X}), \quad \mathbb{B}_r \mathcal{N}^f(\mathbb{X}) := b_f \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathcal{N}^f(\mathbb{X}).$$

Theorem 6.3 (Characterization of Div Bubble Tensors). *For $r \geq 2$, it holds that*

$$\mathbb{B}_r(\text{div}, T; \mathbb{X}) = \bigoplus_{\ell=1}^n \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_r \mathcal{T}^f(\mathbb{X}),$$

and

$$\text{tr}^{\text{div}} : \bigoplus_{\ell=0}^{n-1} \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_r \mathcal{N}^f(\mathbb{X}) \rightarrow \text{tr}^{\text{div}} \mathbb{P}_r(T; \mathbb{X})$$

is a bijection. Consequently

$$\dim \mathbb{B}_r(\text{div}, T; \mathbb{X}) = \sum_{\ell=1}^n \binom{n+1}{\ell+1} \binom{r-1}{\ell} \left[\ell \binom{n}{k} + \binom{n-\ell}{n-k+1} - \binom{n}{k-1} \right].$$

Proof. Notice that the trace operator is applied to the second component in the tensor product $\text{dy}_\sigma^f \otimes * \text{dy}_i^f$ and $\text{tr}_F^{\text{div}}(* \text{dy}_i^f) = \det(V) \mathbf{n}_F \cdot \hat{\mathbf{t}}_i^f \text{dx}_F = 0$ if $i \in \mathcal{T}^f$ and $f \in \Delta(F)$. The modification in $\pi_{\mathbb{X}}(\text{dy}_\sigma^f \otimes * \text{dy}_i^f)$ will use the pair index in \mathcal{T}^f and thus remains the normal trace free. For $f \notin \Delta(F)$, the bubble function $b_f|_F = 0$.

So we have verified $\bigoplus_{\ell=1}^n \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_r \mathcal{T}^f(\mathbb{X}) \subseteq \mathbb{B}_r(\text{div}, T; \mathbb{X})$.

The rest is the same as Lemma 3.5. \square

We propose the following finite element for the constraint tensor \mathbb{X} .

Lemma 6.4. For each $f \in \Delta_\ell(T)$, choose a t - n basis $\{\mathbf{t}_1^f, \dots, \mathbf{t}_\ell^f, \mathbf{n}_1^f, \dots, \mathbf{n}_{n-\ell}^f\}$ for $\ell = 1, \dots, n-1$. Let $\{\text{dy}_i\}$ be the corresponding basis of Alt^1 , and $\{\text{d}\hat{\mathbf{y}}_i\}$ be its dual basis. The shape function space $\mathbb{P}_r(T; \mathbb{X})$ is uniquely determined by the DoFs

$$\omega(\mathbf{v}_i), \quad i = 0, \dots, n, \omega \in \mathbb{X}, \quad (52a)$$

$$(\omega, \eta)_f, \quad \eta \in \mathbb{P}_{r-(\ell+1)}(f) \otimes \{\text{d}\hat{\mathbf{y}}_\sigma^f \otimes * \text{d}\hat{\mathbf{y}}_i^f \mid (\sigma, i) \text{ is free}, i \in \mathcal{N}^f\}, f \in \Delta_\ell(T), \ell = 1, \dots, n-1, \quad (52b)$$

$$(\omega, \eta)_T, \quad \eta \in \mathbb{B}_r(\text{div}, T; \mathbb{X}). \quad (52c)$$

Proof. Recall the duality

$$\langle \text{dy}_\sigma^f \otimes * \text{dy}_i^f, \text{d}\hat{\mathbf{y}}_\eta^f \otimes * \text{d}\hat{\mathbf{y}}_j^f \rangle = \delta_{\sigma, \eta} \delta_{i, j}$$

for all $1 \leq i, j \leq n$, and $\sigma, \eta \in \Sigma(1 : k, 1 : n)$. In $\pi_{\mathbb{X}}(\text{dy}_\sigma^f \otimes * \text{dy}_i^f) = \text{dy}_\sigma^f \otimes * \text{dy}_i^f - \epsilon(i, \tau) \epsilon(i_1, \tau) \text{dy}_{\sigma_1}^f \otimes * \text{dy}_{i_1}^f$, the pair index (σ_1, i_1) is not free and thus the duality still holds for all free indices

$$\langle \pi_{\mathbb{X}}(\text{dy}_\sigma^f \otimes * \text{dy}_i^f), \text{d}\hat{\mathbf{y}}_\eta^f \otimes * \text{d}\hat{\mathbf{y}}_j^f \rangle = \delta_{\sigma, \eta} \delta_{i, j}, \quad (\sigma, i) \text{ and } (\eta, j) \text{ are free.}$$

Now assume (52a)–(52b) vanishes. For $\omega = \sum_{\text{free}(\sigma, i)} c_{(\sigma, i)} b_f \pi_{\mathbb{X}}(\text{dy}_\sigma^f \otimes * \text{dy}_i^f)$, the DoF $(\omega, c_{(\sigma, i)} \text{d}\hat{\mathbf{y}}_\sigma^f \otimes * \text{d}\hat{\mathbf{y}}_i^f)_f = 0$ will imply $c_{(\sigma, i)} = 0$ for all free indices (σ, i) and $i \in \mathcal{N}^f$. Coupling with the property of the bubble function, we can prove by the forward substitution argument for $\ell = 0, 1, \dots, n-1$ (see the proof of Lemma 3.5), all normal components $\mathbb{B}_r \mathcal{N}^f(\mathbb{X})$ for all $f \in \Delta_\ell(T)$ will vanish and thus only tangential components are left, i.e., $\omega \in \bigoplus_{\ell=1}^n \bigoplus_{f \in \Delta_\ell(T)} \mathbb{B}_r \mathcal{T}^f(\mathbb{X})$. By Theorem 6.3, $\omega \in \mathbb{B}_r(\text{div}, T; \mathbb{X})$ and vanishing (52c) will imply $\omega = 0$. \square

DoF (52b) is in the spirit of the Petrov–Galerkin method, where the test function $\text{d}\hat{\mathbf{y}}_\sigma^f \otimes * \text{d}\hat{\mathbf{y}}_i^f$ is different from the trial function $\pi_{\mathbb{X}}(\text{dy}_\sigma^f \otimes * \text{dy}_i^f)$. This change is important as $\text{Prox}(\text{d}\hat{\mathbf{y}}_\sigma^f) \otimes \text{Prox}(* \text{d}\hat{\mathbf{y}}_i^f) = c \text{Prox}(\text{d}\hat{\mathbf{y}}_\sigma^f) \otimes \mathbf{n}_i^f$ for $i \in \mathcal{N}^f$ will contain the normal component only, which will determine the normal trace to ensure the $H(\text{div})$ -conformity.

In view of the vector proxy, usually we can choose an orthonormal basis for \mathcal{T}^f so that $\mathbf{t}_i^f = \hat{\mathbf{t}}_i^f$ for $i \in \mathcal{T}^f$. We use the normal vector $\{\mathbf{n}_i^f\}$ to define DoFs while use its dual basis $\{\hat{\mathbf{n}}_i^f\}$ to expand the shape function.

Similar to [30], we can write out an explicit basis function

$$\phi_\alpha \pi_{\mathbb{X}}(\text{dy}_\sigma^f \otimes * \text{dy}_i^f), \quad \alpha \in \mathbb{T}_r^\ell(\hat{f}), (\sigma, i) \text{ is free}, f \in \Delta_\ell(T), \ell = 0, \dots, n,$$

where ϕ_α is the nodal basis of Lagrange element at lattice point α , and $\mathbb{T}_r^\ell(\hat{f})$ is the set of lattice points whose geometric embedding is in the interior of f .

6.3. $H(\text{div})$ -Conforming finite element spaces

We shall glue local finite element spaces to form an $H(\text{div})$ -conforming subspace of $H(\text{div}, \Omega; \mathbb{X})$ by choosing a global t - n basis $\{\mathbf{t}_1^f, \dots, \mathbf{t}_\ell^f, \mathbf{n}_1^f, \dots, \mathbf{n}_{n-\ell}^f\}$, i.e., depending only on f not the element containing f .

Theorem 6.5 ($H(\text{div})$ -Conforming Finite Element with Global t - n Bases). For each $f \in \Delta_\ell(\mathcal{T}_h)$, $\ell = 0, \dots, n-1$, choose a t - n basis $\{\mathbf{t}_1^f, \dots, \mathbf{t}_\ell^f, \mathbf{n}_1^f, \dots, \mathbf{n}_{n-\ell}^f\}$ depending only on f . Let $\{\text{dy}_i\}$ be the corresponding basis of Alt^1 , and $\{\text{d}\hat{\mathbf{y}}_i\}$ be its dual basis. Then the following DoFs

$$\omega(\mathbf{v}_i), \quad i = 0, \dots, n, \omega \in \mathbb{X}, \quad (53a)$$

$$(\omega, \eta)_f, \quad \eta \in \mathbb{P}_{r-(\ell+1)}(f) \otimes \{\text{d}\hat{\mathbf{y}}_\sigma^f \otimes * \text{d}\hat{\mathbf{y}}_i^f \mid (\sigma, i) \text{ is free}, i \in \mathcal{N}^f\}, f \in \Delta_\ell(\mathcal{T}_h), \ell = 1, \dots, n-1, \quad (53b)$$

$$(\omega, \eta)_T, \quad \eta \in \mathbb{B}_r(\text{div}, T; \mathbb{X}), T \in \mathcal{T}_h, \quad (53c)$$

will determine a space $V_h \subset H(\text{div}, \Omega; \mathbb{X})$.




Fig. 8. The normal constraint block (in blue) and the free block (in green) for an $f \in \Delta_1(T)$ with $n = 4, k = 2$. In the free block, no index is in a normal constraint. In the normal constraint block, each row contains at least one index in a normal constraint. The red circles will contribute to the bubble spaces. The white circle denotes the pair index of each constraint sequence which is not a free index. (For interpretation of the references to color in this figure legend, the reader is referred to the web version of this article.)

Proof. By Lemma 6.4, on each simplex, DoFs (53) will define a function $\omega \in \mathbb{P}_r(T; \mathbb{X})$. We only need to verify the trace is uniquely determined by (53a)–(53b).

For a face $F \in \Delta_{n-1}(\mathcal{T}_h)$, we have a normal vector \mathbf{n}_F depending only on F and the formula

$$\langle \mathrm{tr}_F^{\mathrm{div}} \omega, d\hat{y}_\sigma^f \rangle = \langle \omega, d\hat{y}_\sigma^f \otimes \star dn_F \rangle.$$

For $f \in \Delta(F)$, as $n_F \in \mathcal{N}^f$, we can expand $n_F = \sum_{i=1}^{n-\ell} c_i^f n_i^f$ and $\star dn_F = \sum_{i=1}^{n-\ell} c_i^f \star dn_i^f = \sum_{i \in \mathcal{N}^f} \tilde{c}_i^f \star d\hat{y}_i^f$.

When (σ, i) is free and $i \in \mathcal{N}^f$, $(\omega, \mathrm{d}\hat{y}_\sigma^f \otimes * \mathrm{d}\hat{y}_i^f)_f$ is given by the DoFs (53a)–(53b).

Then consider the case (σ, i) is not free and $i \in \mathcal{N}^f$. Namely $(\sigma, i) = (i_1 + \tau, i_1)$ is a pair index for a constraint sequence. As $i_1 = i \in \mathcal{N}^f$, this constraint is a normal constraint, i.e., all $i_m \in \mathcal{N}^f$ for $m = 1, \dots, n - k + 1$. We can express $\epsilon(i_1, \tau) \mathrm{d}\hat{y}_{\sigma}^* \otimes \mathrm{d}\hat{y}_i^* = \hat{e}_{\tau}^A - \sum_{m=2}^{n-k+1} \epsilon(i_m, \tau) \mathrm{d}\hat{y}_{i_1+\tau}^* \otimes \mathrm{d}\hat{y}_{i_m}^*$, where $\hat{e}_{\tau}^A \in \mathbb{X}^{\perp}$ is defined in (49). As $\omega \in \mathbb{X}$ and $\hat{e}_{\tau}^A \in \mathbb{X}^{\perp}$, $\langle \omega, \hat{e}_{\tau}^A \rangle = 0$ and consequently

$$(\omega, \epsilon(i_1, \tau) d\hat{y}_\sigma^f \otimes * d\hat{y}_i^f)_f = - \sum_{m=2}^{n-k+1} \epsilon(i_m, \tau)(\omega, d\hat{y}_{i_m+\tau}^f \otimes * d\hat{y}_{i_m}^f)_f.$$

Notice that the index $(i_m + \tau, i_m)$ is free and $i_m \in \mathcal{N}^f$ for $m \geq 2$. So $(\omega, d\hat{y}_\sigma^\otimes * d\hat{y}_i^f)_f$ can be also determined by DoFs (53a)–(53b) even (σ, i) is not free.

It follows that

$$(\mathrm{tr}_F^{\mathrm{div}} \omega, \eta)_f, \quad f \in \Delta_\ell(F), \eta \in \mathbb{P}_{r-(\ell+1)}(f) \otimes \mathrm{Alt}^k, \ell = 0, \dots, n-1.$$

can be determined by DoFs (53a)–(53b). As $\text{tr}_I^{\text{div}} \omega \in \mathbb{P}_r(F) \otimes \text{Alt}^k$, by the uni-solvence of the vector Lagrange element, we conclude $\text{tr}_E^{\text{div}} \omega$ is uniquely determined by DoFs (53a)–(53b). \square

DoF (53a) implies the continuity at vertices. We argue that the continuity at vertices is also necessary. Take a vertex in $\Delta_0(T)$, for example v_0 . Then $(\mathbf{a}_{F_i})(v_0)$ is determined by the row vector $(\mathbf{a}_{F_i})|_{F_i} \in \mathcal{K}$ for $i = 1, \dots, n$, where $F_i \in \Delta_{n-1}(T)$ is the face opposite to v_i . If it is continuous on each face not on vertices, the number of elements in $(\mathbf{a}_F)(v_0)$ is $\dim \text{Alt}^k$ for each face. Running i from 1 to n , $\mathbf{a}(v_0)$ is determined by $n \dim \text{Alt}^k$ conditions, which is more than $\dim \mathbb{X}$. In other words, the constraint makes the tensor product of vector DoFs fails and introduce additional smoothness.

6.4. Facewise redistribution

DoFs (53a)–(53b) implies stronger continuity on the normal plane. We shall further redistribute some DoFs to faces $F \in \Delta_{n-1}(T)$.

To do so, we introduce the concept of the *free block* and the *normal constraint block*. For a fixed σ , the row vector $(\sigma, i), i = \ell + 1 : n$ is called a *free row* if no index is in a normal constraint. All free rows will form a sub-matrix called the *free block*. The rest is called the *normal constraint block*, which contains all normal constraints. Do not confuse the free row with the free index. All indices of a free row are free. But a row with all free indices may not be a free row. A free index can be associated to a normal constraint sequence. See the second and third rows in Fig. 8.

For indices in the free block, the pair index is in the tangential component, which will not change the normal trace. For a normal constraint, the pair index is still in the normal constraint block and changing a basis of \mathcal{N}^f may destroy the $H(\text{div})$ -conformity.

Therefore a global basis $\{\mathbf{n}_i^f\}$ of \mathcal{N}^f is chosen to impose the normal constraints for the normal constraint block. For example, DoFs (39b) for $\mathbf{n}_i^T \sigma \mathbf{n}_j$ in the $H(\text{div}, \mathbb{S})$ element are normal constraints.

For a free row, we have a vector in \mathbb{R}^n consists of $[a_{\sigma,1}, \dots, a_{\sigma,\ell}, a_{\sigma,\ell+1}, \dots, a_{\sigma,n}]^T$. The first ℓ components are in the tangential component and their values are determined locally as the element-wise div bubble polynomials. The part $[a_{\sigma,\ell+1}, \dots, a_{\sigma,n}]^T$ is in the normal component and the corresponding DoFs can be redistributed facetwisely by using the face normal basis $\{\mathbf{n}_F, f \subseteq F, F \in \Delta_{n-1}(T)\}$ of \mathcal{N}^f . In short, a free row is just like a vector $H(\text{div})$ element.

Theorem 6.6 (*$H(\text{div})$ -Conforming Finite Element with Face Redistribution*). For each $f \in \Delta_\ell(\mathcal{T}_h)$, $\ell = 0, \dots, n-1$, choose a global t - n basis $\{\mathbf{t}_1^f, \dots, \mathbf{t}_\ell^f, \mathbf{n}_1^f, \dots, \mathbf{n}_{n-\ell}^f\}$ depending only on f . Let $\{\mathbf{d}\mathbf{y}_i\}$ be the corresponding basis of Alt^1 , and $\{\mathbf{d}\hat{\mathbf{y}}_i\}$ be its dual basis. The following DoFs

$$\omega(\mathbf{v}_i), \quad i = 0, \dots, n, \omega \in \mathbb{X}, \quad (54a)$$

$$(\omega, \eta)_f, \quad \eta \in \mathbb{P}_{r-(\ell+1)}(f) \otimes \{\mathbf{d}\hat{\mathbf{y}}_\sigma^f \otimes \mathbf{d}\hat{\mathbf{y}}_i^f \mid (\sigma, i) \text{ is free and in the normal constraint block}\}, \quad (54b)$$

$$f \in \Delta_\ell(\mathcal{T}_h), \ell = 1, \dots, n-1,$$

$$(\text{tr}_F^{\text{div}} \omega, \eta)_f, \quad \eta \in \mathbb{P}_{r-(\ell+1)}(f) \otimes \{\mathbf{d}\hat{\mathbf{y}}_\sigma^f \mid \sigma \text{ is in the free block}\}, F \in \Delta_{n-1}(\mathcal{T}_h), f \in \Delta_\ell(F), \ell = 1, \dots, n-1, \quad (54c)$$

$$(\omega, \eta)_T, \quad \eta \in \mathbb{B}_r(\text{div}, T; \mathbb{X}), T \in \mathcal{T}_h, \quad (54d)$$

will determine a space $V_h \subset H(\text{div}, \Omega; \mathbb{X})$.

Proof. [1] Local unisolvence on an element. We write DoF (54c) as

$$(\text{tr}_F^{\text{div}} \omega, \mathbf{d}\hat{\mathbf{y}}_\sigma^f)_f = (\omega, \mathbf{d}\hat{\mathbf{y}}_\sigma^f \otimes \star \mathbf{d}\mathbf{n}_F^T)_f, \quad f \subseteq F, \quad (55)$$

where $\{\mathbf{n}_F^T, f \subseteq F, F \in \Delta_{n-1}(T)\}$ is the face normal basis of \mathcal{N}^f in element T . As $\{\mathbf{n}_{F_i}^T, i \in f^*\}$ and $\{\mathbf{n}_j^f, j = 1, \dots, n-\ell\}$ are different bases of the same space \mathcal{N}^f , DoF (55) will also determine

$$(\omega, \mathbf{d}\hat{\mathbf{y}}_\sigma^f \otimes \mathbf{d}\hat{\mathbf{y}}_i^f)_f, \quad i = \ell + 1, \dots, n. \quad (56)$$

Together with (54b), we obtain DoFs (53b). The number of DoFs remains the same as in a free row all indices are free. Then we conclude the unisolvence from Theorem 6.5.

[2] Global conformity across elements. The continuity

$$(\text{tr}_F^{\text{div}} \omega, \eta)_f, \quad \eta \in \mathbb{P}_{r-(\ell+1)}(f) \otimes \{\mathbf{d}\hat{\mathbf{y}}_\sigma^f \mid \sigma \text{ is in the normal constraint block}\}$$

is implied by DoFs (54a)–(54b) as the global t - n basis $\{\mathbf{t}_1^f, \dots, \mathbf{t}_\ell^f, \mathbf{n}_1^f, \dots, \mathbf{n}_{n-\ell}^f\}$ depending only on f . Together with DoF (54c), we conclude $\text{tr}_F^{\text{div}} \omega$ is continuous. \square

Remark 6.7. It seems that we can also try to redistribute the rows in the normal constraint block as DoF (56) can be derived from DoF (55). The problem comes from the fact the basis $\{\mathbf{n}_F^T, f \subseteq F, F \in \Delta_{n-1}(T)\}$ is element-dependent. Expand a global basis vector $\mathbf{n}_i^f = \sum_F c_F^T \mathbf{n}_F^T$ will let (56) be element dependent. For example, for $H(\text{div}, \mathbb{S})$ element, $\mathbf{n}_i^T \tau \mathbf{n}_j$ cannot be redistributed to faces. In the above proof, we transfer (54c) to (56) only for the ease of uni-solvence.

Following the management of DoFs presented in [30], we need to set global and local indexing rules for all DoFs. The global numbering rule is similar to the Lagrange interpolation points. Globally, we can divide the DoFs into those that are shared among simplices and those that are not. The DoFs shared among simplices can be further allocated to the respective sub-simplices. For the DoFs situated on a sub-simplex f , we can choose global normal vectors \mathbf{n}_i^f which share a global DoF labeling, and local tangential vectors \mathbf{t}_i^f which have different labeling in different elements.

We count the size of the normal constraint block. The normal constraint block disappears when $\ell \geq k$ as the length of the constraint $n - k + 1$ will be greater than $n - \ell$ the dimension of the normal plane. That is when $\ell \geq k$, all rows are free and corresponding DoFs can be redistributed to faces.

Consider the case $\ell < k$. If $\sigma \in \Sigma(1 : k, 1 : n)$ is in the constraint block, there exists some $i > \ell$ such that $\tau = \sigma - i \in \Sigma(1 : k - 1, 1 : n)$ satisfies $\tau^c \subseteq [\ell + 1 : n]$, which is equivalent to $\sigma^c \subseteq [\ell + 1 : n]$. Hence the number of rows in the constraint block is $\binom{n-\ell}{n-k}$: among all $n - \ell$ indices of the normal plane, choose $n - k$ indices to form σ^c . When $\ell = 0$, $\binom{n}{n-k} = \binom{n}{k}$, i.e., all rows belong to the constraint block. Consequently DoFs at vertices cannot be redistributed facetwisely. We thus give another justification of the continuity at vertices.

6.5. Discrete inf-sup condition

For a smooth tensor $u = (u_\sigma)$ with index $\sigma \in \Sigma(1 : k, 1 : n)$, let $\text{grad } u$ be a tensor with size $\binom{n}{k} \times n$ give by

$$(\text{grad } u)_{\sigma,i} := \partial_{x_i} u_\sigma.$$

Lemma 6.8. *It holds that*

$$\ker(P_{\mathbb{X}} \text{grad}) = \mathbb{P}_0(T; \mathbb{R}^{\binom{n}{k}}) + \mathbb{X}^\perp x.$$

Proof. Noting that $\text{grad}(\mathbb{P}_0(T; \mathbb{R}^{\binom{n}{k}}) + \mathbb{X}^\perp x) = \mathbb{X}^\perp$, hence

$$\mathbb{P}_0(T; \mathbb{R}^{\binom{n}{k}}) + \mathbb{X}^\perp x \subseteq \ker(P_{\mathbb{X}} \text{grad}).$$

By (31) in [6], $\ker(P_{\mathbb{X}} \text{grad}) \subseteq \mathbb{P}_1(T; \mathbb{R}^{\binom{n}{k}})$. Take $c + Ax \in \ker(P_{\mathbb{X}} \text{grad})$. By $\text{grad}(c + Ax) = A$, we have $P_{\mathbb{X}} A = 0$, i.e. $A \in \mathbb{X}^\perp$. Therefore $\ker(P_{\mathbb{X}} \text{grad}) \subseteq \mathbb{P}_0(T; \mathbb{R}^{\binom{n}{k}}) + \mathbb{X}^\perp x$. \square

We introduce notation $R\mathbb{X} := \ker(P_{\mathbb{X}} \text{grad})$. Examples are $R\mathbb{X} = \text{RM}$ the rigid motion for $\mathbb{X} = \mathbb{S}$, and $R\mathbb{X} = \text{RT}$ for $\mathbb{X} = \mathbb{T}$, where $\text{RT} := \mathbb{P}_0(T; \mathbb{R}^n) + x\mathbb{P}_0(T)$ is the lowest Raviart–Thomas element. In general, $R\mathbb{X}$ is the Whitney form $\mathbb{P}_0 A^k + \kappa_{k+1} \mathbb{P}_0 A^{k+1}$, which is another characterization of $\ker(P_{\mathbb{X}} \text{grad})$.

Operator $P_{\mathbb{X}} \text{grad}$ is the proxy of $P_{\mathbb{X}} d := (-1)^{n-1} \star P_{\mathbb{X}} \star d : \Lambda^{k,0} \rightarrow \Lambda^{k,1}$. Indeed $\int_T \omega \wedge P_{\mathbb{X}} d\eta = (-1)^{n-1} \int_T \omega \wedge \star \star d\eta = \int_T \omega \wedge d\eta$, then the integration by parts holds

$$\int_T d\omega \wedge \eta = (-1)^n \int_T \omega \wedge P_{\mathbb{X}} d\eta + \int_{\partial T} \text{tr}^{\text{div}} \omega \wedge \eta \quad (57)$$

for any $\omega \in H(\text{div}, T; \mathbb{X})$ and $\eta \in \Lambda^{k,0}$. In the matrix and vector proxy, we have

$$\int_T (\text{div } \mathbf{A}) \cdot \mathbf{u} \, dx = - \int_T \mathbf{A} : P_{\mathbb{X}} \text{grad } \mathbf{u} \, dx + \int_{\partial T} (\mathbf{A} \mathbf{n}) \cdot \mathbf{u} \, dS.$$

We consider the finite elements defined in Theorem 6.6. Define the global finite element space

$$\begin{aligned} V_h &:= \{\omega_h \in L^2(\Omega; \mathbb{X}) : \omega_h|_T \in \mathbb{P}_r(T; \mathbb{X}) \quad \forall T \in \mathcal{T}_h, \\ &\quad \text{the DoFs (54a)–(54b) is single-valued across } f \in \Delta_\ell(\mathcal{T}_h) \text{ for } \ell = 0, \dots, n-1, \\ &\quad \text{the DoF (54c) is single-valued across } F \in \Delta_{n-1}(\mathcal{T}_h)\}, \\ Q_h &:= \{q_h \in L^2(\Omega; \Lambda^k) : q_h|_T \in \mathbb{P}_{r-1}(T; \Lambda^k) \quad \forall T \in \mathcal{T}_h\}. \end{aligned} \quad (58)$$

Thanks to Theorem 6.6, $V_h \subset H(\text{div}, \Omega; \mathbb{X})$. We are going to verify the discrete inf-sup condition $\text{div } V_h = Q_h$ if $r \geq n+1$. The following characterization of the range of the div operator on the bubble polynomial space is an abstract version of results (30) and (41) established in [12,16,17].

Lemma 6.9. *For each $T \in \mathcal{T}_h$, it holds*

$$\text{div } \mathbb{B}_r(\text{div}, T; \mathbb{X}) = \mathbb{P}_{r-1}(T; \Lambda^k) \cap R\mathbb{X}^\perp. \quad (59)$$

Proof. When $r = 0, 1$, (59) is obviously true as both sides are zero. We thus consider $r \geq 2$.

Apply the integration by parts (57) to get

$$\text{div } \mathbb{B}_r(\text{div}, T; \mathbb{X}) \subseteq (\mathbb{P}_{r-1}(T; \Lambda^k) \cap R\mathbb{X}^\perp).$$

Next we focus on the proof of the equality. For simplicity, write $d\lambda_{\sigma^*} \otimes d\lambda_{[\sigma(0), \sigma(i)]^*}$ as $\phi_{\sigma,i}$ for each $\sigma \in \Sigma(0 : n-k, 0 : n)$ and $i = 1, \dots, n-k$. By Theorem 5.7, $\{\phi_{\sigma,i}\}_{\sigma \in \Sigma(0:n-k,0:n), i=1,\dots,n-k}$ is a basis of \mathbb{X} , whose dual basis (appropriate rescaling of (47)) is denoted by $\{\psi_{\tau,j}\}_{\tau \in \Sigma(0:n-k,0:n), j=1,\dots,n-k}$, that is $\psi_{\tau,j} \in \mathbb{X}$, $\langle \phi_{\sigma,i}, \psi_{\tau,j} \rangle = 1$ for $\sigma = \tau$ and $i = j$, otherwise it vanishes.

Consider the edge $e = e_{\sigma(0), \sigma(i)}$. The vector proxy of $d\lambda_{[\sigma(0), \sigma(i)]^*}$ is proportional to t^e . Coupled with the edge bubble function $b_e = \lambda_{\sigma(0)} \lambda_{\sigma(i)}$, the vector function $b_e t^e$ satisfies

$$\mathbf{n}_F \cdot b_e t^e|_F = 0, \quad F \in \Delta_{n-1}(T),$$

as if the edge $e \not\subseteq F$, then $b_e|_F = 0$; otherwise $\mathbf{n}_F \cdot t^e = 0$. Therefore $\lambda_{\sigma(0)} \lambda_{\sigma(i)} \phi_{\sigma,i} \in \mathbb{B}_2(\text{div}, T; \mathbb{X})$.

If $\text{div } \mathbb{B}_r(\text{div}, T; \mathbb{X}) \subset (\mathbb{P}_{r-1}(T; \Lambda^k) \cap R\mathbb{X}^\perp)$, then there exists $u \in \mathbb{P}_{r-1}(T; \Lambda^k) \cap R\mathbb{X}^\perp$ satisfying $(u, \text{div } \omega)_T = 0$ for any $\omega \in \mathbb{B}_r(\text{div}, T; \mathbb{X})$. Equivalently

$$(P_{\mathbb{X}} \text{grad } u, \omega)_T = 0 \quad \forall \omega \in \mathbb{B}_r(\text{div}, T; \mathbb{X}).$$

By expressing $P_{\mathbb{X}} \text{grad } u = \sum_{\sigma \in \Sigma(0:n-k,0:n)} \sum_{i=1}^{n-k} q^{\sigma,i} \psi_{\sigma,i}$ with $q^{\sigma,i} \in \mathbb{P}_{r-2}(T)$, we choose

$$\omega = \sum_{\sigma \in \Sigma(0:n-k,0:n)} \sum_{i=1}^{n-k} \lambda_{\sigma(0)} \lambda_{\sigma(i)} q^{\sigma,i} \phi_{\sigma,i} \in \mathbb{B}_r(\text{div}, T; \mathbb{X}).$$

Then we have

$$\sum_{\sigma \in \Sigma(0:n-k,0:n)} \sum_{i=1}^{n-k} (\lambda_{\sigma(0)} \lambda_{\sigma(i)} q^{\sigma,i}, q^{\sigma,i})_T = 0.$$

Therefore $q^{\sigma,i} = 0$ for all i and σ and consequently $u = 0$. \square

Employing the same argument as the proof of [Proposition 4.8](#), the discrete inf-sup condition follows from [\(59\)](#).

Lemma 6.10. *Let $r \geq n+1$. Let $V_h \subset H(\text{div}, \Omega; \mathbb{X})$ be the finite element space defined in [Theorem 6.5](#) and Q_h be the space defined in [\(58\)](#). It holds*

$$\|q_h\|_0 \lesssim \sup_{\omega_h \in V_h} \frac{(\text{div } \omega_h, q_h)}{\|\omega_h\|_0 + \|\text{div } \omega_h\|_0} \quad \forall q_h \in Q_h.$$

For $\ell \geq k$, all DoFs can be redistributed facewisely as no normal constraint block exists. We can further modify the DoFs to get the discrete inf-sup condition with degree $r \geq k+1$ relaxing the requirement $r \geq n+1$ for $k = 1, \dots, n-2$.

Theorem 6.11 (*H(div)-Conforming Finite Element with a Better Inf-Sup Condition*). *Let $1 \leq k \leq n-2$ and $r \geq k+1$. For each $f \in \Delta_\ell(\mathcal{T}_h)$, $\ell = 0, \dots, n-1$, choose a global t - n basis $\{t_1^f, \dots, t_\ell^f, n_1^f, \dots, n_{n-\ell}^f\}$ depending only on f . Let $\{dy_i\}$ be the corresponding basis of Alt^1 , and $\{dy_i^*\}$ be its dual basis. The DoFs*

$$\omega(v_i), \quad i = 0, \dots, n, \omega \in \mathbb{X}, \quad (60a)$$

$$(\omega, \eta)_f, \quad \eta \in \mathbb{P}_{r-(\ell+1)}(f) \otimes \{dy_\sigma^f \otimes dy_i^f \mid (\sigma, i) \text{ is free and in the constraint block}\}, f \in \Delta_\ell(\mathcal{T}_h), \ell = 1, \dots, k-1, \quad (60b)$$

$$(\text{tr}_F^{\text{div}} \omega, \eta)_f, \quad \eta \in \mathbb{P}_{r-(\ell+1)}(f) \otimes \{dy_\sigma^f \mid \sigma \text{ is in the free block}\}, F \in \Delta_{n-1}(\mathcal{T}_h), f \in \Delta_\ell(F), \ell = 1, \dots, k-1, \quad (60c)$$

$$(\text{tr}_F^{\text{div}} \omega, \eta)_F, \quad \eta \in [\mathbb{P}_1(F) \oplus (\mathbb{B}_{r,k}(F) \cap \mathbb{P}_1^\perp(F))] \otimes \text{Alt}^k, F \in \Delta_{n-1}(\mathcal{T}_h), \quad (60d)$$

$$(\omega, \eta)_T, \quad \eta \in \mathbb{B}_r(\text{div}, T; \mathbb{X}), T \in \mathcal{T}_h,$$

will determine a space $V_h \subset H(\text{div}, \Omega; \mathbb{X})$.

Proof. The condition $k \leq n-2$ is to ensure

$$\dim \mathbb{B}_{k+1,k}(F) = |\Delta_k(F)| = \binom{n}{k+1} \geq n = \dim \mathbb{P}_1(F) \quad \text{if } k \leq n-2.$$

So that we can modify the face DoF to [\(60d\)](#). Vanishing DoFs [\(60a\)–\(60c\)](#) will imply $\text{tr}_F^{\text{div}} \omega \in \mathbb{B}_{r,k}(F) \otimes \text{Alt}^k$, which can be decomposed into [\(60d\)](#). \square

As $\eta \in \mathbb{P}_1(F) \otimes \text{Alt}^k$ is included in DoF [\(60d\)](#), we acquire the following discrete inf-sup condition by applying the same argument as the proof of [Lemma 6.10](#).

Corollary 6.12. *Let $1 \leq k \leq n-2$ and $r \geq k+1$. Let $V_h \subset H(\text{div}, \Omega; \mathbb{X})$ be the finite element space defined in [Theorem 6.11](#) and Q_h be the space defined in [\(58\)](#). It holds*

$$\|q_h\|_0 \lesssim \sup_{\omega_h \in V_h} \frac{(\text{div } \omega_h, q_h)}{\|\omega_h\|_0 + \|\text{div } \omega_h\|_0} \quad \forall q_h \in Q_h.$$

Declaration of competing interest

The authors declare the following financial interests/personal relationships which may be considered as potential competing interests: Long Chen reports financial support was provided by National Science Foundation. Xuehai Huang reports financial support was provided by National Natural Science Foundation of China. Xuehai Huang reports financial support was provided by Natural Science Foundation of Shanghai. If there are other authors, they declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Data availability

No data was used for the research described in the article.

Acknowledgments

The first author was supported by DMS-2012465 and DMS-2309785. The second author was supported by the National Natural Science Foundation of China Project 12171300, and the Natural Science Foundation of Shanghai, China 21ZR1480500.

References

- [1] Arnold Douglas N, Falk Richard S, Winther Ragnar. Finite element exterior calculus, homological techniques, and applications. *Acta Numer* 2006;15:1–155.
- [2] Arnold Douglas N, Falk Richard S, Winther Ragnar. Finite element exterior calculus: From Hodge theory to numerical stability. *Bull Amer Math Soc* 2010;47(2):281–354.
- [3] Arnold Douglas N. Finite element exterior calculus. CBMS-NSF Regional Conference Series in Applied Mathematics, 93, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA; 2018, p. xi+120.
- [4] Chen Long, Huang Xuehai. Decoupling of mixed methods based on generalized Helmholtz decompositions. *SIAM J Numer Anal* 2018;56(5):2796–825.

- [5] Chen Long, Huang Xuehai. Finite element complexes in two dimensions (in Chinese). *Sci. Sin. Math.* 2024;54:1–34.
- [6] Arnold Douglas N, Hu Kaibo. Complexes from complexes. *Found Comput Math* 2021;21(6):1739–74.
- [7] Johnson C, Mercier B. Some equilibrium finite element methods for two-dimensional elasticity problems. *Numer Math* 1978;30(1):103–16.
- [8] Arnold Douglas N, Douglas Jr Jim, Gupta Chaitan P. A family of higher order mixed finite element methods for plane elasticity. *Numer Math* 1984;45(1):1–22.
- [9] Arnold Douglas N, Winther Ragnar. Mixed finite elements for elasticity. *Numer Math* 2002;92:401–19.
- [10] Quenneville-Belair Vincent. A new approach to finite element simulations of general relativity. Ann Arbor, MI: ProQuest LLC; 2015, p. 113, Thesis (Ph.D.)—University of Minnesota.
- [11] Chen Long, Huang Xuehai. Discrete hessian complexes in three dimensions. In: *The virtual element method and its applications*. SEMA SIMAI springer ser., vol. 31, Springer, Cham; 2022, p. 93–135.
- [12] Hu Jun, Liang Yizhou. Conforming discrete Gradgrad-complexes in three dimensions. *Math Comp* 2021;90(330):1637–62.
- [13] Hu Jun, Zhang Shangyou. Finite element approximations of symmetric tensors on simplicial grids in \mathbb{R}^n : the lower order case. *Math Models Methods Appl Sci* 2016;26(9):1649–69.
- [14] Arnold Douglas N, Awanou Gerard, Winther Ragnar. Finite elements for symmetric tensors in three dimensions. *Math Comp* 2008;77(263):1229–51.
- [15] Adams Scot, Cockburn Bernardo. A mixed finite element method for elasticity in three dimensions. *J Sci Comput* 2005;25(3):515–21.
- [16] Hu Jun, Zhang ShangYou. A family of symmetric mixed finite elements for linear elasticity on tetrahedral grids. *Sci China Math* 2015;58(2):297–307.
- [17] Hu Jun. Finite element approximations of symmetric tensors on simplicial grids in \mathbb{R}^n : the higher order case. *J Comput Math* 2015;33(3):283–96.
- [18] Falk Richard S, Winther Ragnar. Local bounded cochain projections. *Math Comp* 2014;83(290):2631–56.
- [19] Christiansen Snorre H, Munthe-Kaas Hans Z, Owren Brynjulf. Topics in structure-preserving discretization. *Acta Numer* 2011;20:1–119.
- [20] Arnold Douglas N, Falk Richard S, Winther Ragnar. Geometric decompositions and local bases for spaces of finite element differential forms. *Comput Methods Appl Mech Engrg* 2009;198(21–26):1660–72.
- [21] Gopalakrishnan J, García-Castillo LE, Demkowicz LF. Nédélec spaces in affine coordinates. *Comput Math Appl* 2005;49(7–8):1285–94.
- [22] Christiansen Snorre H, Hu Jun, Hu Kaibo. Nodal finite element de Rham complexes. *Numer Math* 2018;139(2):411–46.
- [23] Chen Long, Huang Xuehai. Geometric decompositions of the simplex lattice and smooth finite elements in arbitrary dimension. 2021.
- [24] Chen Long, Huang Xuehai. Finite element de Rham and Stokes complexes in three dimensions. *Math Comp* 2024;93(345):55–110.
- [25] De Siqueira Denise, Devloo Philippe RB, Gomes Sônia M. Hierarchical high order finite element approximation spaces for $H(\text{div})$ and $H(\text{curl})$. In: Kreiss Gunilla, Lötstedt Per, Målqvist Axel, Neytcheva Maya, editors. *Numerical mathematics and advanced applications 2009*. Berlin, Heidelberg: Springer Berlin Heidelberg; 2010, p. 269–76.
- [26] De Siqueira Denise, Devloo Phillipe RB, Gomes Sônia M. A new procedure for the construction of hierarchical high order Hdiv and Hcurl finite element spaces. *J Comput Appl Math* 2013;240:204–14.
- [27] Castro Douglas A, Devloo Philippe RB, Farias Aginaldo M, Gomes Sônia M, de Siqueira Denise, Durán Omar. Three dimensional hierarchical mixed finite element approximations with enhanced primal variable accuracy. *Comput Methods Appl Mech Engrg* 2016;306:479–502.
- [28] Chen Long, Huang Xuehai. Finite elements for div- and divdiv-conforming symmetric tensors in arbitrary dimension. *SIAM J Numer Anal* 2022;60(4):1932–61.
- [29] Hu Kaibo, Lin Ting, Shi Bowen. Finite elements for symmetric and traceless tensors in three dimensions. 2023.
- [30] Chen Chunyu, Chen Long, Huang Xuehai, Wei Huayi. Geometric decomposition and efficient implementation of high order face and edge elements. *Commun Comput Phys* 2024;35(4):1045–72.
- [31] Nicolaides RA. On a class of finite elements generated by Lagrange interpolation. II. *SIAM J Numer Anal* 1973;10:182–9.
- [32] Nédélec JC. A new family of mixed finite elements in \mathbb{R}^3 . *Numer Math* 1986;50:57–81.
- [33] Brezzi Franco, Douglas Jr Jim, Durán Ricardo, Fortin Michel. Mixed finite elements for second order elliptic problems in three variables. *Numer Math* 1987;51(2):237–50.
- [34] Brezzi Franco, Douglas Jim, Marini LD. Two families of mixed finite elements for second order elliptic problems. *Numer Math* 1985;47(2):217–35.
- [35] Stenberg Rolf. A nonstandard mixed finite element family. *Numer Math* 2010;115(1):131–9.
- [36] Costabel Martin, McIntosh Alan. On Bogovskii and regularized Poincaré integral operators for de Rham complexes on Lipschitz domains. *Math Z* 2010;265(2):297–320.
- [37] Huang Xuehai, Zhang Chao, Zhou Yaqian, Zhu Yangxing. New low-order mixed finite element methods for linear elasticity. *Adv Comput Math* 2024;50(2). Paper No. 17, 31.
- [38] Licht Martin W. On basis constructions in finite element exterior calculus. *Adv Comput Math* 2022;48(2). Paper No. 14, 36.
- [39] Chen Long, Hu Jun, Huang Xuehai. Fast auxiliary space preconditioners for linear elasticity in mixed form. *Math Comp* 2018;87(312):1601–33.
- [40] Christiansen Snorre H. On the linearization of Regge calculus. *Numer Math* 2011;119(4):613–40.