

## MODEL THEORY AND THE QWEP CONJECTURE

ISAAC GOLDBRING

ABSTRACT. We observe that Kirchberg’s QWEP conjecture is equivalent to the statement that  $C^*(\mathbb{F})$  is elementarily equivalent to a QWEP  $C^*$  algebra. We also make a few other model-theoretic remarks about WEP and LLP  $C^*$  algebras.

For the sake of simplicity, all  $C^*$  algebras in this note are assumed to be unital.

Suppose that  $B$  is a  $C^*$  algebra and  $A$  is a subalgebra. We say that  $A$  is *relatively weakly injective* in  $B$  if there is a u.c.p. map  $\phi : B \rightarrow A^{**}$  such that  $\phi|_A = \text{id}_A$ ; such a map is referred to as a *weak conditional expectation*. (We view  $A$  as canonically embedded in its double dual.) A  $C^*$  algebra  $A$  is said to have the *weak expectation property* (or be WEP) if it is relatively weakly injective in every extension and  $A$  is said to be *QWEP* if it is the quotient of a WEP algebra.

*Kirchberg’s QWEP conjecture* states that every separable  $C^*$  algebra is QWEP. In the seminal paper [10], Kirchberg proved that the QWEP conjecture is equivalent to the Connes Embedding Problem (CEP), namely that every finite von Neumann algebra embeds into an ultrapower of the hyperfinite  $\text{II}_1$  factor.

If  $\mathbb{F}$  is the free group on countably many generators, then using the fact that  $C^*(\mathbb{F})$  is surjectively universal, we see that the QWEP conjecture is equivalent to the statement that  $C^*(\mathbb{F})$  is QWEP. The main point of this note is to point to an a priori weaker equivalent statement of the QWEP conjecture:

**THEOREM 1.** *The QWEP conjecture is equivalent to the statement that  $C^*(\mathbb{F})$  is elementarily equivalent to a QWEP  $C^*$  algebra.*

---

Received January 6, 2016; received in final form March 17, 2016.  
2010 *Mathematics Subject Classification.* 46L05, 03C98.

Here, two  $C^*$  algebras  $A$  and  $B$  are *elementarily equivalent* if they have the same first-order theories in the sense of model theory. (Here, we work in the signature for *unital*  $C^*$  algebras.)

The next lemma is probably well known to the experts but since we could not locate it in the literature we include a proof here.

**PROPOSITION 2.** *Let  $A$  be a  $C^*$  algebra and  $\omega$  a nonprincipal ultrafilter on some (possibly uncountable) index set.*

- (1) *Suppose that  $A$  is a subalgebra of  $B$  and that  $B$  admits a u.c.p. map into  $A^\omega$  that restricts to the diagonal embedding of  $A$ . Then  $A$  is relatively weakly injective in  $B$ .*
- (2)  *$A$  is relatively weakly injective in  $A^\omega$ .*

*Proof.* Part (1) is almost identical to the easy direction of [10, Corollary 3.2(ii)] except there he works with the corona algebra instead of the ultrapower. In any event, the proof is easy so we give it here: Suppose that  $\phi : B \rightarrow A^\omega$  is a u.c.p. map restricting to the diagonal embedding of  $A$ . Let  $\theta : A^\omega \rightarrow A^{**}$  be the u.c.p. ultralimit map, that is  $\theta((a_n)^\bullet) := \lim_\omega a_n$  (ultra-weak limit). Then the desired weak expectation  $\psi : B \rightarrow A^{**}$  is given by  $\psi := \theta \circ \phi$ . (2) follows immediately from (1) by taking a nonprincipal ultrafilter  $\omega'$  on a big enough index set so as to allow for an embedding  $A^\omega$  into  $A^{\omega'}$  that restricts to the diagonal embedding of  $A$  into  $A^{\omega'}$ .  $\square$

Theorem 1 follows immediately from the following proposition.

**PROPOSITION 3.** *The set of  $C^*$  algebras with QWEP forms an axiomatizable class.*

*Proof.* We use the semantic test for axiomatizability, namely we show that the class of QWEP algebras is closed under isomorphism, ultraproduct, and ultraroot. (See [1, Proposition 5.14].) Clearly the class of QWEP algebras is preserved under isomorphism. To see that it is closed under ultraproducts, it suffices to note that it is closed under products [10, Corollary 3.3(i)] and (obviously) closed under quotients. To see that it is closed under ultraroots, we use the fact that  $A$  is relatively weakly injective in its ultrapower (Proposition 2(2)) together with the fact that QWEP passes to relatively weakly injective unital subalgebras [10, Corollary 3.3(iii)].  $\square$

**REMARK 4.** Proposition 3 is false with QWEP replaced by WEP: in [7] the authors show that the ultrapower of  $\mathcal{B}(H)$  does not have WEP.

**REMARK 5.** Inductive limits of QWEP algebras are again QWEP (see [2, Lemma 13.3.6]), so the class of QWEP algebras is  $\forall\exists$ -axiomatizable.

There is one other model-theoretic way to settle the QWEP conjecture; we refer the reader to [6] for the definition of existential embeddings.

PROPOSITION 6. *The QWEP conjecture is equivalent to the statement that there is a QWEP  $C^*$  algebra  $A$  containing  $C^*(\mathbb{F})$  as a subalgebra such that the inclusion is an existential embedding of operator systems.*

*Proof.* Suppose that  $A$  is as in the conclusion of the proposition. Then there is a u.c.p. embedding  $A \hookrightarrow C^*(\mathbb{F})^\omega$  whose restriction to  $C^*(\mathbb{F})$  is the diagonal embedding. It follows that  $C^*(\mathbb{F})$  is relatively weakly injective in  $A$ , whence it is itself QWEP by the aforementioned result of Kirchberg.  $\square$

The previous proposition appeared as [6, Corollary 2.24] but with QWEP replaced by WEP.

In [3], the authors ask whether or not every  $C^*$  algebra is elementarily equivalent to a nuclear  $C^*$  algebra. It seems that the authors there were unaware of the fact that if their question had a positive answer, then the QWEP conjecture (and hence CEP) would also be settled. Nevertheless, in the forthcoming manuscript [5], the authors settle this question in the negative by showing that neither  $C_r^*(\mathbb{F})$  nor  $\prod_\omega M_n$  have nuclear models.

A question of Kirchberg, first appearing in print in [11], asks something seemingly more modest than the QWEP conjecture: is there an example of a non-nuclear  $C^*$  algebra that has both WEP and the *local lifting property* (LLP)? Indeed, Kirchberg showed that the QWEP conjecture is equivalent to the statement that the LLP implies WEP. Let us call the statement that there exists a non-nuclear  $C^*$  algebra that has both WEP and LLP the *weak QWEP conjecture*.

PROPOSITION 7. *Let  $A$  be either  $C_r^*(\mathbb{F})$  or  $\prod_\omega M_n$ . If  $A$  is elementarily equivalent to a  $C^*$  algebra  $B$  with LLP, then  $B$  yields a positive solution to the weak QWEP conjecture.*

*Proof.* Since  $A$  is QWEP (for the case of  $C_r^*(\mathbb{F})$ , see [2, Proposition 3.3.8]), we have that  $B$  is also QWEP by Proposition 3; since  $B$  has LLP, we see that  $B$  also has WEP [10, Corollary 2.6(ii)].  $B$  is not nuclear from the aforementioned result in [5].  $\square$

We end this note with something only tangentially related. First, a preparatory result.

PROPOSITION 8. *Suppose that  $A$  is a nonseparable  $C^*$  algebra with a cofinal collection of separable subalgebras that have WEP. Then  $A$  has WEP.*

*Proof.* Suppose that  $A \subseteq B$ ; we must show that  $A$  is relatively weakly injective in  $B$ . To see this, for each separable  $C \subseteq A$  with WEP, there is a weak conditional expectation  $\phi_C : B \rightarrow C^{**} \subseteq A^{**}$ . By taking a pointwise-ultraweak limit of  $\phi_C$  as  $C$  ranges over a cofinal family of separable subalgebras with WEP, we get a witness to the fact that  $A$  is relatively weakly injective in  $B$ .  $\square$

The following first appeared as Theorem 3.1 of [4].

**COROLLARY 9.** *The theory of unital  $C^*$  algebras does not have a model companion, meaning that the class of existentially closed  $C^*$  algebras is not axiomatizable.*

*Proof.* Suppose that  $T$  is the model companion of the theory of  $C^*$  algebras, so the models of  $T$  are precisely the existentially closed  $C^*$  algebras. By [6, Corollary 2.4], all models of  $T$  have WEP. Let  $A$  be a model of  $T$ . Then  $A^\omega$  has a cofinal collection of WEP separable subalgebras, namely the separable elementary substructures of  $A^\omega$ . By Proposition 8,  $A^\omega$  has WEP, whence  $A$  is subhomogeneous by [7, Corollary 4.14]. In particular,  $A$  is finite. Since existentially closed  $C^*$  algebras are purely infinite by [6, Corollary 2.7], we have a contradiction.  $\square$

The advantage of the previous proof over the one in [4] is that the latter proof invokes a serious result of Haagerup and Thorbjørnsen [8], while the above proof ultimately relies instead on the (fundamental but more elementary) work of Junge and Pisier [9].

**Acknowledgments.** The work here was partially supported by NSF CAREER Grant DMS-1349399. The author would like to thank Ilijas Farah and Thomas Sinclair for useful discussions around this work.

#### REFERENCES

- [1] I. Ben Yaacov, A. Berenstein, C. W. Henson and A. Usvyatsov, *Model theory for metric structures*, Model theory with applications to algebra and analysis, vol. 2, London Math. Soc. Lecture Note Ser., vol. 350, Cambridge University Press, Cambridge, 2008, pp. 315–427. MR 2436146
- [2] N. P. Brown and N. Ozawa,  *$C^*$ -algebras and finite-dimensional approximations*, Grad. Studies in Math., vol. 88, AMS, Providence, RI, 2008. MR 2391387
- [3] K. Carlson, E. Cheung, I. Farah, A. Gerhardt-Bourke, B. Hart, L. Mezuman, N. Sequeira and A. Sherman, *Omitting types and AF algebras*, Arch. Math. Logic **53** (2014), 157–169. MR 3151403
- [4] C. Eagle, I. Farah, E. Kirchberg and A. Vignati, *Quantifier elimination in  $C^*$  algebras*, preprint.
- [5] I. Farah, B. Hart, M. Lupini, L. Robert, A. Tikuisis, A. Vignati and W. Winter, *The model theory of  $C^*$  algebras*, preprint; available at [arXiv:1602.08072](https://arxiv.org/abs/1602.08072).
- [6] I. Goldbring and T. Sinclair, *On Kirchberg’s embedding problem*, J. Funct. Anal. **269** (2015), 155–198. MR 3345606
- [7] I. Goldbring and T. Sinclair, *Omitting types in operator systems*, to appear in Indiana Univ. Math. J.
- [8] U. Haagerup and S. Thorbjørnsen, *A new application of random matrices:  $\text{Ext}(C_{\text{red}}^*(F_2))$  is not a group*, Ann. of Math. (2) **162** (2005), 711–775. MR 2183281
- [9] M. Junge and G. Pisier, *Bilinear forms on exact operator spaces and  $\mathcal{B}(H) \otimes \mathcal{B}(H)$* , Geom. Funct. Anal. **5** (1995), 329–363. MR 1334870
- [10] E. Kirchberg, *On non-semisplit extensions, tensor products, and exactness of group  $C^*$ -algebras*, Invent. Math. **112** (1993), 449–489. MR 1218321

- [11] N. Ozawa, *On the QWEP conjecture*, Internat. J. Math. **15** (2004), 501–530.  
MR 2072092

ISAAC GOLDBRING, DEPARTMENT OF MATHEMATICS, STATISTICS, AND COMPUTER SCIENCE, UNIVERSITY OF ILLINOIS AT CHICAGO, SCIENCE AND ENGINEERING OFFICES M/C 249, 851 S. MORGAN ST., CHICAGO, IL 60607-7045, USA AND DEPARTMENT OF MATHEMATICS UNIVERSITY OF CALIFORNIA, IRVINE 340 ROWLAND HALL (BLDG.# 400) IRVINE, CA 92697-3875, USA

*E-mail address:* [isaac@math.uci.edu](mailto:isaac@math.uci.edu); *URL:* <http://www.math.uci.edu/~isaac>