


Math 2B : Midterm # 1 Sample

This exam consists of 5 questions. Problems # 1-3 are worth 15 points each and problems # 4 and 5 are worth 20 points each. There is a total of 85 available points. Read directions for each problem carefully. Please show all work needed to arrive at your solutions. Label all graphs. Clearly indicate your final answers.

- 1.) a.) Estimate the area under the graph of $f(x) = x^2 + x$ from $x = 0$ to $x = 3$ using 3 approximating rectangles and left endpoints. Width of each rectangle = 1.

$$\begin{aligned} & 1 \cdot f(0) + 1 \cdot f(1) + 1 \cdot f(2) \\ & = 1 \cdot 0 + 1 \cdot 2 + 1 \cdot 6 = 8. \end{aligned}$$

- b.) Estimate the area under the graph of $f(x) = x - 1$ from $x = 0$ to $x = 6$ using 3 rectangles and midpoint approximation method. width = 2



$$\begin{aligned} & 2 \cdot f(1) + 2 \cdot f(3) + 2 \cdot f(5) \\ & = 2 \cdot 0 + 2 \cdot 2 + 2 \cdot 4 = 12 \end{aligned}$$

- c.) Find an expression for the area under the graph of $f(x) = x^2 + x$ from $x = 2$ to $x = 5$ as a limit of a Riemann sum. (You do not need to evaluate.)

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{5-2}{n} \cdot f(x_i^*)$$

We'll use right endpoints:

$$x_i^* = 2 + i \cdot \left(\frac{5-2}{n} \right)$$

$$\lim_{n \rightarrow \infty} \sum_{i=1}^n \frac{3}{n} \cdot \left[\left(2 + i \frac{3}{n} \right)^2 + \left(2 + i \frac{3}{n} \right) \right]$$

2.) Evaluate each of the following indefinite integrals:

a.) $\int x\sqrt{3x^2-1} dx$ $u = 3x^2 - 1$ $du = 6x dx$ $\frac{1}{6} du = x dx$

$$= \int \frac{1}{6} \sqrt{u} du = \frac{1}{6} \cdot \frac{2}{3} \cdot u^{\frac{3}{2}} + C$$

$$= \frac{1}{9} (3x^2 - 1)^{\frac{3}{2}} + C$$

b.) $\int \frac{(1-\sin^2 x)}{\cos x} dx = \int \frac{\cos^2 x}{\cos x} dx = \int \cos x dx = \sin x + C$

$$\cos^2 x + \sin^2 x = 1, \text{ so } \cos^2 x = 1 - \sin^2 x$$

c.) $\int \sin(7\theta + 5) d\theta$ $u = 7\theta + 5$ $\frac{1}{7} du = d\theta$

$$\int \frac{1}{7} \sin(u) du = -\frac{1}{7} \cos(u) + C$$

$$= -\frac{1}{7} \cos(7\theta + 5) + C$$

3.) a.) Find the average value of the function $f(x) = \tan^3 x \sec^2 x$ on the interval $[0, \frac{\pi}{4}]$.


$$\text{average} = \frac{1}{\frac{\pi}{4} - 0} \int_0^{\frac{\pi}{4}} \tan^3 x \sec^2 x \, dx$$

$$u = \tan x \quad du = \sec^2 x \, dx$$

$$= \frac{4}{\pi} \int_{\tan 0}^{\tan \frac{\pi}{4}} u^3 \, du = \frac{4}{\pi} \int_0^1 u^3 \, du = \frac{4}{\pi} \frac{u^4}{4} \Big|_0^1 = \frac{1}{\pi}$$

b.) A particle moves along a line so that its velocity at time t is $v(t) = |2 - t|$. Find the displacement of the particle during the time period $0 \leq t \leq 3$.

$$\text{Displacement} = \int_0^3 |2 - t| \, dt$$



$$= \text{Area} = \frac{1}{2} \cdot 2 \cdot 2 + \frac{1}{2} \cdot 1 \cdot 1 = \frac{5}{2}$$

4.) a.) Complete the blanks in the following statement of the Fundamental Theorem of Calculus.

Fundamental Theorem of Calculus:

Suppose f is continuous on $[a, b]$.

If $g(x) = \int_a^x f(t) dt$, then $g'(x) = \underline{f(x)}$.

$\int_a^b f(x) dx = \underline{F(b) - F(a)}$, where F is any antiderivative of f .

b.) Use the Fundamental Theorem of Calculus to evaluate the following.

i.) $\frac{d}{dy} \int_2^y \frac{\sin t}{t^2 + 3} dt$

$\frac{\sin y}{y^2 + 3}$

ii.) $\frac{d}{dx} \int_x^{x^4} \sqrt{t} dt$

$\sqrt{x^4} \cdot 4x^3 - \sqrt{x}$

c.) Answer each of the following questions. No work or explanations are needed.

i.) If $f(t)$ is measured in dollars per year and t in years, what are the units of $\int_0^{10} f(t) dt$?
dollars


ii.) True or False: All continuous functions have derivatives.

false (not all)

iii.) True or False: All continuous functions have antiderivatives.

true $\int_0^x f(t) dt$

iv.) Below is the graph of a function $v(t)$. Let $g(x) = \int_0^x v(t) dt$.




Find each of the following:

$g(0) = \underline{0}$ $g(2) = \underline{3}$
 $2 + \frac{1}{2} \cdot 1 \cdot 2$

$g'(1) = \underline{2}$ $g'(4) = \underline{-1}$
FTC 1

5.) Let S be the region bounded by $y = x^3$ and $y = \sqrt{x}$.

a.) Find the area of region S .



Intersect at

$$x^3 = \sqrt{x}$$

$$x = 0 \quad \& \quad x = 1$$

$$\int_0^1 \sqrt{x} - x^3 \, dx$$

$$= \left. \frac{2}{3} x^{\frac{3}{2}} - \frac{x^4}{4} \right|_0^1$$


$$= \left(\frac{2}{3} - \frac{1}{4} \right)$$

Reality check: it is positive.

b.) i.) Find the volume obtained by revolving the region S about the x axis.


$$\begin{aligned} & \int_0^1 \pi (\sqrt{x})^2 - \pi (x^3)^2 dx \\ &= \int_0^1 \pi x - \pi x^6 dx \\ &= \left. \frac{\pi x^2}{2} - \frac{\pi x^7}{7} \right|_0^1 \\ &= \pi \left(\frac{1}{2} - \frac{1}{7} \right) \end{aligned}$$

ii.) Set up an integral to find the volume obtained by revolving the region S about the y axis.
(You do not need to evaluate the integral.)



$$\int_0^1 \pi (y^{\frac{1}{3}})^2 - \pi (y^2)^2 dy$$

iii.) Set up an integral to find the volume obtained by revolving the region S about the line $y = 5$.
(You do not need to evaluate the integral.)


~~$$\int_0^1 \pi (\sqrt{x} + 5)^2 - \pi (x^3 + 5)^2 dx$$~~

$$\int_0^1 \pi (5 - x^3)^2 - \pi (5 - \sqrt{x})^2 dx$$