

# COUNTEREXAMPLES TO MAXIMAL REGULARITY FOR OPERATORS IN DIVERGENCE FORM

SEBASTIAN BECHTEL, CONNOR MOONEY, AND MARK VERAAR

ABSTRACT. In this paper, we present counterexamples to maximal  $L^p$ -regularity for a parabolic PDE. The example is a second-order operator in divergence form with space and time-dependent coefficients. It is well-known from Lions' theory that such operators admit maximal  $L^2$ -regularity on  $H^{-1}$  under a coercivity condition on the coefficients, and without any regularity conditions in time and space. We show that in general one cannot expect maximal  $L^p$ -regularity on  $H^{-1}(\mathbb{R}^d)$  or  $L^2$ -regularity on  $L^2(\mathbb{R}^d)$ .

## 1. INTRODUCTION

Let  $V$  and  $H$  be complex Hilbert spaces such that  $V \hookrightarrow H$  densely and continuously. Identifying  $H^*$  with its dual, we can view  $H$  as a subspace of  $V^*$ , which is the dual of  $V$ . We start with the abstract problem

$$(CP) \quad \begin{cases} u' - \mathcal{A}u = f, & \text{on } (0, 1), \\ u(0) = 0. \end{cases}$$

Here  $\mathcal{A} : (0, 1) \rightarrow \mathcal{L}(V, V^*)$  is strongly measurable and  $f \in L^2(0, 1; V^*)$ . Moreover, we suppose that there are  $\Lambda, \lambda > 0$  such that for all  $t \in (0, 1)$  and  $v \in V$  one has

$$\|\mathcal{A}(t)v\|_{V^*} \leq \Lambda\|v\|_V, \quad \operatorname{Re}\langle \mathcal{A}(t)v, v \rangle \geq \lambda\|v\|_V^2.$$

By Lions' theory [18] it is known that (CP) has a unique weak solution  $u \in H^1(0, 1; V^*) \cap L^2(0, 1; V)$ . More generally (see [8, XVIII.3.5]), if  $f \in L^2(0, 1; V^*) + L^1(0, 1; H)$ , then (CP) has a unique weak solution  $u \in L^2(0, 1; V)$  such that  $u' \in L^2(0, 1; V^*) + L^1(0, 1; H)$ .

In this paper we study two problems concerning the regularity of  $u$ .

**Problem 1.** *Let  $p \in (1, \infty) \setminus \{2\}$ . Under what condition on  $\mathcal{A}$  does the following hold: for all  $f \in L^p(0, 1; V^*)$  there is a unique weak solution  $u$  which is in  $L^p(0, 1; V)$ ?*

Using the equation this also gives  $u \in H^{1,p}(0, 1; V^*)$ . Problem 1 can equivalently be formulated to the question whether there is a constant  $C > 0$  such that for all step functions  $f$  valued in  $H$  and with  $u$  the unique solution to (CP) given by Lions' result for  $p = 2$  one has the estimate

$$\|u\|_{L^p(0,1;V)} \leq C\|f\|_{L^p(0,1;V^*)}.$$

---

*Date:* January 11, 2024.

*2010 Mathematics Subject Classification.* Primary: 35K90; Secondary: 35B65.

*Key words and phrases.* Maximal  $L^p$ -regularity, counterexamples, Lions' problem, divergence form operators.

It is well-known that regularity results fail for the endpoint cases  $p = 1$  and  $p = \infty$  unless  $V = V^*$  and thus  $\mathcal{A}$  is a family of bounded operators on  $V$  (see [6, 13] and also [15, Thm. 17.4.4 & Cor. 17.4.5]). Hence, we can safely concentrate on the case  $p \in (1, \infty)$  in this paper.

The second problem is a variation of Lions' problem [18, p. 68], who originally asked this question for symmetric  $\mathcal{A}$ .

**Problem 2.** *Under what condition on  $\mathcal{A}$  does the following hold: for all  $f \in L^2(0, 1; H)$  the unique weak solution  $u$  satisfies  $u' \in L^2(0, 1; H)$ ?*

Again using the equation one also has  $\mathcal{A}u \in L^2(0, 1; H)$ . However, it is unclear what this tells about the regularity of  $u$  since the domain of the operators  $\mathcal{A}(t)$  in  $H$  are not easy to describe in general.

Both problems have in common (at least if  $p > 2$ ) that they imply regularity of  $u$  such as  $u \in C^\alpha([0, 1]; [H, V]_\lambda)$  for some  $\alpha > 0$  and  $\lambda \in (0, 1)$ . Such properties are for instance useful in the study of non-linear problems. They follow from standard interpolation estimates and Sobolev embeddings. In Lions' general  $L^2$ -setting one can only obtain  $C([0, 1]; H)$  or Hölder regularity of small order, compare with Theorem 2.2.

If  $\mathcal{A}$  is autonomous, that is to say, the family  $\mathcal{A}(t)$  does not depend on  $t$ , then in both problems the answer is affirmative. Indeed, in Problem 1 this can be concluded from Lions' result and [15, Thm. 17.2.31]. Concerning Problem 2, this follows from a result of de Simon [21].

Otherwise, some conditions are needed. In the case of Problem 1 it is sufficient that the mapping  $t \mapsto \mathcal{A}(t) \in \mathcal{L}(V, V^*)$  is (piecewise relatively) continuous [2, 20]. Without any continuity it is only known that there is some  $\varepsilon > 0$  depending on  $\Lambda, \lambda$  such that Problem 1 holds with  $p \in (2 - \varepsilon, 2 + \varepsilon)$ , see Theorem 2.2. For Problem 2 the situation is even more delicate. Based on an abstract counterexample for the Kato square root problem due to McIntosh, Dier constructed a first nonsymmetric counterexample to Problem 2 in his PhD thesis, see also [3, Sec. 5]. Moreover, Fackler constructed a counterexample that is symmetric and  $C^{\frac{1}{2}}$ -Hölder continuous [11]. To the contrary, with slightly more regularity (for instance  $C^{\frac{1}{2} + \varepsilon}$ -Hölder regularity), many positive results were given [1, 4, 7, 9, 12, 14, 20]. We also recommend the survey [3] for an overview of Problem 2.

The counterexamples due to Dier and Fackler are abstract and not differential operators. As is highlighted by the solution to the Kato square root problem [5], the extra structure of a differential operator can be beneficial compared to the general situation. It is hence of interest to find counterexamples to Problems 1 and 2 that are differential operators. For Problem 2 this was explicitly pointed out in [12, Problem 6.1] and [3, Prop. 12.1]. To be more precise with our setting, we work with  $H = L^2(\mathbb{R}^d)$  and  $V = H^1(\mathbb{R}^d)$ . If  $B: [0, 1] \times \mathbb{R}^d \rightarrow \mathbb{C}^{d \times d}$  is elliptic in the sense that there exist constants  $\Lambda, \lambda > 0$  such that for all  $t \in [0, 1]$  and  $x \in \mathbb{R}^d$  one has

$$(1.1) \quad |B(t, x)\xi| \leq \Lambda|\xi|, \quad \operatorname{Re} B(x, t)\xi \cdot \bar{\xi} \geq \lambda|\xi|^2 \quad (\xi \in \mathbb{C}^d),$$

then consider the problem

$$(P) \quad \begin{cases} u' - \operatorname{div}(B\nabla u) = f, & \text{on } (0, 1), \\ u(0) = 0, \end{cases}$$

where  $f \in L^2(0, 1; H^{-1}(\mathbb{R}^d))$ . In the notation of (CP) we have put  $\mathcal{A}(t)v = -\operatorname{div}(B(t)\nabla v)$ . Lions' theory yields a unique solution  $u$  in the regularity class

$H^1(0, 1; L^2(\mathbb{R}^d)) \cap L^2(0, 1; H^1(\mathbb{R}^d))$ . Problems 1 and 2 ask if for all elliptic coefficients one has the regularity  $u \in L^p(0, 1; H^1(\mathbb{R}^d))$  or  $u' \in L^2(0, 1; L^2(\mathbb{R}^d))$  when the forcing term  $f$  is taken from  $L^p(0, 1; H^{-1}(\mathbb{R}^d))$  or  $L^2(0, 1; L^2(\mathbb{R}^d))$ . The answer is negative in both cases and the respective counterexamples will be the content of the present article. More precisely, based on a construction of the second-named author [19], we will obtain equations whose solution fails to have higher  $L^r(L^s)$ -integrability (Theorem 2.3). Based on this result, we produce a counterexample to Problem 1 for every  $p \neq 2$  in Theorem 2.4. In particular, this shows sharpness of Theorem 2.2 which we mentioned briefly above. Concerning Problem 2, we will show in Theorem 2.6 that it fails in the worst possible way for general elliptic problems in divergence form.

## 2. MAXIMAL REGULARITY IN HILBERT SPACES

**2.1. Known results.** We present the results regarding Problem 1 that were already discussed in the introduction.

A function  $u \in L^2(0, 1; V) \cap H^1(0, 1; V^*)$  is called a solution to (CP) if for all  $t \in [0, 1]$ ,

$$u(t) - \int_0^t \mathcal{A}u ds = \int_0^t f ds,$$

where the integrals are defined as Bochner integrals in  $V^*$ .

One can check that  $u$  is a solution to (CP) if and only if for all  $\phi \in C_c^\infty((0, 1); V)$  one has

$$(2.1) \quad - \int_0^1 \langle u(s), \phi'(s) \rangle ds - \int_0^1 \langle \mathcal{A}(s)u(s), \phi(s) \rangle ds = \int_0^1 \langle f(s), \phi(s) \rangle ds.$$

**Theorem 2.1** (Lions). *In the situation of (CP) there is a unique solution  $u \in L^2(0, 1; V) \cap H^1(0, 1; V^*)$ . Constants in the maximal regularity estimate depend only on  $\Lambda$  and  $\lambda$ .*

Based on a perturbation principle for isomorphisms in complex interpolation scales due to Sneiberg [22], Lions' result can be extended to  $p$  close to 2, see [10, Thm. 4.2].

**Theorem 2.2.** *In the situation of Lions' result (Theorem 2.1) there exists  $\varepsilon > 0$  depending on  $\Lambda$ ,  $\lambda$  and the pair  $(V, H)$  such that Problem 1 has a positive solution for  $p \in (2 - \varepsilon, 2 + \varepsilon)$ .*

It will be clear from Theorem 2.4 that one cannot improve the latter to all  $p \in (1, \infty)$ , even if  $f \in L^p(0, 1; H)$ .

**2.2. Failure of  $L^r(L^s)$ -integrability for variational solutions.** The following counterexample is based on a construction due to the second-named author [19] and is the basis for our subsequent counterexamples to Problems 1 and 2.

**Theorem 2.3** (Failure of higher integrability). *Let  $d \geq 2$ . For every  $s, r \in (1, \infty)$  with  $\frac{2}{r} + \frac{d}{s} < \frac{d}{2}$  there exists an elliptic matrix  $B: [0, 1] \times \mathbb{R}^d \rightarrow \mathbb{C}^{d \times d}$  and  $f \in L^\infty(0, 1; L^2(\mathbb{R}^d))$  such that the unique solution  $u$  to (P) satisfies  $u \notin L^r(0, 1; L^s(\mathbb{R}^d))$ .*

*Proof.* Pick  $\frac{2}{r} + \frac{d}{s} < \mu < \frac{d}{2}$ .

**Step 1:** general setup. We repeat part of the construction in [19]. There exist a Lipschitz function  $w: \mathbb{R}^d \rightarrow \mathbb{C}$  and an elliptic matrix  $a: \mathbb{R}^d \rightarrow \mathbb{C}^{d \times d}$  with the

subsequent properties. The functions  $a$  and  $w$  are uniformly bounded, smooth away from zero, Lipschitz in zero, and for all multiindices  $\alpha$ ,  $|x| \geq 1$ ,

$$(2.2) \quad |\partial^\alpha w(x)| \leq C_\alpha |x|^{-|\alpha|-\mu},$$

$$(2.3) \quad |\partial^\alpha a(x)| \leq C_\alpha |x|^{-|\alpha|}.$$

These estimates follow from the definitions of  $w$  and  $a$  given in [19, Sec. 3]. Similar as in [19, Eq. (5), Thm 2.2, and Cor 2.3] define

$$\begin{aligned} B(t, x) &:= a(x/(1-t)^{1/2}) \\ \zeta(t, x) &:= (1-t)^{-\mu/2} e^{-\frac{\mu}{2} \log(1-t)} w(x/(1-t)^{1/2}). \end{aligned}$$

These were constructed so that  $\zeta \in L^2_{\text{loc}}([0, 1) \times \mathbb{R}^d)$ ,  $\nabla \zeta \in L^2_{\text{loc}}([0, 1) \times \mathbb{R}^d)$  and the equation

$$(2.4) \quad \partial_t \zeta(t, x) = \text{div}(B(t, x) \nabla \zeta(t, x))$$

is satisfied pointwise for  $t \in (0, 1)$  and  $x \in \mathbb{R}^d \setminus \{0\}$ . The latter means that

$$(\zeta, \partial_t \phi)_{L^2((0,1) \times \mathbb{R}^d)} = (B \nabla \zeta, \nabla \phi)_{L^2((0,1) \times \mathbb{R}^d)}$$

holds for all  $\phi \in C_c^\infty((0, 1) \times \mathbb{R}^d)$ . It is clear from the constructions that the weak derivative  $\partial_t \zeta$  exists on  $(0, T)$  as an  $L^2(B_R)$ -valued function, for any  $T < 1$  and  $R > 0$ . Now let

$$\begin{aligned} u(t, x) &= t \zeta(t, x) \eta(x), \\ f(t, x) &= \eta(x) \zeta(t, x) - t \sum_{k, \ell=1}^d [B_{k, \ell}(t, x) + B_{\ell, k}(t, x)] \partial_k \zeta(t, x) \partial_\ell \eta(x) \\ &\quad - t \zeta(t, x) \text{div}(B(t, x) \nabla \eta(x)), \end{aligned}$$

where the cut-off function  $\eta \in C_c^\infty(\mathbb{R}^d)$  is such that  $\eta(x) = 1$  for  $|x| \leq 1$ ,  $\eta(x) = 0$  for  $|x| \geq 2$  and  $0 \leq \eta \leq 1$  on  $\mathbb{R}^d$ .

We claim that  $u \in L^2(0, 1; H^1(\mathbb{R}^d)) \cap H^1(0, 1; H^{-1}(\mathbb{R}^d))$ ,  $f \in L^\infty(0, 1; L^2(\mathbb{R}^d))$  and  $u$  is the unique solution to (P). Then, using (2.4) it is elementary to check that for  $t \in (0, 1)$  and  $x \in \mathbb{R}^d \setminus \{0\}$  one pointwise has

$$\begin{cases} \partial_t u(t, x) - \text{div}(B(t, x) \nabla u(t, x)) = f(t, x), \\ u(t, 0) = 0. \end{cases}$$

Moreover, it also holds in distributional sense on  $(0, 1) \times \mathbb{R}^d$ . Hence, by density and (2.1), it is a solution to (P).

Therefore, it remains to check  $u \in L^2(0, 1; H^1(\mathbb{R}^d))$  and  $f \in L^\infty(0, 1; L^2(\mathbb{R}^d))$  in order to find that  $u$  is the unique solution to (P) provided by Theorem 2.1.

**Step 2:**  $u \in L^2(0, 1; H^1(\mathbb{R}^d))$ . Fix  $t \in (0, 1)$  and let  $\alpha$  be a multiindex with  $|\alpha| \leq 1$ . By definition of  $\zeta$  and using (2.2) one has

$$\begin{aligned} \|\partial^\alpha \zeta(t, \cdot)\|_{L^2(B_2)} &\leq (1-t)^{-\frac{\mu+|\alpha|}{2} + \frac{d}{4}} \|\partial^\alpha w\|_{L^2(B_{2/(1-t)^{1/2}})} \\ &\leq (1-t)^{-\frac{\mu+|\alpha|}{2} + \frac{d}{4}} \left[ C_{d,w} + C_0 \|\cdot\|^{-\mu-|\alpha|} \right]_{L^2(B_{2/(1-t)^{1/2}} \setminus B_1)} \\ &\leq (1-t)^{-\frac{\mu+|\alpha|}{2} + \frac{d}{4}} \left[ C_{d,w} + C_0 C_d (1-t)^{\frac{\mu+|\alpha|}{2} - \frac{d}{4}} \right] \\ &\leq C_{d,w} (1-t)^{-\frac{\mu+|\alpha|}{2} + \frac{d}{4}} + C'_{d,w} \end{aligned} \tag{2.5}$$

for some constant  $C'_{d,w}$  only depending on  $d$  and  $w$  and with

$$C_{d,w} = |B_1|^{1/2} \sup_{x \in B_1} (|w(x)| + |\nabla w(x)|).$$

Note that  $C_{d,w}$  is finite since  $w$  is Lipschitz. Recall that  $\frac{d}{4} - \frac{\mu}{2}$  is positive. Thus, by definition of  $u$  and using (2.5) we find

$$\|u(t, \cdot)\|_{L^2(\mathbb{R}^d)} \leq t \|\zeta(t, \cdot)\|_{L^2(B_2)} \leq C_{d,w} + C'_{d,w}.$$

Thus,  $u$  is bounded as an  $L^2(\mathbb{R}^d)$ -valued function and therefore in particular  $u \in L^2(0, 1; L^2(\mathbb{R}^d))$ . Similarly,

$$\begin{aligned} \|\nabla u(t, \cdot)\|_{L^2(\mathbb{R}^d)} &\leq \|\nabla \zeta(t, \cdot)\|_{L^2(B_2)} + \|\nabla \eta\|_\infty \|\zeta(t, \cdot)\|_{L^2(B_2 \setminus B_1)} \\ &\leq C_{d,w} (1-t)^{-\frac{\mu+1}{2} + \frac{d}{4}} + C'_{d,w} + C'_{d,w} \|\eta\|_\infty. \end{aligned}$$

By choice of  $\mu$  we have  $-\frac{\mu+1}{2} + \frac{d}{4} > -\frac{1}{2}$ , therefore also  $\nabla u \in L^2(0, 1; L^2(\mathbb{R}^d))$ .

**Step 3:**  $f \in L^\infty(0, 1; L^2(\mathbb{R}^d))$ . To estimate the norm of  $f$  we consider all parts separately. The fact that  $\eta \zeta \in L^\infty(0, 1; L^2(\mathbb{R}^d))$  follows from (2.5). Due to the support properties of  $\nabla \eta$  and the boundedness of  $B$  it suffices to prove a uniform estimate for  $\|\partial_m \zeta(t, \cdot)\|_{L^2(B_2 \setminus B_1)}$  to estimate  $B_{k,\ell} \partial_m \zeta \partial_n \eta$ . For this calculate with (2.2) that

$$(2.6) \quad \begin{aligned} \|\partial_m \zeta(t, \cdot)\|_{L^2(B_2 \setminus B_1)} &= (1-t)^{-\frac{\mu}{2} - \frac{1}{2}} \|\partial_m w(\cdot / (1-t)^{1/2})\|_{L^2(B_2 \setminus B_1)} \\ &\leq C_1 \|\cdot\|^{-1-\mu} \|w\|_{L^2(B_2 \setminus B_1)}. \end{aligned}$$

It remains to estimate  $\zeta \operatorname{div}(B \nabla \eta)$ . Again, due to the support properties of  $\nabla \eta$ , it suffices to consider  $x \in B_2 \setminus B_1$ . First, by (2.2) and definition of  $\zeta$ ,

$$|\zeta(t, x)| \leq (1-t)^{-\mu/2} |w(x/(1-t)^{1/2})| \leq C_{1,0}.$$

Thus it remains to estimate  $\|\partial^\alpha B_{k,\ell}(t, \cdot) \partial^\beta \eta\|_{L^2(B_2 \setminus B_1)}$  for  $|\alpha| \leq 1$  and  $1 \leq |\beta| \leq 2$ . Since  $B_{k,\ell}$  and  $\partial^\beta \eta$  are uniformly bounded, it is enough to show that  $\|\partial^\alpha B_{k,\ell}(t, \cdot)\|_{L^2(B_2 \setminus B_1)}$  is uniformly bounded for  $|\alpha| = 1$ . Indeed, this follows from the analogous calculation to (2.6) using (2.3) instead of (2.2).

**Step 4:**  $u \notin L^r(0, 1; L^s(\mathbb{R}^d))$ . Similar to (2.5) but using  $B_1 \subseteq B_{1/(1-t)^{1/2}}$ ,

$$\|\zeta(t, \cdot)\|_{L^s(B_1)} = (1-t)^{-\frac{\mu}{2} + \frac{d}{2s}} \|w\|_{L^s(B_{1/(1-t)^{1/2}})} \geq (1-t)^{-\frac{\mu}{2} + \frac{d}{2s}} \|w\|_{L^s(B_1)}.$$

Therefore, with  $t \in (\frac{1}{2}, 1)$ ,

$$\|u(t, \cdot)\|_{L^s(\mathbb{R}^d)} \geq t \|\zeta(t, \cdot)\|_{L^s(B_1)} \geq \frac{1}{2} (1-t)^{-\frac{\mu}{2} + \frac{d}{2s}} \|w\|_{L^s(B_1)}.$$

By choice of  $\mu$ ,  $r$  and  $s$  we have  $-\frac{\mu}{2} + \frac{d}{2s} < -\frac{1}{r}$ , therefore  $u \notin L^r(0, 1; L^s(\mathbb{R}^d))$ .  $\square$

**2.3. Negative result concerning problem 1.** The following result shows that for time-dependent operators in the variational setting, maximal  $L^2$ -regularity cannot be extrapolated to maximal  $L^p$ -regularity for  $p \neq 2$  besides the small interval given in Theorem 2.2. It answers Problem 1 in a negative way in the setting of elliptic differential operators.

**Theorem 2.4** (Failure of extrapolation of maximal  $L^p$ -regularity). *Let  $d \geq 2$ . For every  $p \in (1, \infty) \setminus \{2\}$  there exists  $B : [0, 1] \times \mathbb{R}^d \rightarrow \mathbb{C}^{d \times d}$  elliptic and  $f \in L^p(0, 1; H^{-1}(\mathbb{R}^d))$  such that the unique solution  $u$  to (P) satisfies  $u \notin L^p(0, 1; H^1(\mathbb{R}^d))$ .*

*Proof.* We divide the proof into two cases.

**Case 1:**  $p > 2$ . We appeal to Theorem 2.3. If  $d \geq 3$  put  $r = p$  and  $s = 2^* := \frac{2d}{d-2}$  and if  $d = 2$  put  $r = p$  and  $s > \frac{2p}{p-2}$ . In both cases the condition  $\frac{2}{r} + \frac{d}{s} < \frac{d}{2}$  is satisfied, so that Theorem 2.3 yields  $B : [0, 1] \times \mathbb{R}^d \rightarrow \mathbb{C}^{d \times d}$  elliptic and  $f \in L^\infty(0, 1; L^2(\mathbb{R}^d)) \subseteq L^p(0, 1; L^2(\mathbb{R}^d))$  such that the unique solution  $u$  to (2.3) satisfies  $u \notin L^r(0, 1; L^s(\mathbb{R}^d))$ . This implies  $u \notin L^p(0, 1; H^1(\mathbb{R}^d))$  since otherwise the Sobolev embedding yields a contradiction to the previous assertion.

**Case 2:**  $p < 2$ . We use a duality argument that resembles [16, Thm. 6.2]. By the first part there are  $f \in L^{p'}(0, 1; H^{-1}(\mathbb{R}^d))$ , elliptic coefficients  $B : [0, 1] \times \mathbb{R}^d \rightarrow \mathbb{C}^{d \times d}$  and a solution  $u \in L^2(0, 1; H^1(\mathbb{R}^d)) \cap H^1(0, 1; H^{-1}(\mathbb{R}^d))$  to the equation

$$(2.7) \quad \begin{cases} \partial_t u - \operatorname{div}(B \nabla u) = f, \\ u(0) = 0, \end{cases}$$

such that  $u \notin L^{p'}(0, 1; H^1(\mathbb{R}^d))$ . Assume for the sake of contradiction that for every  $A : [-1, 0] \times \mathbb{R}^d \rightarrow \mathbb{C}^{d \times d}$  and  $\tilde{g} \in L^2(-1, 0; H^{-1}(\mathbb{R}^d))$  there is a unique solution  $\tilde{v} \in L^2(-1, 0; H^1(\mathbb{R}^d)) \cap H^1(-1, 0; H^{-1}(\mathbb{R}^d))$  to the problem

$$(2.8) \quad \begin{cases} \partial_t \tilde{v} - \operatorname{div}(A \nabla \tilde{v}) = \tilde{g}, \\ \tilde{v}(-1) = 0, \end{cases}$$

satisfying the estimate

$$(2.9) \quad \|\partial_t \tilde{v}\|_{L^p(-1, 0; H^{-1}(\mathbb{R}^d))} + \|\tilde{v}\|_{L^p(-1, 0; H^1(\mathbb{R}^d))} \leq C \|\tilde{g}\|_{L^p(-1, 0; H^{-1}(\mathbb{R}^d))},$$

where the constant  $C > 0$  does not depend on  $\tilde{v}$  and  $\tilde{g}$ . By translation the question on  $(-1, 0)$  is equivalent to that on  $(0, 1)$ . Both intervals are related by the transformation  $v \rightarrow -v$ . We write for instance  $\tilde{u}(t) = u(-t)$  to translate  $u$  to a function on  $(-1, 0)$  and vice versa. Specialize  $A = (\tilde{B})^*$  in (2.8). Since  $u(0) = 0 = v(1)$  we can use integration by parts to obtain

$$\begin{aligned} \int_0^1 \langle g, u \rangle dt &= \int_{-1}^0 \langle \tilde{g}, \tilde{u} \rangle dt = \int_{-1}^0 \langle (\tilde{v})', \tilde{u} \rangle + ((\tilde{B})^* \nabla \tilde{v} | \nabla \tilde{u})_2 dt \\ &= \int_0^1 -\langle v', u \rangle + (\nabla v | B \nabla u)_2 dt \\ &= \int_0^1 \langle v, u' \rangle + (\nabla v | B \nabla u)_2 dt. \end{aligned}$$

Plug in (2.7) to deduce

$$\int_0^1 \langle g, u \rangle dt = \int_0^1 \langle v, f \rangle dt.$$

Hence, using (2.9) we can estimate

$$\begin{aligned} \left| \int_0^1 \langle g, u \rangle dt \right| &\leq \|v\|_{L^p(0, 1; H^1(\mathbb{R}^d))} \|f\|_{L^{p'}(0, 1; H^{-1}(\mathbb{R}^d))} \\ &\leq C \|g\|_{L^p(0, 1; H^{-1}(\mathbb{R}^d))} \|f\|_{L^{p'}(0, 1; H^{-1}(\mathbb{R}^d))}. \end{aligned}$$

Since  $g$  was arbitrary, deduce by duality the contradiction  $u \in L^{p'}(0, 1; H^1(\mathbb{R}^d))$ .  $\square$

*Remark 2.5.* For the case  $p > 2$ , we saw that although  $f \in L^\infty(0, 1; L^2(\mathbb{R}^d))$ , the unique solution  $u$  to (P) satisfies  $u \notin L^p(0, 1; H^1(\mathbb{R}^d))$ .

**2.4. Negative result concerning problem 2.** We show that Problem 2 fails in the worst possible way in the case of elliptic operators in divergence form: for every  $\nu \in (1/2, 1]$  there exist coefficients  $B$  and a forcing term  $f \in L^\infty(0, 1; L^2(\mathbb{R}^d))$  such that the unique solution  $u$  to (P) satisfies  $u \notin H^\nu(0, 1; L^2(\mathbb{R}^d))$ . This solves Lions' problem in a negative way also for elliptic operators. It would still be interesting to find a counterexample for  $\nu = 1$  where the coefficients are Hölder continuous of order  $\alpha$  for some  $\alpha \leq 1/2$ , and this would then be an optimal counterexample due to the positive results for which we refer to the survey [3]. The present coefficient function cannot even be continuous, see Remark 2.7. There seems to be some room for improvement in the regularity of  $B$  as the regularity is much worse than  $H^1$  in time.

**Theorem 2.6** (Failure of maximal  $L^2$ -regularity on  $L^2(\mathbb{R}^d)$ ). *Let  $d \geq 2$ . For every  $\nu \in (1/2, 1]$  there exists  $B : [0, 1] \times \mathbb{R}^d \rightarrow \mathbb{C}^{d \times d}$  elliptic and  $f \in L^\infty(0, 1; L^2(\mathbb{R}^d))$  such that the unique solution  $u$  to (P) satisfies  $u \notin H^\nu(0, 1; L^2)$ .*

*Proof.* The strategy is as for the case  $p > 2$  in Theorem 2.4. Let  $\nu \in (1/2, 1]$  and let  $0 < \theta < \frac{1}{2\nu}$ . Then define parameters  $r$  and  $s$  through  $\frac{1}{r} = \frac{1}{2} - \nu\theta$  and  $\frac{1}{s} = \frac{1}{2} - \frac{1-\theta}{d}$ . They satisfy the condition in Theorem 2.4, consequently there is  $B : [0, 1] \times \mathbb{R}^d \rightarrow \mathbb{C}^{d \times d}$  elliptic and  $f \in L^\infty(0, 1; L^2(\mathbb{R}^d))$  such that the unique solution  $u$  to (P) satisfies  $u \notin L^r(0, 1; L^s(\mathbb{R}^d))$ . Suppose that  $u \in H^\nu(0, 1; L^2(\mathbb{R}^d))$ . Since  $u \in L^2(0, 1; H^1(\mathbb{R}^d))$  by Theorem 2.1, complex interpolation (see [15, Theorem 14.7.12]) yields  $u \in H^{\nu\theta}(0, 1; H^{1-\theta}(\mathbb{R}^d))$ . By choice of  $\theta$ , the Sobolev embedding is applicable and gives  $u \in L^r(0, 1; L^s(\mathbb{R}^d))$ , a contradiction.  $\square$

*Remark 2.7.* By construction, the given counterexample is at the same time a counterexample for Problem 1. We have discussed in the introduction that if  $t \mapsto B(t) \in L^\infty(\mathbb{R}^d)$  is (piecewise relatively) continuous, then Problem 1 is automatically true for all  $p \in (1, \infty)$ . Therefore, the function  $B$  in Theorem 2.6 is not continuous as a map into  $L^\infty(\mathbb{R}^d)$ .

The examples show some limitations of what regularity estimates can hold for elliptic operators in divergence form. However, several issues remain for Problems 1 and 2. For instance, the operator  $\mathcal{A}$  used in the above counterexamples is not symmetric/hermitian. We do not know what can be said for the case  $d = 1$ . The only reference we found on the one-dimensional setting is [17], where counterexamples are given in case of  $L^p$ -integrability in time and space for both the divergence and non-divergence setting for the range  $p \in (1, 3/2) \cup (3, \infty)$ . Finally, it would be interesting to see what can be said about Problem 2 if  $\mathcal{A}$  is more regular in time (i.e. continuous or Hölder continuous).

## REFERENCES

- [1] M. ACHACHE and E.M. OUHABAZ. *Lions' maximal regularity problem with  $H^{\frac{1}{2}}$ -regularity in time.* J. Differential Equations **266** (2019), no. 6, 3654–3678. doi:[10.1016/j.jde.2018.09.015](https://doi.org/10.1016/j.jde.2018.09.015)
- [2] W. Arendt, R. Chill, S. Fornaro, and C. Poupaud.  $L^p$ -maximal regularity for nonautonomous evolution equations. *J. Differ. Equations*, 237(1):1–26, 2007.
- [3] W. ARENDT, D. DIER, and S. FACKLER. *J. L. Lions' problem on maximal regularity.* Arch. Math. **109** (2017), no. 1, 59–72. doi:[10.1007/s00013-017-1031-6](https://doi.org/10.1007/s00013-017-1031-6)
- [4] P. AUSCHER and M. EGERT. *On non-autonomous maximal regularity for elliptic operators in divergence form.* Arch. Math. (Basel) **107** (2016), no. 3, 271–284. doi:[10.1007/s00013-016-0934-y](https://doi.org/10.1007/s00013-016-0934-y)

- [5] P. AUSCHER, S. HOFMANN, M. LACEY, A. MCINTOSH, and P. TCHAMITCHIAN. *The solution of the Kato square root problem for second order elliptic operators on  $\mathbb{R}^n$* . Ann. of Math. (2) **156** (2002), no. 2, 633–654. doi:[10.2307/3597201](https://doi.org/10.2307/3597201)
- [6] J-B. BAILLON. *Caractère borné de certains générateurs de semi-groupes linéaires dans les espaces de Banach*. C. R. Acad. Sci. Paris Sér. A-B **290** (1980), no. 16, A757–A760.
- [7] S. BECHTEL. *Weighted non-autonomous  $L^q(L^p)$  maximal regularity for complex systems*. arXiv preprint, available at <https://arxiv.org/abs/2208.02527>.
- [8] R. DAUTRAY and J.L. LIONS. *Mathematical Analysis and Numerical Methods for Science and Technology*, vol. 5. Springer, Berlin, 1992.
- [9] D. DIER and R. ZACHER. *Non-autonomous maximal regularity in Hilbert spaces*. J. Evol. Equ. **17** (2017), no. 3, 883–907. doi:[10.1007/s00028-016-0343-5](https://doi.org/10.1007/s00028-016-0343-5)
- [10] K. DISSER, A. TER ELST, and J. REHBERG. *On maximal parabolic regularity for non-autonomous parabolic operators*. Journal of Differential Equations **262** (2017), 2039–2072. doi:[10.1016/j.jde.2016.10.033](https://doi.org/10.1016/j.jde.2016.10.033).
- [11] S. FACKLER. *J.-L. Lions’ problem concerning maximal regularity of equations governed by non-autonomous forms*. Ann. Inst. H. Poincaré C Anal. Non Linéaire **34** (2017), no. 3, 699–709. doi:[10.1016/j.anihpc.2016.05.001](https://doi.org/10.1016/j.anihpc.2016.05.001)
- [12] S. FACKLER. *Nonautonomous maximal  $L^p$ -regularity under fractional Sobolev regularity in time*. Anal. PDE **11** (2018), no. 5, 1143–1169. doi:[10.2140/apde.2018.11.1143](https://doi.org/10.2140/apde.2018.11.1143)
- [13] S. GUERRE-DELABRIÈRE.  *$L_p$ -regularity of the Cauchy problem and the geometry of Banach spaces*. Illinois J. Math. **39** (1995), no. 4, 556–566.
- [14] B. HAAK and E.M. OUHABAZ. *Maximal regularity for non-autonomous evolution equations*. Math. Ann. **363** (2015), no. 3–4, 1117–1145. doi:[10.1007/s00208-015-1199-7](https://doi.org/10.1007/s00208-015-1199-7)
- [15] T. HYTÖNEN, J. VAN NEERVEN, M. VERAAR, and L. WEIS. *Analysis in Banach spaces. Vol. III. Harmonic Analysis and Spectral Theory*. Ergebnisse der Mathematik und ihrer Grenzgebiete. Springer.
- [16] N. KRYLOV. *Parabolic and elliptic equations with VMO coefficients*. Comm. Partial Differential Equations **32** (2007), no. 1–3, 453–475. doi:[10.1080/03605300600781626](https://doi.org/10.1080/03605300600781626)
- [17] N. V. Krylov. *On parabolic equations in one space dimension*. Commun. Partial Differ. Equations, 41(4):644–664, 2016.
- [18] J.L. LIONS. *Équations différentielles opérationnelles et problèmes aux limites*. Die Grundlehren der mathematischen Wissenschaften, Bd. 111. Springer-Verlag, Berlin-Göttingen-Heidelberg, 1961.
- [19] C. MOONEY. *Singularities of complex-valued solutions to linear parabolic equations*. Int. Math. Res. Not. IMRN (2021), no. 6, 4413–4426. doi:[10.1093/imrn/rnaa117](https://doi.org/10.1093/imrn/rnaa117)
- [20] J. PRÜSS and R. SCHNAUBELT. *Solvability and maximal regularity of parabolic evolution equations with coefficients continuous in time*. J. Math. Anal. Appl. **256** (2001), no. 2, 405–430. doi:[10.1006/jmaa.2000.7247](https://doi.org/10.1006/jmaa.2000.7247)
- [21] L. DE SIMON. *Un’applicazione della teoria degli integrali singolari allo studio delle equazioni differenziali lineari astratte del primo ordine*. Rend. Sem. Mat. Univ. Padova **34** (1964), 205–223.
- [22] I. SNEIBERG. *Spectral properties of linear operators in interpolation families of Banach spaces*. Mat. Issled. **9** (1974), no. 2, 214–229, 254–255.

DELFT INSTITUTE OF APPLIED MATHEMATICS, DELFT UNIVERSITY OF TECHNOLOGY, P.O. BOX 5031, 2600 GA DELFT, THE NETHERLANDS  
*Email address:* [S.Bechtel@tudelft.nl](mailto:S.Bechtel@tudelft.nl)

DEPARTMENT OF MATHEMATICS, UC IRVINE ROWLAND HALL 410C, IRVINE, CA 92697-3875, USA  
*Email address:* [mooneycr@math.uci.edu](mailto:mooneycr@math.uci.edu)

DELFT INSTITUTE OF APPLIED MATHEMATICS, DELFT UNIVERSITY OF TECHNOLOGY, P.O. BOX 5031, 2600 GA DELFT, THE NETHERLANDS  
*Email address:* [M.C.Veraar@tudelft.nl](mailto:M.C.Veraar@tudelft.nl)