

Problem 1. (§2.3: 10)

Spherical curves. Let α be a unit-speed curve with $\kappa > 0, \tau \neq 0$.

(a) If α lies on a sphere of center \mathbf{c} and radius r , show that

$$\alpha - \mathbf{c} = -\rho N - \rho'\sigma B,$$

where $\rho = 1/\kappa$ and $\sigma = 1/\tau$. Thus $r^2 = \rho^2 + (\rho'\sigma)^2$.

(b) Conversely, if $r^2 = \rho^2 + (\rho'\sigma)^2$ has constant value r^2 and $\rho' \neq 0$, show that α lies on a sphere of radius r .

Solution: Assume that α is on the sphere of radius r . Then we have $\|\alpha - \mathbf{c}\|^2 = r^2$.

Taking the derivative, we get

$$(\alpha - \mathbf{c}) \cdot \alpha' = (\alpha - \mathbf{c}) \cdot T = 0.$$

Thus we can write $\alpha - \mathbf{c}$ as a linear combination of N and B :

$$\alpha - \mathbf{c} = aN + bB,$$

where a, b are functions of t . Taking the derivative again and using the Frenet formulas, we get

$$T = \alpha' = a'N + aN' + b'B + bB' = a'N + a(-\kappa T + \tau B) + b'B - b\tau N.$$

From the above equation, we get the following equations

$$1 + a\kappa = 0, \quad a' - b\tau = 0, \quad a\tau + b' = 0.$$

We therefore have

$$a = -\rho, \quad b = -\rho'\sigma.$$

Thus

$$\|\alpha - \mathbf{c}\|^2 = \rho^2 + (\rho'\sigma)^2 = r^2.$$

Conversely, if

$$\alpha - \mathbf{c} = -\rho N - \rho'\sigma B,$$

then

$$\|\alpha - \mathbf{c}\|^2 = \rho^2 + (\rho'\sigma)^2 = r^2.$$

so α is on the sphere of radius r . ■

Problem 2. (§2.3: 11)

Let $\beta, \bar{\beta} : I \rightarrow \mathbb{R}^3$ be unit-speed curves with nonvanishing curvature and torsion. If $T = \bar{T}$, then β and $\bar{\beta}$ are parallel (Ex. 10 of Sec. 2). If $B = \bar{B}$, prove that $\bar{\beta}$ is parallel to either β or the curve $s \mapsto -\beta(s)$.

Solution: We consider $(\beta(s) - \bar{\beta}(s))' = T - \bar{T} = 0$. Thus $\beta(s) - \bar{\beta}(s) = \mathbf{c}$ is a constant. Thus β and $\bar{\beta}$ are parallel.

If $B = \bar{B}$, then $B' = \bar{B}'$ and hence

$$-\tau N = -\bar{\tau} \bar{N}.$$

Thus since both N and \bar{N} are unit vectors, we must have

$$N = \pm \bar{N}, \quad \tau = \pm \bar{\tau}.$$

Thus we have

$$-\kappa T + \tau B = N' = \bar{N}' = -\bar{\kappa} \bar{T} + \bar{\tau} \bar{B}.$$

From the above, we conclude that $T = \bar{T}$. Thus β is either parallel to $\bar{\beta}$ or $-\bar{\beta}$. ■

Problem 3. (§2.4: 1)

Express the curvature and torsion of the curve $\alpha(t) = (\cosh t, \sinh t, t)$ in terms of arc length s measured from $t = 0$.

Solution: We have

$$\alpha' = (\sinh t, \cosh t, 1),$$

$$\alpha'' = (\cosh t, \sinh t, 0),$$

$$\alpha''' = (\sinh t, \cosh t, 0).$$

Thus we have

$$\|\alpha'\| = \sqrt{2} \cosh t,$$

$$\alpha' \times \alpha'' = (-\sinh t, \cosh t, -1),$$

$$\|\alpha' \times \alpha''\| = \sqrt{2} \cosh t,$$

$$(\alpha' \times \alpha'') \cdot \alpha''' = 1.$$

We then have

$$\kappa = \tau = \frac{1}{2 \cosh^2 t}.$$

In order to find the arc-length reparametrization, we solve the differential equation

$$s'(t) = \|\alpha'(t)\| = \sqrt{2} \cosh t, \quad s(0) = 0.$$

We then have

$$s(t) = \sqrt{2} \sinh t.$$

Therefore

$$\kappa(s) = \tau(s) = \frac{1}{2 \cosh^2 t} = \frac{1}{2 + s^2}.$$

■

Problem 4. (§2.4: 4)

Show that the curvature of a regular curve in \mathbb{R}^3 is given by

$$\kappa^2 \nu^4 = \|\alpha''\|^2 - (d\nu/dt)^2.$$

Solution: Since $\nu = ds/dt$, we have

$$\frac{d\nu}{dt} = \frac{d}{dt} \sqrt{\alpha' \cdot \alpha'}$$

Problem 5. (§2.4: 5)

If α is a curve with constant speed $c > 0$, show that

$$\begin{aligned} T &= \alpha'/c, & N &= \alpha''/\|\alpha''\|, & B &= \alpha' \times \alpha''/(c\|\alpha''\|), \\ \kappa &= \frac{\|\alpha''\|}{c^2}, & \tau &= \frac{\alpha' \times \alpha'' \cdot \alpha'''}{c^2\|\alpha''\|^2}. \end{aligned}$$

Problem 6. (§2.4: 6)

(a). If α is a cylindrical helix, prove that its unit vector \mathbf{u} (Thm. 4.5) is

$$\mathbf{u} = \frac{\tau}{\sqrt{\kappa^2 + \tau^2}} T + \frac{\kappa}{\sqrt{\kappa^2 + \tau^2}} B$$

and the coefficients here are $\cos \theta$ and $\sin \theta$ (for θ as in Def. 4.5).

(b). Check (a) for the cylindrical helix in Example 4.2 of Chapter 1.

Problem 7. (§2.4: 12)

If $\alpha(t) = (x(t), y(t))$ is a regular curve in \mathbb{R}^2 , show that its plane curvature (Ex. 8 of Sec. 3) is given by

$$\tilde{\kappa} = \frac{\alpha'' \cdot J(\alpha')}{\nu^3} = \frac{x'y'' - x''y'}{((x')^2 + (y')^2)^{3/2}}.$$

Problem 8. (§2.5: 1)

Consider the tangent vector $\mathbf{v} = (1, -1, 2)$ at the point $\mathbf{p} = (1, 3, -1)$. Compute $\nabla_{\mathbf{v}} W$ directly from the definition, where

(a) $W = x^2 U_1 + y U_2$,

(b) $W = x U_1 + x^1 U_2 - z^2 U_3$.

Problem 9. (§2.5: 3)

If W is a vector field with constant length $\|W\|$, prove that for any vector field V , the covariant derivative $\nabla_V W$ is everywhere orthogonal to W .