

SURPRISING GEOMETRIC PHENOMENA IN HIGH-DIMENSIONAL CONVEXITY THEORY

V. Milman

Abstract

We shall describe asymptotic phenomena for high-dimensional objects, which violate intuition based on traditional studies of objects of low dimension. We will illustrate these phenomena with examples which involve convex bodies and homogeneous structures (both continuous and discrete) for which other parameters play the role of dimension. Our aim will be to convey some of the new intuition about high-dimensional structures, that underlies such counterintuitive results. We will also discuss the notion of “isomorphic” geometry as the study of common geometric features of families of, say, convex bodies, and analyze their unexpected asymptotic behavior as the dimension increases to infinity. The underlying methods use different mathematical tools and are useful in a variety of apparently unrelated mathematical areas.

1. Introduction

In this lecture we discuss results which stand between Geometry and Functional Analysis. The theory was built during the last two decades. It considers geometric problems via a Functional Analysis point of view. Consequently, the “isometric” problems and point of view typical for geometry are substituted by “isomorphic” ones. This became possible with the *asymptotic* approach (with respect to dimension increasing to infinity) to the study of high dimensional spaces. The goal of this lecture is to demonstrate, with some examples and results, a new intuition which corresponds to high dimensional spaces and present some discoveries and highlights of this theory. We avoid the technical part of the theory and will emphasize the geometric insight. In fact, we restrict this presentation to one facet of the theory: isomorphic (asymptotic) geometry and related phenomena. We do not discuss (for example) the very important type-cotype theory with the deep probabilistic connections behind it. To fill these and other omissions I recommend the following books: [MSch86], [P86a], [P86b], [P89a], [TJ88] and surveys [M88a], [M92], [L92], [LM93].

Consider a finite dimensional normed space $X = (\mathbb{R}^n, \|\cdot\|)$. Such a space is defined by its unit ball $K_X = \{x \in \mathbb{R}^n, \|x\| \leq 1\}$. Inversely, if K is a convex centrally-symmetric compact body in \mathbb{R}^n , then $X_K = (\mathbb{R}^n, \|\cdot\|_K)$ is the normed space with the unit ball K . Let $|x|$ be the canonical euclidean norm in \mathbb{R}^n , (x, y) be the standard inner product and \mathcal{D} denote the standard euclidean ball, i.e. $\mathcal{D} = K_{(\mathbb{R}^n, |\cdot|)}$.

A Few Preliminary Examples. We start with three observations:

a) ([GrM84], [MP89]) Fix $0 < \delta < \frac{1}{2}$ (say, $\delta = \frac{1}{4}$). Define the *floating body* K_δ of K as the intersection of the half spaces $\{x \in \mathbb{R}^n \mid (x, \theta) \leq m_\delta(\theta)\}$, $\theta \in \mathbb{R}^n$, where $m_\delta(\theta)$ is defined by

$$\text{Vol} \{x \in K \mid (x, \theta) > m_\delta(\theta)\} = \delta \text{Vol} K .$$

Then, *there is a number $C(\delta)$, independent of dimension n or $K \subset \mathbb{R}^n$, such that for any symmetric convex body K the floating body K_δ is uniformly, up to a factor $C(\delta)$, isomorphic to an ellipsoid; this means that there is an ellipsoid \mathcal{E} , s.t.*

$$\mathcal{E} \subset K_\delta \subset C(\delta)\mathcal{E} .$$

[Note that the initial body K could be very far from any ellipsoid; it could be, say, a cross-polytope (= the unit ball of ℓ_1^n) or a cube, but the described above “regularization” obtained by cutting a fixed portion of volume in any direction brings us to a $C(\delta)$ -neighborhood of an ellipsoid.]

Moreover, this ellipsoid is homothetic to the Legendre ellipsoid of inertia $\mathcal{L}(K)$ of K .

Let us formally introduce a multiplicative geometric distance $d(\cdot, \cdot)$ between convex bodies K and T : $d(K, T) = \inf\{b/a \mid K \subset bT \text{ and } aT \subset K\}$. The Banach-Mazur distance between two normed spaces X_K and X_T is $d(X_K, X_T) = \inf\{d(K, uT) \mid u \in GL_n\}$.

b) *Centroid body* (of Petty [Pet61]) (or “zonoid of inertia” by [MP89]). Consider a body $Z(K) = \frac{1}{|K|} \int_K [0, x] dx$ which is a “Minkowski” integral, meaning that approximating the integral finite sums are Minkowski sums of sets: intervals $[0, x_i]$ with suitable weights. Geometrically, $\partial Z(K)$ is the collection of centroid points (centers of gravity) of “halves” of our body K obtained by cutting K by hyperplanes passing through the origin.

Again, as in a), there is a universal constant $C > 0$, such that $\forall n, \forall K \subset \mathbb{R}^n, d(Z(K), \mathcal{L}(K)) \leq C$ (and $C \leq \sqrt{8}$).

c) *Lattice tiling.* Let the inner part $\overset{\circ}{K} \neq \emptyset$. Denote $K_i = K + x_i$ for $x_i \in \mathbb{R}^n$. We call $\{K_i\}$ a tiling of \mathbb{R}^n (and say that K produces a tiling by shifts) if (i) $\bigcup_i K_i = \mathbb{R}^n$ and (ii) $\overset{\circ}{K}_i \cap \overset{\circ}{K}_j = \emptyset$ for $i \neq j$. A lattice tiling is a tiling such that the set $\{x_i\}$ is a lattice, i.e. $\{x_i\} = AZ^n$, where A is an invertible linear map and $\mathbb{Z}^n \subset \mathbb{R}^n$ is the set of all integer vectors of \mathbb{R}^n .

Trivially, an affine image of the cube $[-1, 1]^n$ produces a lattice tiling and the euclidean ball does not. However, do uniformly isomorphic versions K_n of the euclidean balls \mathcal{D}_n exist which produce a tiling? Surprisingly, the answer is “Yes”. Exactly, *for any integer n there is a convex symmetric body $K_n \subset \mathbb{R}^n, \mathcal{D}_n \subset K_n \subset 3 \cdot \mathcal{D}_n$, such that K_n produces a lattice tiling of \mathbb{R}^n .* (This observation of Alon-Milman follows immediately from some known results of [R50], [Bu72], [Bou87], [Ban90], on lattice covering-packing; see [M92] for details and references.)

Isomorphic Geometry. The three examples above lead us to a notion of “isomorphic ellipsoid”: a family of convex bodies $\{K_\alpha\}$ of infinitely increasing dimension represents an “isomorphic ellipsoid” if there is a constant C and a family of ellipsoids $\{\mathcal{E}_\alpha\}$ such that $\mathcal{E}_\alpha \subset K_\alpha \subset C\mathcal{E}_\alpha$ for every K_α in the family, i.e. $d(K_\alpha, \mathcal{E}_\alpha) \leq C$. So, in example a) we can state that for a fixed δ , $0 < \delta < \frac{1}{2}$, the family of δ -floating bodies $\{K_\delta \mid \forall n \forall K \subset \mathbb{R}^n\}$ is an isomorphic ellipsoid. (We may add: $C(\delta)$ -isomorphic ellipsoid).

So, we naturally derive *isomorphic* geometric objects and *isomorphic* geometric results. Of course, an “isomorphic” geometric object is, in fact, a family of objects in different spaces of increasing dimension and by “isomorphic” geometric properties of such an “isomorphic” object we mean a common property of this family. Classical Geometry (in a fixed dimension) is an isometric theory (or, at most, “almost” isometric, ε -isometric for small $\varepsilon > 0$). “Isomorphic” study means asymptotic behavior determined by some parameter (most often it is dimension $n \rightarrow \infty$) and exact control of dependence of the constants involved on this parameter (say, dimension in most examples in this lecture).

The appearance of such an isomorphic geometric object is a new feature of asymptotic high-dimensional theory. Geometry and Analysis meet here in a non-trivial way. We will meet later *geometric inequalities in isomorphic form*. The study of special examples of such inequalities, “isomorphic isoperimetric inequalities” led to the discovery of the “Concentration Phenomenon” – one of the most powerful tools of asymptotic theory, responsible for many counter-intuitive results in the theory.

2. Another Example – Theorem; Correction of Intuition

Our next example is already a non-trivial theorem originally proved in 1988 (see [M91]).

Theorem 2.1. *There is a universal constant C so that for any n and any centrally-symmetric, compact, convex body $K \subset \mathbb{R}^n$, there are two linear operators $u_1, u_2 \in SL_n$ so that if $P = K \cap u_1K$ and $Q = \text{conv}(P \cup u_2P)$, then the body Q is within distance C of an ellipsoid: that is, there is an ellipsoid \mathcal{E} for which $\mathcal{E} \subset Q \subset C\mathcal{E}$.*

In the language of Functional Analysis, the same fact can be reformulated in the following form:

Theorem 2.2. *For every finite dimensional normed space $X = (\mathbb{R}^n, \|\cdot\|)$ there are three linear operators $T_1, T_2, T_3 \in GL_n$, such that the following relation holds:*

Consider $p(x) = \|T_1x\| + \|T_2x\|$ and an inf-convolution

$$q(x) = p(x) * p(T_3x) \quad [\text{by definition } p(x) * p(ux) \equiv \inf_{y+z=x} \{p(y) + p(uz)\}];$$

then $q(x)$ is C -isomorphic to the standard euclidean norm in \mathbb{R}^n :

$$|x| \leq q(x) \leq C|x| .$$


Note that C does not depend on the dimension n or on the initial norm $\|\cdot\|$.

In fact, an even stronger statement is correct:

Theorem 2.3. *For any four centrally-symmetric convex bodies $K_i \subset \mathbb{R}^n$, $i = 1, 2, 3, 4$, $|K_i| = |\mathcal{D}|$, there are $\{u_i\}_1^4 \subset \text{SL}_n$ such that if $P_1 = u_1K_1 \cap u_2K_2$, $P_2 = u_3K_3 \cap u_4K_4$ and $Q = \text{Conv } P_1 \cup P_2$, then $\frac{1}{C}\mathcal{D} \subset Q \subset C\mathcal{D}$ for some universal constant C independent of the dimension n and $\{K_i\}_1^4$.*

There are many remarkable ellipsoids associated with a given body K . They are recovering different traces of hidden symmetries which exist in any high-dimensional convex body. If K has a “large” group of symmetries then all these ellipsoids coincide. Well known, for example, are the maximal volume ellipsoid inscribed in K (introduced by Löwner and studied by F. John), the minimal volume ellipsoid circumscribed around K or the Legendre and Binet ellipsoids of inertia. There are many other important constructions of ellipsoids associated with K and an ellipsoid which Theorem 2.1 provides is one of them. We call it the M -ellipsoid and know how to identify it a priori through some geometric inequality. Changing coordinates we may consider this ellipsoid \mathcal{E} to be the standard euclidean ball $\mathcal{D} \subset \mathbb{R}^n$; let $w \in \text{GL}_n$ be such that $w\mathcal{E} = \mathcal{D}$. We call the family of *position* of K the family $\{vK\}_{v \in \text{GL}_n}$. So the affine image $wK = \widehat{K}$ is a *position* of K . Applying Theorem 2.1 to K in such a position, i.e. to \widehat{K} , we may additionally claim that $(u_1; u_2)$ are orthogonal pairs of operators and “most” (meaning with high probability with respect to the Haar probability measure of orthogonal groups) pairs $(u_1; u_2) \in O(n) \times O(n)$ will regularize \widehat{K} to isomorphic euclidean ball \mathcal{D} .

Clearly, every position of K produces the unit ball of isometrically the same normed space as X_K . It is an interesting feature of an (asymptotic) high dimensional theory of convex sets which we will meet throughout the lecture that we are, in fact, forced to consider the family of all positions of a given K (that is all affine images uK , $u \in \text{GL}_n$) even when we are aiming at some volume inequalities or other properties of an individual K . (See K. Ball [B91] for the use of the family of positions in isoperimetric type problems.)




Let us try to develop an intuition of high-dimensional spaces. To start with, we should understand how to draw “high-dimensional” pictures, how high-dimensional convex bodies “look”.

The first non-intuitive point is that 2 or 3 dimensional “pictures” of a high-dimensional convex body should have a “hyperbolic” form! The reason is that the volume of parallel hypersections decays exponentially after passing the median level.

This is a *fact* observed in [GrM84]. Yes, K is a convex set but the rate of volume decay has a crucial influence on the geometry and we should find a way to visualize it in our pictures.

Now, with such a picture in mind, let us try, intuitively, to understand the theorem.



We use the following notation: $|K|$ defines the volume of K and $K^\circ = \{x \in \mathbb{R}^n \mid (x, y) \leq 1 \text{ for every } y \in K\}$ is the polar body of K .

Let $|K| = |\mathcal{D}|$; for some $R > 1$ almost no $(n - 1)$ -dimensional volume belongs to the intersection $R \cdot \partial \mathcal{D} \cap K$. Then find a turn u_1 s.t.

$$(R\partial \mathcal{D} \cap K) \cap (R\partial \mathcal{D} \cap u_1 K) = \emptyset .$$

So $P = K \cap u_1 K \subset R \cdot D$.

But K is convex (finally, we recall this fact), so consider $P^\circ \supset \frac{1}{R}D$. Do the same operation with the body P° : $Q^\circ = P^\circ \cap u_2 P^\circ \subset RD$ and it looks as though we succeeded:

$$\frac{1}{R}D \subset Q \subset R \cdot D .$$

However: When does exponential decay start? What is the value of R ? For many classical ellipsoids (say, maximal vol. ellipsoids), R may be as large as $\sim \sqrt{n / \log n}$. For some other ellipsoids $R \sim \log n$. It was necessary to construct a special ellipsoid (better to say, to prove its existence) for which $R \sim \text{Const}$. However, this is not yet enough. The result above is a “metric one”: we want $(K \cap u_1 K \cap \partial(R \cdot D))$ to be empty, not just to have small volume; so more corrections are necessary, but the idea is correct. Just instead of estimating decay of measure, we should consider decay of coverings.

3. Isomorphic Geometric Inequalities

Let us return to the ellipsoid which appeared in Theorem 2.1. It was originally constructed in connection with some geometric inequality (but, again, inequality in isomorphic form).

Theorem 3.1. *There exists a universal constant C , such that for any convex symmetric body K in \mathbb{R}^n , there is an ellipsoid \mathcal{M}_K , $|\mathcal{M}_K| = |K|$, such that for any (not only convex) set T*

$$(i) \quad \frac{1}{C} |\mathcal{M}_K + T|^{1/n} \leq |K + T|^{1/n} \leq C |\mathcal{M}_K + T|^{1/n}$$

where $K + T = \{x + y \mid x \in K, y \in T\}$ is the Minkowski addition of sets. (We ignore measurability problems; they are not crucial here). And for any convex symmetric set T

$$(ii) \quad \frac{1}{C} |\mathcal{M}_K \cap T|^{1/n} \leq |K \cap T|^{1/n} \leq C |\mathcal{M}_K \cap T|^{1/n} .$$

Define *volume radius* of K by $\text{vrad } K = (|K|/|\mathcal{D}|)^{1/n}$. Then Theorem 3.1 states that substituting K by some special ellipsoid \mathcal{M}_K in (i) and (ii) essentially preserves volume radii.

Of course, it immediately implies reversion of Brunn-Minkowski inequality (which was the initial result in this direction). Recall that Brunn-Minkowski inequality states that for any two compact sets (say, convex) K and T in \mathbb{R}^n ,

$$|K + T|^{1/n} \geq |K|^{1/n} + |T|^{1/n} .$$

The reverse Brunn-Minkowski inequality is the following statement:

Theorem 3.2 [M86]. $\exists C$ such that for any integer n and any two convex bodies K and T there is a relative position $\widehat{K} = uK$, $u \in SL_n$, of say, K such that

$$|\widehat{K} + T|^{1/n} \leq C (|K|^{1/n} + |T|^{1/n}) .$$

(Also the same inequality holds for the polars $\widehat{K}^\circ, T^\circ$).

This is a very easy consequence of Theorem 3.1(i) (but it precedes it).

The original proof of Theorem 3.2 was followed by three other proofs: two by G. Pisier ([P89a], [P89b]) and another one of [M88b]. All four proofs being different in their technical presentations added a number of facets to the understanding of the phenomenon behind Theorems 3.1 and 3.2. One such addition is the following *covering property*.

Let $N(K, T) = \min\{N \mid \exists \{x_i\}_1^N \subset \mathbb{R}^n, K \subset \bigcup_1^N (x_i + T)\}$.

An equivalent statement for Theorem 3.2 (or Theorem 3.1) is the following fact: *There is a number C such that for any $n \in \mathbb{N}$, any K (convex symmetric) there is an ellipsoid \mathcal{M}_K , $|K| = |\mathcal{M}_K|$, such that*

$$N(K, \mathcal{M}_K) \leq e^{Cn} \quad \text{and} \quad N(\mathcal{M}_K, K) \leq e^{Cn} .$$

This is, of course, the best possible estimate (up to a value of universal constant C). Note that, for any ellipsoid \mathcal{E} , $N(2\mathcal{E}, \mathcal{E}) \sim C_0^n$ (for some $C_0 > 1$) but K may be very far from any ellipsoid.

Of course, the ellipsoid described above is not unique and there are many with different additional properties.

Such additional information is a decay of the functions $N(K, t\mathcal{M}_K)$ and $N(\mathcal{M}_K, tK)$ when t increases. Pisier [P89a] showed that, for any $p < 2$ there is an M -ellipsoid \mathcal{E} such that $N(\mathcal{E}, tK) + N(K, t\mathcal{E}) \leq e^{C_p n/t^p}$ for some constant C_p depending on $p < 2$ only.

Remarkably, in fact, ellipsoids do not play a special role in these facts. We may take *any* convex symmetric body P and the family of its positions $\mathcal{P} = \{uP\}_{u \in GL_n}$. Then the same facts are true for \mathcal{P} instead of the family of ellipsoids. Say:

$\exists C$ such that for any convex body K , there is a position $P_K \in \mathcal{P}$ such that $|K| = |P_K|$ and $N(P_K, K) + N(K, P_K) \leq e^{Cn}$.

Also P_K can be put in inequalities (i) and (ii) of Theorem 3.1 instead of \mathcal{M}_K . So, we have some (high-dimensional) *Principle*:

*The affine family of **any** fixed convex symmetric body (say P) is rich enough to substitute any other body (K above) in an essential part of volume computations.*

This reflects a new intuition about high dimension: instead of the expected increasing of essentially different bodies in \mathbb{R}^n with increasing $n \rightarrow \infty$ (diversity of possibilities) we observe their *decreasing* to essentially one (any) body with its affine class.

In many results we described till now, convexity is not a crucial assumption (but *high* dimension is). One may substitute convexity by another, very weak condition, and Theorems 2.1, 2.2 and 3.1, 3.2 still hold. Define a convolution of two bodies $K \square_T = \bigcup_{\substack{x \in K \\ y \in T}} [x, y]$, were $[x, y]$ is the interval joining points x and y .

Let K be a *quasi-convex* star body, i.e.

- (i) $tK \subset K$, $0 \leq t \leq 1$ (star-body condition)
- (ii) $K \square K \subset C \cdot K$

(We say that K is C -quasi-convex. Example: $K_{\ell_p^n}$, the unit ball of ℓ_p^n space for $0 < p < 1$, is a $C(p)$ -quasi-convex for some constant $C(p)$ independent of dimension n .) Clearly, $C = 1$ iff K is convex.

Then Theorems 3.1 and 3.2 are true for quasi-convex bodies with constant C in these theorems depending on the quasi-convexity constant *only* ([BBP95]). Using this fact, also Theorems 2.1 and 2.2 are extended to a quasi-convex

setting ([M95]). I repeat one statement on the language of functions because it corresponds better to the original goal: the study of functions on the sphere S^n .

Let $f \in C(S^{n-1})$, $f > 0$ and even. Consider its homogeneous extension to \mathbb{R}^n by $\widehat{f}(x) = |x|f(\frac{x}{|x|})$, for $x \neq 0$. Let $K = \{x \in \mathbb{R}^n \mid \widehat{f}(x) \leq 1\}$ be a C -quasi-convex body. Apply Theorem 2.2 to the function $\widehat{f}(x) \equiv \|x\|$ (it is not a norm now but a *quasi-norm*). Then the function $q(x)$ we obtained is C_1 -isomorphic to the standard euclidean norm in \mathbb{R}^n where C_1 depends *only* on the C -quasi-convexity constant of K .

As we will see below, these facts reflect the probabilistic nature of high dimension. And I mean by this more than just the fact that we are using probabilistic techniques in many steps of the proofs. In the next section we will discuss the so called *concentration phenomenon* but here I would like to mention one very recent result which gives a probabilistic description for the norm of an operator

$$A : \ell_2^n \rightarrow X .$$

Of course, the norm $\|A\| = \sup\{\|Ax\| \mid |x| \leq 1\}$ is not a probabilistic notion. However, up to a universal factor, i.e. in isomorphic form, it is (!) (see [MSch96]):

there is a universal constant C such that

$$\frac{1}{C} \sqrt{n/k} M(A) \leq \|A\| \leq C \sqrt{n/k} M(A)$$

where $M(A) = \int_{S^{n-1}} \|Ax\| d\mu(x)$, $\mu(x)$ is the rotation invariant probability measure on the sphere $S^{n-1} = \{x \mid |x| = 1\}$, and $k = k(A)$ has also a probability meaning:

$$k = k(A) = \max \left[k \in \mathbb{N} \mid \text{Prob}\{E \in G_{n,k} \mid \|Ax\| \leq 2M(A)|x| \right. \\ \left. \text{for } x \in E\} \geq 1 - \frac{k}{n+k} \right] .$$

Here $G_{n,k}$ is the Grassmann manifold of k -dimensional subspaces of \mathbb{R}^n and the probability measure on $G_{n,k}$ is defined by the euclidean structure ℓ_2^n on \mathbb{R}^n .

4. Isoperimetric Inequalities in Isomorphic form; Concentration Phenomenon

One of the major tools in the development of asymptotic high dimensional theory are isomorphic forms of isoperimetric inequalities. Let us introduce a general framework for such problems, and then let us describe a few examples to give the feeling of what we mean by *isomorphic* form of inequalities.

Let (X, ρ, μ) be a metric space (ρ is a metric) with a probability measure μ . For $A \subset X$, let $A_\varepsilon = \{x \in X \mid \rho(x, A) \leq \varepsilon\}$. Define

$$\alpha(X; \varepsilon) = \sup \left\{ \mu(X \setminus A_\varepsilon) \mid \forall A, \mu(A) \geq \frac{1}{2} \right\} .$$

So, the value $\alpha(\varepsilon)$ corresponds to a solution of the “ ε -isoperimetric problem” for sets of $\frac{1}{2}$ -measure.

Only in very few cases, solutions can be described explicitly and $\alpha(X, \varepsilon)$ calculated exactly. The case of a euclidean sphere S^n – solved by P. Lévy [L51] in 1919 is such a lucky case and the most classical one after \mathbb{R}^n . The calculation gives an estimate $\alpha(S^{n+1}, \varepsilon) \leq \sqrt{\frac{\pi}{8}} e^{-\varepsilon^2 n/2}$. We consider here $S^n \subset \mathbb{R}^{n+1}$ equipped with geodesic distance ρ_r and rotation invariant probability measure μ .

This means that for (any) $A_n \subset S^{n+1}$, $\mu(A_n) = \frac{1}{2}$, and $\varepsilon > 0$, $\mu((A_n)_\varepsilon)$ tends to 1 when $n \rightarrow \infty$ (and exponentially quickly). This is not a geometric point of view. Classical geometry fixes a space (meaning “ n ”) and considers the dependence on $\varepsilon > 0$ and then it does not observe the phenomenon! However, we do not consider one set “ A ” but a family of sets $\{A_n \subset S^{n+1}, \mu(A_n) \geq \frac{1}{2}\}$ and a statement which is an isomorphic geometric inequality is a uniform bound for ε -extensions of sets of this family.

At the same time in many (most) interesting cases, we know how to estimate $\alpha(X; \varepsilon)$ without any idea of the structure of the extremal sets. The above estimate for S^n happens to be typical and implies the so called “concentration phenomenon”. Typical estimates for many natural families $\{X_n\}$ have a similar form:

$$\alpha(X_n, \varepsilon) \leq c_1 e^{-c_2 \varepsilon^2 n} . \quad (*)$$

However, the exact structure of extremal sets or exact value of $\alpha(X; \varepsilon)$ are not known even for such “simple” cases as the torus $X = \mathbb{T}^n = \prod_i^n S_i^1$ equipped with Haar probability measure and distance $d(x, y) = \frac{1}{\sqrt{n}} \sqrt{\sum_1^n \rho(x_i, y_i)^2}$ where ρ is the geodesic distance on S^1 and $x = (x_i)_{i=1}^n, y = (y_i)_{i=1}^n$, or Stiefel manifold $W_{n,2}$ – the unit tangent bundle of S^{n-1} . (Both these examples satisfy (*).)

Another important geometric example (where no extremal sets are known, as well as the exact value of $\alpha(\varepsilon)$) ([GrM83]):

$$\alpha(\text{SO}_n, \varepsilon) \leq \sqrt{\frac{\pi}{8}} e^{-\varepsilon^2 n/8} .$$

Such geometric estimates, asymptotic by increasing “dimension” n , form one of the major technical tools of Asymptotic Theory, and the isomorphic form of the inequalities is enough for all applications [and also the only one typically available]. So the isomorphic (=asymptotic) view on isoperimetric problems has freed us from the necessity of solving (exactly) the isoperimetric problems.

[It is an extremely well developed direction which uses different methods and with many people involved; besides those mentioned above there are Alon, Amir, Borell, Ledoux, Marton, Maurey, Schechtman, Sudakov, Talagrand, Tsirelson and others. (See the books [MSch86], [LT91] and surveys [M88a] and [T95] for details.)]

Let us outline why a bound of the form (*) is so crucial. Consider a 1-Lip function $f(x)$ defined on (X, ρ, μ) , i.e. $|f(x) - f(y)| \leq \rho(x, y)$.

Denote by L_f being the median of $f(x)$, i.e. $\mu\{x \in X \mid f(x) \geq L_f\} \geq \frac{1}{2}$ and $\mu\{x \in X \mid f(x) \leq L_f\} \geq \frac{1}{2}$. Then

$$\mu\{x \in X \mid |f(x) - L_f| < \varepsilon\} \geq 1 - 2\alpha(X, \varepsilon).$$

So, if the value of $\alpha(X, \varepsilon)$ is very small, then values of Lipschitz function “concentrate” by a measure around one value. This is the case when $X = S^n$ and dimension n is large, as well as for \mathbb{T}^n or SO_n for large n . It is, in fact, a general property of high-dimensional metric probability spaces which we call “concentration phenomenon”. Many different techniques were developed to treat different examples of metric probability spaces and to prove concentration phenomenon for them. Let us describe one discrete example considered by B. Maurey [Mau79].

Let Π_n be the group of permutations of $[1, \dots, n]$ equipped with the counting probability measure $\mu(A) = \frac{\#A}{n!}$ (for any $A \subset \Pi_n$) and Hamming distance $\rho(s, t) = \frac{\#\{i \in [1, \dots, n] \mid s(i) \neq t(i)\}}{n}$ for any two permutations $s, t \subset \Pi_n$. Then $\alpha(\Pi_n, \varepsilon) \leq c_1 e^{-c_2 \varepsilon^2 n}$ for absolute constants c_1 and $c_2 > 0$. Again, we have an inequality of “isoperimetric type” but in a new, isomorphic version, which is enough for what we call “concentration phenomenon”.

This phenomenon is responsible for many “unexpected”, “strange” properties of high dimensional spaces. Behind intuitively “expected” properties of high dimensional spaces stands the behavior of ε -entropy which increases exponentially with increase in dimension. However, this exponential increase is compensated by the exponential effect of the concentration phenomenon. As a result, we often observe only linear behavior where a priori intuition expects an exponential. Some Examples:

Approximation by Minkowski sums. Let $A + B = \{x + y \mid x \in A, y \in B\}$ be the Minkowski sum of two sets A and B in \mathbb{R}^n . Let $I_i = [-x_i, x_i] \subset \mathbb{R}^n$ be intervals of length, say, 1. Consider $T = \sum_{i=1}^N I_i$. We want to approximate a Euclidean ball by such sums, that is, for a given $\varepsilon > 0$ we would like to have $d(T, D) \leq 1 + \varepsilon$. Obviously, if $N = n$ then $d(T, D) \geq \sqrt{n}$ and, by an entropy consideration, it looks as if we need at least an exponential by n number of intervals to achieve a good approximation to D . However, an easy geometric interpretation of an old result from [FLM77] (see also [Go85], [Sch88]) shows that there exist intervals $I_i \subset \mathbb{R}^n$, $i = 1, \dots, N_0$, for $N_0 \leq c \frac{n}{\varepsilon^2}$ (c is a numerical constant) such that $d(\sum_{i=1}^{N_0} I_i, D) \leq 1 + \varepsilon$.

This direction was later treated intensively as part of a program of estimating the number of intervals needed for approximating zonoids by a sum of intervals (in [Sch87], [BLM89a] and [BLM88]). It is shown there that the above situation is essentially preserved when we substitute intervals by other convex bodies or if we consider approximation by sums of other convex bodies instead of D . For example [BLM88], [Schm91].

Theorem 4.1. *Let a convex compact body $K \subset \mathbb{R}^n$ be given. There exist orthogonal operators $A_i \in SO_n$, $i = 1, \dots, N_0$ for $N_0 < c \frac{n}{\varepsilon^2}$ such that $T = \frac{1}{N_0} \sum_1^{N_0} A_i K$ satisfies $d(T, D) \leq 1 + \varepsilon$ (as usual, c is a numerical constant).*

Classical symmetrizations. Surprisingly very few symmetrizations are necessary to transform any convex body K to ε -neighborhood of a euclidean ball. Just $\frac{1}{2}n \log n + c(\varepsilon)n$ Minkowski symmetrizations will do the job [BLM88].

The situation with Steiner symmetrization is less clear: $\exists c > 0$ s.t. starting with $\forall K \subset \mathbb{R}^n$, $|K| = |\mathcal{D}|$, we derive T , $c\mathcal{D} \leq T \leq (1 + \varepsilon)\mathcal{D}$, after properly chosen $c(\varepsilon)n \log n$ Steiner symmetrizations (note it is again an isomorphic result) [BLM89b]. However, there is resistance for deriving a $(1 + \varepsilon)$ -approximation by a small number of symmetrizations. Some partial results for higher dimensional symmetrizations are due to Tsolomitis [Ts96].

Another application of concentration estimate (*) is the following: again, let $f(x) \in C(X)$ be a Lipschitz function $|f(x_1) - f(x_2)| \leq \rho(x_1, x_2)$. Then $E|f| \equiv \|f\|_{L_1(X)} \sim L_{|f|}$ (median of $|f|$) if $\alpha(X, \varepsilon)$ is very small. (So as we have already observed, f is “almost constant” with high probability.) But also

$$\|f\|_{L_p(X)} \sim L_{|f|} \sim \|f\|_{L_1(X)},$$

and we see that L_p -norms are equivalent for Lip. functions on X with small $\alpha(X; \varepsilon)$.

Brunn-Minkowski inequality implies similar properties for any convex body K ([Bor75], [GrM84], [MP89]):

$$\forall n, \forall K \subset \mathbb{R}^n, |K| = 1 \quad \text{and} \quad 0 < p, q < \infty$$

$$\text{a) } \|f\|_{L_p(K)} \equiv \left(\int_K |f(x)|^p dx \right)^{1/p} \leq c_{p,q} \|f\|_{L_q(K)} \quad \text{for } \forall \text{ linear function } f(x)$$

$$\text{b) } \text{For any } \|\cdot\| \text{ and } 0 < p, q < \infty : \|(\|x\|)\|_{L_p(K)} \leq c_{p,q} \|(\|x\|)\|_{L_q(K)}$$

One side of such inequalities are just Hölder inequalities; so another side is “reverse” Hölder inequalities which we also call Khinchine type inequalities because $K = \left[-\frac{1}{2}, \frac{1}{2}\right]^n$ in a) gives Khinchine inequalities and in b) gives Kahane’s inequalities.

These cases have a probability interpretation (they correspond to independent random variables uniformly distributed on $\left[-\frac{1}{2}, \frac{1}{2}\right]$). However, from our point of view, the role of the “independence” condition, as well as other standard probability type conditions such as, say, “martingale”, is to create a “high dimensional” convex body; but *any* n -dimensional convex body will serve the same purpose. The reason behind applications of these conditions is the high dimension created and not, for example, independence.

So, probability type ideas and approach are combined here with high-dimensional convexity theory. In fact, the starting point for P. Lévy’s study

of isoperimetric inequality for S^n (and its consequence – concentration phenomenon for the sphere S^n) was a geometric interpretation by E. Borel (~ 1914) of the law of large numbers.

Let $C^n = [-1, 1]^n$ be a cube in \mathbb{R}^n with the standard Euclidean distance “dist”. Then $\text{diam } C^n = 2\sqrt{n}$. Consider a linear functional f , $f(x) = \sum_1^n x_i$ (i.e. $\text{Ker } f = (1, \dots, 1)^\perp$). Then $(1/2^n) \text{Vol}\{x \in C^n : \text{dist}(x, \text{Ker } f) \geq \varepsilon\sqrt{n}\} = P\{|\frac{1}{n} \sum_1^n \xi_i| > \varepsilon \mid (\xi_i)_{i=1}^n \text{ are uniformly distributed on } [-1, 1] \text{ independent random variables}\} \leq c \exp(-\frac{\varepsilon^2 n}{2})$. Therefore, “most” of the volume of C^n is concentrated near a “small slice” (relative to the diameter). However, the concentration phenomenon for a cube C^n implies a similar estimate for any 1-Lip. function on C^n and not only for a particular linear functional. So, in a sense, the concentration phenomenon is a “non-linear” version of deviation-inequality type results in Probability.

Returning to a) and b) equivalences above, Bourgain [Bou91] extended a) to any polynomials (with a constant $c_{p,q}(d)$ depending also on the degree d of the polynomial). An interesting case of a) and b) is $q \rightarrow 0$. It was shown in [MP89] that $c_{p,q}$ are independent of q for linear functionals and recently Latala [La96] showed that constants $c_{p,q}$ are independent of q also in b).

The last example of family of spaces X_n with a concentration property I would like to demonstrate is also discrete. It is taken from [AIM85] and is useful in Computer Science (see also [Al86]).

Let $G = (V, E)$ be a connected graph on $|V| = n$ vertices. We equip V with the counting probability measure $\mu(A \subset V) = |A|/|V|$ and with the path metric: $\rho(x, y) = \{\text{the smallest number of edges in a path which joins } x \text{ and } y\}$.

Let $L^2(V)$ denote the space of real valued functions on V with the usual scalar product (f, g) and the usual norm $\|f\| = \sqrt{(f, f)}$ induced by it. Consider the quadratic form

$$(Qf, f) = \sum_{e \in E} (f(e^+) - f(e^-))^2$$

defined on $f \in L^2(V)$ where e^+ and e^- denote the vertices joined by the edge e . Let $0 = \lambda_0 < \lambda_1 = \lambda_1(G) \leq \lambda_2 \leq \dots \leq \lambda_{n-1}$ be the eigenvalues of Q . Clearly, Q plays a role of Laplacian for the graph $G = (V, E)$. Let $d = \max\{d(v) \mid v \in V\}$ be the maximum degree of a vertex of the graph G . Then the concentration function of (V, ρ, μ) is estimated by

$$\alpha(V; \varepsilon) \leq \frac{1}{2} \exp\left\{-\varepsilon \sqrt{\lambda_1/2d} \log 2\right\}.$$

(Note that ε is an integer in this case and ranges from 1 to the diameter of V .)

The most important examples of the use of this estimate are the so-called Cayley graphs. Let V be a finite group and $S \subset V$ some set of generators. Assume $S = S^{-1}$ and the identity $e \notin S$. We join v and u from V by an edge if $u = s^{-1}v$ for some $s \in S$. Then a path distance in such a graph, $G = (V; E)$, is

the word distance in V induced by S . The degree d of G is equal to $|S|$. Consider $L_2(V)$ and let π be the left regular representation of $V : \pi(t)f(v) = f(t^{-1}v)$. So if $\{\epsilon_t\}_{t \in V}$ is the natural basis of L_2 then $\pi(t)\epsilon_v = \epsilon_{tv}$ and the Laplacian $Q = |A|Id - \sum_{s \in S} \pi(s)$. This is a self-adjoint operator and $\lambda_1(Q) = \{|S| - \text{the second largest positive eigenvalue of } A(G) = \sum_{s \in S} \pi(s)\} \geq |S| - \|A\|_{L_2^0}$ where $L_2^0 \oplus \{\text{const.}\} = L_2(V)$.

To use this construction successfully we have to find natural families of groups V_i , $|V_i| = n_i \rightarrow \infty$, and generators $S_i \subset V_i$, $|S_i| = d_i \leq \text{Const.}$, such that $\|A(G_i)\|_{L_2^0} \leq |S_i| - \varepsilon$ for some positive fixed ε . This brings us to the well-known in Representation Theory T -property of Kazhdan [K67], which gives us a number of most interesting examples, say $V_m = \text{SL}_k(\mathbb{Z}/m\mathbb{Z})$, $k \geq 2$ fixed and $m \rightarrow \infty$.

There is a (right) choice of generators $S_m \subset V_m$, $|S_m| = 4$, such that $\alpha(V_m; t) \leq \frac{1}{2} \exp(-c_k t)$. Note that the bound does not depend on p . It is a very strong concentration property and implies families of expanders – special bipartite graphs which play a central role in the construction of fast algorithms (see, e.g. [Al86], [AlSp92]). I also recommend the books [S90] and [Lu94] which show how to apply delicate results from Number Theory to choose the best possible sets of generators S_m . Many other fascinating related results can be found in these books.

5. Duality and connected isomorphic inequalities

Let K be a convex compact body in \mathbb{R}^n and $K = -K$. Then the *polar* body (or the *dual* body) of K is $K^\circ = \{x \in \mathbb{R}^n \mid (x, y) \leq 1 \text{ for } \forall y \in K\}$. Obviously $X_K^* = X_{K^\circ}$. It is well known how important duality is in the study of convexity and also in Functional Analysis. The notion has its roots, in fact, in mechanics. Legendre and Binet ellipsoids of inertia of K are, up to a factor, the dual of each other (see [J37] or [MP89]). Clearly, a larger body has a smaller polar. It is possible to give (and in many different forms) a precise quantitative form to this relation.

5.1. For example, the product of the volume radii of K and K° remains between two universal constants: there is a universal constant $c > 0$ such that $\forall n$ and $\forall K \subset \mathbb{R}^n$ as above

$$c \leq \left(\frac{|K| \cdot |K^\circ|}{|\mathcal{D}|^2} \right)^{1/n} \leq 1.$$

(The right side is the classical Blaschke-Santaló inequality, and the left side is [BM85].)

These inequalities imply, perhaps more surprisingly, a similar relation for covering numbers ([KM87]): there are universal constants C and $c > 0$ such that $\forall n$ and for every two convex symmetric bodies K and $T \subset \mathbb{R}^n$

$$c \leq \left(\frac{N(K, T)}{N(T^\circ, K^\circ)} \right)^{1/n} \leq C.$$

5.2. The next two results are an interpretation of facts from Local Theory in the language of classical convexity. However, I do not know of any other proofs than the very delicate and non-trivial analysis of the corresponding normed spaces by purely Functional Analysis methods.

5.2a. [FLM77] Let P be a centrally symmetric polytope, $v(P) = \#$ of its vertices and $f(P) = \#$ of its faces. Then $\log v(P) \cdot \log f(P) \geq cn$ for some universal constant $c > 0$ (and for some polytopes it is exact). (Note the importance of the symmetry condition $P = -P$; otherwise, for simplex S ,

$$v(S) = n + 1 \quad \text{and} \quad f(S) = n + 1 ;$$

so exponents are a result of symmetry!)

5.2b. Let $w(K)$ be the mean width of K (again $K = -K$). (It is called $b(K)$ in Convex Geometry; we use the notation $M^*(K)$ in Local Theory; let $\|x\|^*$ be the norm with K° being the unit norm; then $\frac{1}{2}w(K) = M^*(K) = \int_{S^{n-1}} \|x\|^* d\mu(x)$.)

Then \exists a position $uK = \widehat{K}$, $u \in \text{SL}_n$, of K , s.t.

$$w(\widehat{K}) \cdot w(\widehat{K}^\circ) \leq C \log n$$

(or, more strongly, $\leq c \log d_K$ where $d_K = d(X_K, \ell_2^n) = \min\{d(K, \mathcal{E}) \mid \mathcal{E} \text{ is an ellipsoid}\}$).

[This is a combination of results of Lewis [Lew79]/Figiel-Tomczak [FT79]/Pisier [P82].] Note that it is a different type of isomorphic geometric inequality (not a volume-type) and one of the most important tool in the proofs of all isomorphic volume type inequalities we discussed before. It uses solutions of some variational problems, non-trivial duality arguments and, most important, the estimate of Pisier for Rademacher projections which is a non-trivial application of very concrete Harmonic analysis methods in abstract Functional Analysis.

5.3. There is an even deeper connection between sections of K and K° .

5.3a. Fix $\kappa > 0$. Let $k + m \leq [(1 - \kappa)n]$. $\forall K$, for “most” [=high probability] sections E of dimension k and most sections F of K° of dimension m ,

$$w(K \cap E) \cdot w(K^\circ \cap F) \leq \frac{8}{\kappa} .$$

Also, in fact, for some universal C

$$\int_{G_{n,k}} w(K \cap E) d\mu(E) \cdot \int_{G_{n,m}} w(K^\circ \cap F) d\mu(F) \leq \frac{C}{\kappa} .$$

This is surprising: polarity is a *global* construction, so why does it connect so precisely with local structure, i.e. with the structure of sections?

5.3b. In fact, even the maximal width $r(K \cap E) = \text{diam}(K \cap E)$ of a central section E of K of dimension, say, $n/2$, can be estimated in average by the mean-width: for some universal constant C ,

$$\int_{G_{n,n/2}} \text{diam}(K \cap E) d_\mu(E) \leq Cw(K) .$$

In Local Theory, we call this “the low M^* -estimate” (see [M85], [M90], [PT86], [Go88], or any of the books mentioned earlier). In its analytic form it is used as a bridge between results of a Functional Analytic nature and Geometric type Global results.

5.4. Another type of geometric (isomorphic) inequality connecting a body K with its polar K° is the following:

Let $r_\ell = \max\{r \mid r\mathcal{D} \subset \frac{1}{\ell} \sum_{i=1}^{\ell} u_i K, u_i \in O(n)\}$.

Then $\exists c > 0$ such that, $\forall n, \forall K \subset \mathbb{R}^n$, convex symmetric compact body

$$r_2(K) \cdot r_3(K^\circ) \geq c .$$

In fact, in this case we know that $c \geq 0.0388$ (see [M91]).

Applying this fact to, say, the unit cubes $C^n = [-1, 1]^n$, we immediately see that $\exists u \in O(n)$ s.t.

$$c\sqrt{n}\mathcal{D} \subset \frac{C^n + uC^n}{2} \subset \sqrt{n}\mathcal{D}$$

for an absolute constant $c > 0$. This particular very interesting case was derived long ago by Kashin [Kas77].

6. Local and Global Asymptotic Theories

At the heart of the global results presented in this lecture stand methods of Functional Analysis. By global properties we refer to properties of the original body in question, while the local properties pertain to the structure of lower dimensional sections and projections of the body, i.e. to the linear structure of a normed space in the spirit of functional analysis.

From the beginning of the 70's the needs of Geometric Functional analysis led to a deep investigation of the linear structure of finite dimensional normed spaces. Dvoretzky's theorem was the initial result in this direction. But it had to develop a long way before this structure was understood well enough that it could be used for the study of the global properties of a space.

The culmination of this study was an understanding of the fact that subspaces (and quotient spaces) of proportional dimension behave very predictably. This was the bridge between the problems of Functional Analysis and the Global Asymptotic properties of convex sets. The following theorem was one of the first links between local properties and global structure [M85]:

Theorem 6.1. Fix $0 < \lambda < 1$. Every finite dimensional normed space $X = (\mathbb{R}^n, \|\cdot\|)$ contains a subspace sX and a quotient $q(sX) = F$ of a subspace sX such that

$$\text{i) } k = \dim(F) \geq \lambda n \quad \text{ii) } d(F, \ell_2^k) \leq C(\lambda) \sim \frac{1}{1-\lambda} \log \frac{1}{1-\lambda} .$$

We know to-day how to rewrite almost every fact of Local Theory in a Global form. For example, the above theorem is (for $\lambda = \frac{1}{2}$) a “Local form” of the “Global” Theorems 2.1 and 2.2. (λ close to 1 has no global analogue.) Also Theorem 4.1 may be considered a global form of Dvoretzky’s Theorem. However, only recently in one case did we succeed to put this analogy in an exact quantitative form, and I am ending this lecture with this example.

Let $|\cdot|$ define the standard euclidean norm in \mathbb{R}^n , X be \mathbb{R}^n with norm $\|\cdot\|$ and $M = \int_{S^{n-1}} \|x\| d\mu(x)$. Let $k = k(X; |\cdot|) \leq n$ be the largest integer such that

$$\mu_{G_{n,k}} \left(\left\{ E ; \frac{M}{2}|x| \leq \|x\| \leq 2M|x| \text{ for all } x \in E \right\} \right) > 1 - \frac{k}{n+k} .$$

Let t be the smallest integer such that there are orthogonal transformations $u_1, \dots, u_t \in O(n)$ with

$$\frac{M}{2}|x| \leq \frac{1}{t} \sum_{i=1}^t \|u_i x\| \leq 2M|x| , \quad \text{for all } x \in \mathbb{R}^n .$$

So $k = k(X, |\cdot|)$ is a quantitative expression for a Local property of a space X equipped with the given euclidean structure; at the same time, t gives us a quantitative form for the corresponding global property of the space X .

Theorem 6.2 [MSch96]. For some universal constant C

$$C^{-1}n \leq kt \leq Cn .$$

Concluding Remarks.

There are a number of reasons for and origins of order behavior. One may mention “repetition” which creates order, as statistics demonstrates. What I attempted to show in this lecture was that very high dimensions, or more generally, high parametric families are another source of order. Of course, time limitations have restricted me to very specific objects and structures.

Let me note in conclusion a few remarks about the techniques used in this theory, which stand behind the scenes of my lecture. Besides the concentration phenomenon which absorbed a huge diversity of methods, the probabilistic nature of a high dimensional space implies that probability notions and parameters should play an important, perhaps central, role. Indeed, notions of

the type and cotype of a space which were developed in the 70s (through works of Maurey, Pisier, Kwapien, Krivine and others) give a “right” classification of normed spaces (convex bodies) by their probabilistic behaviour. Also Harmonic Analysis techniques enter the abstract theory of Convexity through these notions (Pisier; Bourgain). Let me also note the use of delicate martingale theory, p -stable probability laws (Pisier, Johnson-Schechtman) and the construction of “random normed spaces” (Gluskin; Szarek) with a very interesting behaviour, not observed in concrete examples.

I would like to thank Miriam Hercberg for helping me to edit this text and artfully weaving computer graphics into \TeX . I also thank Noga Alon and Leonid Polterovich for their useful remarks on the first draft of this talk.

References

Books and Surveys.

- [AlSp92] N. Alon, J.H. Spencer, *The Probabilistic Method*, Wiley Interscience, 1992.
- [LT91] M. Ledoux, M. Talagrand, *Probability in Banach Spaces*, *Ergebnisse der Mathematik und ihrer Grenzgebiete* 23, Springer Verlag, 1992.
- [L51] P. Levy, *Problèmes concrets d’analyse fonctionnelle*, Gauthier Villars, Paris 1951.
- [L92] J. Lindenstrauss, Almost spherical sections, their existence and their applications, *Jber. Deutsch. Math.-Vereinig. Jubiläumstagung 1990*, Teubner, Stuttgart, 39-61 (1992).
- [LM93] J. Lindenstrauss, V. Milman, The local theory of normed spaces and its applications to convexity, In “*Handbook of Convex Geometry*”, Vol. B, 1149-1220 (P.M. Gruber and J.M. Wills, eds), North Holland (1993).
- [Lu94] A. Lubotzky, *Discrete Groups, Expanding Graphs and Invariant Measures*, *Progress in Mathematics* 125, Birkhauser Verlag, 1994.
- [M88a] V.D. Milman, The heritage of P. Levy in geometrical functional analysis, *Asterisque* 157/158, 273-301 (1988).
- [M92] V. Milman, Dvoretzky Theorem – 30 years later, *Geometric and Functional Analysis (GAFA)* 2:4, 455-479 (1992).
- [MP89] V.D. Milman, A. Pajor, Isotropic position and inertia ellipsoids and zonoids of the unit ball of a normed n -dimensional space, *GAFA-Seminar 87-88*, *Springer Lecture Notes in Math.* 1376, 64-104, (1989).
- [MSch86] V.D. Milman, G. Schechtman, *Asymptotic theory of finite dimensional normed spaces*, Springer-Verlag, *Lecture Notes in Mathematics* 1200, 156pp. (1986).
- [P86a] G. Pisier, *Factorization of linear operators and geometry of Banach spaces*, *CBMS Regional Conference Series no. 60*, AMS, 1-154 (1986). (Second printing with corrections, 1987).

- [P86b] G. Pisier, Probabilistic methods in the geometry of Banach spaces, CIME, Varenna, 1985, Springer Lecture Notes in Mathematics 1206, 167-241 (1986).
- [P89a] G. Pisier, The Volume of Convex Bodies and Banach Space Geometry, Cambridge Tracts in Math. 94 (1989).
- [S90] P. Sarnak, Some Applications of Modular Forms, Cambridge Tracts in Mathematics 99, Cambridge University Press, 1990.
- [T95] M. Talagrand, Concentration of measure and isoperimetric inequalities in product spaces, IHES, Publications Mathématiques 81, 205pp. (1995).
- [TJ88] N. Tomczak-Jaegermann, Banach-Mazur Distances and Finite Dimensional Operator Ideals, Pitman, 1988.

Articles.

- [Al86] N. Alon, Eigenvalues and expanders, *Combinatorica* 6 (1986), 83-96.
- [AlM85] N. Alon, V.D. Milman, λ_1 , isoperimetric inequalities for graphs and superconcentrators, *J. Combinatorial Theory, Ser. B* 38, N1, 73-88 (1985).
- [AmM80] D. Amir, V.D. Milman, Unconditional and symmetric sets in n -dimensional normed spaces, *Israel J. of Math.*, Vol. 37, 1-2, 3-20 (1980).
- [B91] K. Ball, Volume ratios and a reverse isoperimetric inequality, *J. London Math. Soc.* 44 (1991).
- [Ban90] W. Banaszczyk, On the lattice packing-covering ratio of finite dimensional normed spaces. *Colloq. Math.* 59:1, 31-33 (1990).
- [BBP95] J. Bastero, J. Bernués, A. Peña, An extension of Milman's reverse Brunn-Minkowski inequality, *GAFSA* 5:3, 572-581 (1995).
- [Bob96] S.G. Bobkov, An isoperimetric inequality on the discrete cube, and an elementary proof of the isoperimetric inequality in Gauss space, preprint.
- [BobH96] S.G. Bobkov, C. Houdré, Isoperimetric constants for product probability measures, preprint.
- [Bor75] C. Borell, The Brunn-Minkowski inequality in Gauss spaces, *Inventiones Math.* 30, 207-216 (1975).
- [Bou87] J. Bourgain, On lattice packing of convex symmetric sets in \mathbb{R}^n , Springer Lecture Notes in Math. 1267, 5-12 (1987).
- [Bou89] J. Bourgain, Bounded orthogonal systems and the $\Lambda(p)$ -set problem. *Acta* 162 (1989), 227-245.
- [Bou91] J. Bourgain, On the distribution of polynomials on high dimensional convex sets, Israel Seminar (GAFSA) 1989-90 (J. Lindenstrauss, V.D. Milman, eds.) Springer Lecture Notes in Math. 1469, 127-137 (1991).
- [BLM88] J. Bourgain, J. Lindenstrauss, V. Milman, Minkowski sums and symmetrizations, GAFSA-Seminar Notes '86-87, Springer Lecture Notes in Math. 1317, 44-66 (1988).

- [BLM89a] J. Bourgain, J. Lindenstrauss, V. Milman, Approximation of zonoids by zonotopes, *Acta Math.* 162, 73-141 (1989).
- [BLM89] J. Bourgain, J. Lindenstrauss, V. Milman, Estimates related to Steiner symmetrizations, *Springer Lecture Notes in Math.*, v. 1376, 264-273 (1989).
- [BM85] J. Bourgain, V.D. Milman, Sections euclidiennes et volume des corps symétriques convexes dans R^n , *C.R Acad. Sc. Paris*, t.300 Ser. 1, No. 13, 435-438 (1985) (see also, J. Bourgain, V.D. Milman, New volume ratio properties of convex symmetric bodies in R^n , *Invent. Math.* 88, 319-340 (1987)).
- [Bu72] G.Y. Butler, Simultaneous packing and covering in euclidean space, *Pro. London Math. Soc.* 3:25, 721-735 (1971).
- [FLM77] T. Figiel, J. Lindenstrauss, V.D. Milman, The dimension of almost spherical sections of convex bodies, *Acta Math.* 139:1-2, 53-94 (1977).
- [FT79] T. Figiel, N. Tomczak-Jaegermann, Projections onto Hilbertian subspaces of Banach spaces, *Israel J. Math.* 33, 155-171 (1979).
- [Gl81] E.D. Gluskin, The diameter of the Minkowski compactum is roughly equal to n , *Funct. Anal. Appl.* 15, 72-73 (1981).
- [Go85] Y. Gordon, Some inequalities for Gaussian processes and applications, *Israel J. Math.* 50, 265-289 (1985).
- [Go88] Y. Gordon, On Milman's inequality and random subspaces which escape through a mesh in \mathbb{R}^n , *GAFa*, *Israel Funct. Analysis Seminar (86/87)*, *Springer Lecture Notes* 1317, 84-106 (1988).
- [Gr83] M. Gromov, Filling Riemannian manifolds, *J.D.G.* 18, 1-147 (1983).
- [Gr88] M. Gromov, Dimension, non-linear spectra and width, *GAFa*, *Israel Funct. Analysis Seminar (86/87)*, *Springer Lecture Notes* 1317, 132-184 (1988).
- [GrM83] M. Gromov, V.D. Milman, A topological application of the isoperimetrical inequality, *American J. of Math.* 105:4, 843-854 (1983).
- [GrM84] M. Gromov, V.D. Milman, Brunn Theorem and a concentration of volume of convex bodies, *GAFa Seminar Notes*, Tel Aviv University, Israel 1983-1984, Exp. V, 12pp.
- [IST76] I.A. Ibragimov, V.N. Sudakov, B.S. Tsirelson, Norms of Gaussian sample functions, *Lecture Notes Math.* 550, 20-41 (1976).
- [J37] F. John, Polar correspondence with respect to convex region, *Duke Math. J.* 3:2, 355-369 (1937) (University of Kentucky).
- [JS82] W.B. Johnson, G. Schechtman, Embedding ℓ_p^m into ℓ_1^n , *Acta Math* 149, 71-85 (1982).
- [Kas77] B. Kašin, Sections of some finite dimensional sets and classes of smooth functions, *Izv. Acad. Nauk SSR* 41, 334-351 (1977) (in Russian).
- [K67] D.A. Kazhdan, Connection of the dual space of a group with the structure of its closed subgroups, *Functional Anal. Appl.* 1, 63-65 (1967).

- [KM87] H. König, V.D. Milman, On the covering numbers of convex bodies, GAFA-Seminar Notes, Springer Lecture Notes in Math. 1267, 82-95 (1987).
- [La96] R. Latała, On the equivalence between geometric and arithmetic means for logconcave measures, preprint.
- [Le92] M. Ledoux, A heat semigroup approach to concentration on the sphere and on a compact Riemannian manifold, GAFA 2, 221-224 (1992).
- [Lew79] D. Lewis, Ellipsoids defined by Banach ideal norms, *Mathematika* 26, 18-29 (1979).
- [Ma96] K. Marton, A measure concentration inequality for contracting Markov chains, GAFA 6:3, 556-571 (1996).
- [Mau79] B. Maurey, Construction de suites symétriques, C.R.A.S., Paris 288, 679-681 (1979).
- [M85] V.D. Milman, Almost Euclidean quotient spaces of subspaces of a finite dimensional normed space, *Proceedings AMS* 94:3, 445-449 (1985).
- [M86] V.D. Milman, An inverse form of the Brunn-Minkowski inequality with applications to local theory of normed spaces, C.R. Acad. Sc. Paris 302, Ser. 1, No. 1, 25-28 (1986).
- [M88b] V.D. Milman, Isomorphic symmetrization and geometric inequalities, GAFA-Seminar Notes '86-87, Springer Lecture Notes in Math. 1317, 107-131 (1988).
- [M90] V. Milman, A note on a low M^* -estimate, In "Geometry of Banach Spaces" (eds. P.F.X. Müller and W. Schachermayer), London Mathematical Society Lecture Notes Series 158, 219-229 (1990).
- [M91] V. Milman, Some applications of duality relations, GAFA Seminar 1989-90, Springer Lecture Notes in Math. 1469, 13-40 (1991).
- [M95] V. Milman, Isomorphic Euclidean regularization of quasi-norms in \mathbb{R}^n , C.R. Acad. Sci. Paris, 321, 879-884 (1995).
- [MSch96] V.D. Milman, G. Schechtman, Global vs. local asymptotic theories of finite dimensional normed spaces, preprint 1996.
- [PT86] A. Pajor, N. Tomczak-Jaegermann, Subspaces of small codimension of finite dimensional Banach spaces, *Pro. Amer. Math. Soc.* 97, 637-642 (1986).
- [Pet61] C.M. Petty, Centroid surfaces, *Pacific J. Math.* 11, 1535-1547 (1961).
- [P82] G. Pisier, Holomorphic semi-groups and the geometry of Banach spaces, *Annals of Math.* 115, 375-392 (1982).
- [P89b] G. Pisier, A new approach to several results of V. Milman, *Journal für die Reine und Angew. Math.* 393, 115-131 (1989).
- [R50] C.A. Rogers, A note on coverings and packing, *J. London Math. Soc.* 25, 327-331 (1950).
- [Sch81] G. Schechtman, Levy type inequality for a class of metric spaces, in "Martingale Theory in Harmonic Analysis and Banach Spaces", 211-215, Springer-Verlag, 1981.

- [Sch87] G. Schechtman, More on embedding subspaces of L_p in ℓ_r^n , *Compositio Math.* 61, 159-170 (1987).
- [Sch88] G. Schechtman, A remark concerning the dependence on ε in Dvoretzky's Theorem, *GAFSA Seminar Notes 87-88*, Springer Lecture Notes in Math. 1376, 274-277 (1988).
- [Schm91] M. Schmuckenschlager, On the dependence on ε in a theorem of J. Bourgain, J. Lindenstrauss, V. Milman, *GAFSA Seminar Notes 89-90*, Springer Lecture Notes in Math. 1469, 166-173 (1991).
- [SuTs78] V.N. Sudakov, B. Tsirelson, Extremal properties of half-spaces for spherically invariant measures, *J. Soviet. Math.* 9, 9-18 (1978) [translated from *Zap. Nauch. Sem. L.O.M.I.* 41, 14-24 (1974)].
- [Sz83] S.J. Szarek, The finite dimensional basis problem, with an appendix on nets of Grassman manifold, *Acta Math.* 151, 153-179 (1983).
- [Ts96] A. Tsolomitis, Quantitative Steiner/Schwartz-type symmetrizations, *Geometriae Dedicata* 60:2, 187-206 (1996).

V. MILMAN, Sackler Faculty of Exact Sciences, Tel Aviv University, Tel Aviv 69978, Israel